



Frontiers International Conference
on Wastewater Treatment



PROCEEDINGS
Frontiers International Conference on
Wastewater Treatment (FICWTM)

May 21–24, 2017

Palermo, Italy



Frontiers International Conference
on Wastewater Treatment



PROCEEDINGS
Frontiers International Conference on
Wastewater Treatment (FICWTM)

May 21–24, 2017

Palermo, Italy

**Proceedings of Frontiers International Conference on Wastewater
Treatment
May 21-14, 2017, Palermo, ITALY.**

How to cite the full proceedings:

Mannina, G., 2017. Proceedings of Frontiers International Conference on
Wastewater Treatment
May 21-14, 2017, Palermo, ITALY.

How to cite an individual paper:

Author, A., Author, B., Author, C..., 2016. This is the title of your paper. In:
Mannina, G., 2017. Proceedings of Frontiers International Conference on
Wastewater Treatment
May 21-14, 2017, Palermo, ITALY.

Peer Review:

Each paper has been peer reviewed by at least three independent reviewers
with possible outcomes of reject, revise, and accept.

Bacterial community structure of an IFAS-MBRs wastewater treatment plant

P. Cinà^{*}, G. Bacci^{**}, G. Gallo^{*}, M. Capodici^{***}, A. Cosenza^{***}, D. Di Trapani^{***}, R. Fani^{**}, G. Mannina^{***}, A.M. Puglia

* Laboratory of Molecular Microbiology and Biotechnology, STEBICEF Department, University of Palermo, Viale delle Scienze ed.16, 90128 Palermo, Italy.

**Laboratory of Microbial and Molecular Evolution, Department of Biology, University of Florence, Florence, Italy.

*** Dipartimento di Ingegneria Civile, Ambientale, Aerospaziale, dei Materiali, Università di Palermo, Viale delle Scienze, Ed. 8, 90100, Palermo, Italy.

Summary

The principal objective of domestic or industrial wastewater treatment is generally to make effluents less hazardous to human or environment health. It is well known that nutrients particularly, nitrogen (N) and phosphorus (P) compounds may have adverse environmental impacts (e.g., eutrophication, toxicity towards the aquatic organisms, etc...). Biological nutrient removal (BNR) from domestic wastewater has been extensively investigated in the last years. In an University Cape Town- Moving Bed Membrane BioReactor (UCT-MB-MBR), BNR is usually based on in-series anaerobic, anoxic and aerobic reactors where N and P removal is accomplished by heterotrophic denitrifying bacteria and polyphosphate-accumulating organisms (PAOs). In particular, the biological phosphorous removal is usually achieved through the growth of PAOs, able to accumulate and store P as intracellular polyphosphate (poly-P) under alternating anaerobic/aerobic conditions. Thus, the knowledge of bacterial communities involved in wastewater treatment plant is crucial for a better understanding of the biological processes that allow the nutrient removal and, perspective, for improving plant BNR performance. In this work, the bacterial community putatively involved in BNR events of a UCT-MBMBR pilot plant was elucidated by both culture-dependent and metagenomics DNA analyses. The presence of bacterial isolates belonging to *Bacillus* (in the anoxic compartment) and to *Acinetobacter*, *Stenotrophomonas*, *Rhodococcus*, *Escherichia* and *Aeromonas* (in the aerobic compartment) is in agreement with the nitrification/denitrification processes observed in the plant. Moreover, the study of bacterial community structure by NGS revealed a microbial diversity suggesting a biochemical complexity which can be further explored and exploited to improve UCT-MBMBR plant performance.

Bacterial Communities; NGS; Biological Nutrient Removal; Wastewater treatment plant; Membrane Bioreactors; MBBR; Enhanced biological phosphorus removal; IFAS-MBR

1. Introduction

The principal objective of domestic or industrial wastewater treatment is generally to make effluents less hazardous to human or environment health. It is well known that nutrients (particularly, nitrogen and phosphorus compounds) may have adverse environmental impacts (e.g., eutrophication, toxicity towards the aquatic organisms, etc...) (Wang et al., 2006). Biological nutrient removal (BNR) from domestic wastewater has been extensively investigated and developed in the last years. In systems aimed at the BNR in-series anaerobic, anoxic and aerobic reactors are required (Wanner et al., 1992; Mannina et al., 2016) where nitrogen (N) and phosphorus (P) removal is accomplished by heterotrophic denitrifying bacteria and polyphosphate-accumulating organisms (PAOs), requiring a carbon

source (Naessens et al., 2012). In particular, the biological phosphorus removal is usually achieved through the growth of PAOs, able to accumulate P and to store it as intracellular polyphosphate (poly-P) under alternating anaerobic/aerobic conditions (Li et al., 2013). Thus, knowledge of bacterial communities that take root in wastewater treatment plant is indispensable for a better understanding of the biological processes that allow the nutrient removal and perspective for improving plant BNR performance. In this study bacterial communities of an Integrated Fixed Film Activated Sludge (IFAS) University Cape Town (UCT) membrane bioreactor (MBR) was investigated.

2. Methods

Plant design. The IFAS-UCT-MBR pilot plant was characterized by three in-series reactors: one anaerobic (volume 62 L), one anoxic (volume 102 L) and one aerobic (volume 211 L) compartment according to the UCT scheme.




Figure 2.1 Schematic lay-out of IFAS-UCT-MBR pilot plant

The IFAS-UCT-MBR pilot plant was operated for 60 days and was fed with a mixture of real wastewater (deriving from the University buildings and characterized by higher ammonia content compared to typical domestic wastewater) and synthetic wastewater. In particular, MBBR processes rely on the use of small plastic carriers elements (courtesy of amitec Co.Ltd, carriers density= 0.95 g cm^3) that are kept in constant motion throughout the entire volume of the reactor, for biofilm growth. The plastic carriers were a 15 and 40% filling fraction, corresponding to a net surface area of $75 \text{ and } 200 \text{ m}^2 \text{ m}^{-3}$ in the anoxic and aerobic reactor, respectively.

Identification of bacterial isolates. Wastewater aliquots of $100 \mu\text{L}$ from the aerobic and anoxic tanks were collected, serially diluted and, then, plated on Luria Bertami (LB), Mannitol Soya flour (MS) and R2YE agar-media (Kieser et al., 2000). The plates were incubated at 30°C until appearance of microbial colonies (2-5 days). The bacterial colonies, selected on the basis of pigmentation and morphology, were repeatedly plated on agar-media to obtain pure cultures. The bacterial isolates were characterized on the base of their 16S rDNA sequence using the universal bacterial primers 27F and 1492R (Frank et al., 2008) for 16S rDNA amplification by colony PCR as previously described (Gallo et al. 2012; Milanese et al., 2015). The PCR products were purified by using NucleoSpin Gel and PCR Clean-up (MACHERY-NAGEL, Germany). Sequencing was performed by BMR Genomics srl. Phylogenetic

relationships to known species were inferred by neighbour-joining, using the software Mega6 (Tamura et al., 2013).

Metagenomic analysis based on 16S rDNA sequencing. Metagenomic DNA was extracted from 50 mL aliquots of wastewater from aerobic, anoxic and anaerobic compartment, respectively. In addition, two colonized carriers from aerobic and anoxic compartments were used to obtain Metagenomic DNA. Since CTAB performed better in reducing humic contamination, it was used in the buffer for sodium dodecyl sulfate (SDS)-based DNA extraction. The metagenomic analysis was performed through new generation sequencing (NGS) by Illumina platform (BMR Genomics srl).

3. Results and discussion

After 2–5 days of incubation, bacterial colonies appeared on the surface of agar-medium plates inoculated with serially diluted wastewater aliquots. The colonies showed different phenotypes: most had a white translucent pigmentation, someone white with a matt pigmentation and one showed orange pigmentation. A total of 14 isolates (8 and 6 from aerobic and anoxic tank, respectively) were selected and obtained as pure cultures. From all the 14 isolates, a 16S rDNA sequence was obtained to carry out phylogenetic analysis. As all the isolates from anoxic tank belonged to *Bacillus* genus, the isolates from aerobic tank belonged to five different bacterial genera: *Acinetobacter* (2 isolates), *Stenotrophomonas* (2 isolates), *Rhodococcus* (2 isolates), *Escherichia* (1 isolates) and *Aeromonas* (1 isolates). Interestingly, they are all implied in nitrification/denitrification processes (Di Bonaventura et al., 2004).

The 16S rDNA-based metagenomic analysis revealed 12 (Operational taxonomic unit) OTU. In general, the bacterial community composition of anaerobic, anoxic and aerobic compartments are quite similar; indeed, the most relevant differences are quantitatively observed for the less represented OTUs. In particular, the most representative OTUs were: Saprospiraceae (27.30%), Rhodocyclaceae (26.61%), Sphingobacteriales (12.96%) in aerobic compartment (Fig. 3.1a); Rhodocyclaceae (22.69%), Saprospiraceae (19.99%), and Sphingobacteriales (10.54%) in anoxic compartment (Fig. 3.1b); Rhodocyclaceae (21.60%), Saprospiraceae (19.20%), and Sphingobacteriales (10.29%) in anaerobic compartment (Fig 3.1e). Concerning the colonized carriers, the bacterial community composition almost paralleled that one of the corresponding compartment, with Saprospiraceae and Rhodocyclaceae the most abundant OTUs in both cases. However, the less abundant OTUs *Rhodanobacter*, in aerobic and anoxic conditions, and *Thermomonas* and *Clostridium sensu strictu*, in aerobic condition, were observed only in the carries.




Figure 3.1 Bacterial OTU relative abundance in a) aerobic tank, b) anoxic tank, c) aerobic tank carriers, d) anoxic tank carriers e) anaerobic tank.

4. Conclusions

The bacterial community putatively involved in BNR events of a IFAS-UCT-MBR pilot plant was elucidated by both culture-dependent and metagenomics DNA analyses. The presence of bacterial isolates in the anoxic and aerobic compartments is in agreement with the nitrification/denitrification processes observed in the plant (Mannina et al, unpublished results). Moreover, the study of microbial community structure by NGS revealed a microbial diversity and suggests a biochemical complexity which has to be further explored and exploited to improve IFAS-UCT-MBR plant performance.

Acknowledgments

This work forms part of a research project supported by grant of the Italian Ministry of Education, University and Research (MIUR) through the Research project of national interest PRIN2012 (D.M. 28 dicembre 2012 n. 957/Ric – Prot. 2012PTZAMC) entitled “Energy consumption and GreenHouse Gas (GHG) emissions in the wastewater treatment plants: a decision support system for planning and management – <http://ghgfromwwtp.unipa.it>” in which prof. Giorgio Mannina is the Principal Investigator.

References

- Di Bonaventura, G., Spedicato, G., D'Antonio, D., Robuffo, I., Piccolomini, R. 2004. Biofilm Formation by *Stenotrophomonas maltophilia*: Modulation by Quinolones, Trimethoprim- Sulfamethoxazole, and Ceftazidime antimicrob. Agents chemother. 48(8), 151–160.
- Frank JA, Reich CI, Sharma S, Weisbaum JS, Wilson BA, Olsen GJ (2008) Critical evaluation of two primers commonly used for amplification of bacterial 16S rRNA genes. Appl Environ Microbiol 74:2461–2470



-
- Gallo G, Baldi F, Renzone G, Gallo M, Cordaro A, Scaloni A, Puglia AM (2012) Adaptive biochemical pathways and regulatory networks in *Klebsiella oxytoca* BAS-10 producing a biotechnologically relevant exopolysaccharide during Fe (III)-citrate fermentation. *Microb Cell Fact* 11:152
- Kieser Y, Bibb MJ, Buttner MJ, Chater KF, Hopwood DA (2000) *Practical Streptomyces genetics*. The John Innes Foundation, Norwich
- Li, C., Wang, T., Zheng, N., Zhang, J., Ngo, H.H., Guo, W., Liang, S. 2013. Influence of organic shock loads on the production of N₂O in denitrifying phosphorus removal process. *Bioresour. Technol.* 141, 160–166.
- Mannina, G., Capodici, M., Cosenza, A., Di Trapani, D., 2016. Carbon and nutrient biological removal in a University of Cape Town membrane bioreactor: Analysis of a pilot plant operated under two different C/N ratios. *Chemical Engineering Journal* 296 (2016) 289–299
- Milanesi C, Cresti M, Costantini L, Gallo M, Gallo G, Crognale S, Faleri C, Gradi A, Baldi F (2015) Spoilage of oat bran by sporogenic microorganisms revived from soil buried 4000 years ago in Iranian archaeological site. *Int Biodeter Biodegr* 104:83–91
- Naessens, W., Maere, T., Nopens, I., 2012. Critical review of membrane bioreactor models Part 1: biokinetic and filtration models. *Bioresour. Technol.* 122, 95–106.
- Tamura K, Stecher G, Peterson D, FilipSKI A, Kumar S (2013) MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30:2725–2729
- Wang, X.J., Xia, S.Q., Chen, L., Zhao, J.F., Renault, N.J., Chovelon, J.M., 2006. Nutrients removal from municipal wastewater by chemical precipitation in a moving bed biofilm reactor, *Process Biochem.* 41(4), 824–828.
- Wang, Z., Xu, J., Li, Y., Wang, K., Wang, Y., Hong, Q., Li, W.J., Li, S.P. 2010. *Rhodococcus jialingiae* sp. nov., an actinobacterium isolated from sludge of a carbendazim wastewater treatment facility. *Int. J. Syst. Evol. Microbiol.* 60, 378–81.
- Wanner, J., Cech, J.S., Kos, M., 1992. New process design for biological nutrient removal, *Water Sci. Technol.* 25(4-5), 445–448.