

ATTILIO CARAPEZZA & GABRIELLA LO VERDE

FIRST RECORD OF *HALYOMORPHA HALYS* (STÅL, 1855)
(Hemiptera Pentatomidae) IN SICILY

SUMMARY

The occurrence of the brown marmorated stink bug, *Halyomorpha halys* (Stål, 1855), in Sicily is here recorded for the first time. The species, native of Eastern Asia, was accidentally introduced in North America and Europe, and first found in northern Italy in 2012. In Sicily, the insect was detected in November 2017 in two different areas of Palermo town, the neighbourhood of the maritime port and the Botanical Garden. Detection of *H. halys* in Sicily was expected, due to the high spread rate shown in the different areas in which it was introduced.

RIASSUNTO

Ritrovamento in Sicilia di Halyomorpha halys (Stål, 1855) (Hemiptera Pentatomidae). Viene segnalata la presenza in Sicilia di *Halyomorpha halys* (Stål, 1855). La specie, originaria dell'Asia sud orientale, è stata accidentalmente introdotta in America ed Europa, e trovata in Italia per la prima volta nel 2012. In Sicilia è stata rinvenuta in due diverse parti della città di Palermo, nei pressi dei Cantieri Navali e vicino all'Orto Botanico. Il ritrovamento in Sicilia dell'eterottero era prevedibile, alla luce della grande velocità di dispersione manifestata in altre aree di nuova introduzione.

INTRODUCTION

The brown marmorated stink bug *Halyomorpha halys* (Stål, 1855) (Heteroptera Pentatomidae), is native to South-Eastern countries of Asia (China, Taiwan, Japan, Korea, Vietnam) (LEE *et al.*, 2013). The highly invasive behavior of *H. halys* caused a rapid expansion throughout the world (ZHU *et al.*, 2016). The insect was accidentally introduced into the United States, where it

was collected in 1998 in Pennsylvania, and then detected in 42 states (LESKEY *et al.*, 2012a, USDA-NIFA SCRI, 2017) and in Canada (FOGAIN & GRAFF, 2011). In Europe it was caught in a light trap in 2004 in Liechtenstein, and reported from Switzerland in 2007 (ARNOLD, 2009; WERMELINGER *et al.*, 2008; EPPO, 2013). In the following years, *H. halys* spread to several other countries: France, Italy, Greece, Hungary, Romania, Austria, Serbia, Spain, Russia, Bulgaria (HECKMANN, 2012; CALLOT & BRUA, 2013; MILONAS & PARTSINEVELOS, 2014; ŠEAT, 2015; MACAVEI *et al.*, 2015; DIOLI *et al.*, 2016; MITYUSHEV, 2016; SIMON, 2016).

In Italy, it was first detected in 2012 in the province of Modena, and then found in many northern and central regions (MAISTRELLO *et al.*, 2013, 2014, 2016; PANSA *et al.*, 2013; BARISELLI *et al.*, 2016) and in Sardinia (DIOLI *et al.*, 2016). For Sicily, occasional detection is reported without information about place and date (COSTI *et al.*, 2017).

The present record is the first for the town of Palermo, Sicily, where two specimens were detected in November 2017 in two different places: in the neighborhood of the maritime port and at the Botanical Garden (Fig. 1). In both cases the proximity of the seaport could explain the introduction; moreover, the Botanical Garden often holds exhibitions of exotic plants involving nursery owners from other Italian regions*.



Fig. 1 — Detection sites of *Halyomorpha halys* (Stål, 1855) in Palermo town, November 2017

* On 1st December, during the proof revision of this paper, a further Sicilian record of *H. halys* from Ispica (Ragusa) was posted on a social network (Rodolfo Occhipinti Sata Srl).

IDENTIFICATION

Halyomorpha halys is a large (12–17 mm) Pentatomidae (Figs 2a, 2b) belonging to the subfam. Pentatominae, tribe Cappaeini; it is dorsally yellowish




Fig. 2 — a. adult female of *Halyomorpha halys* (Stål, 1855) (Italy, Emilia, Sasso Marconi, 16.VI.2017, L. Colacurcio leg., A. Carapezza coll.); b. same, ventral view; c. adult female of *Rhyphogaster nebulosa* (Poda, 1761) (Italy, Sicily, Godrano, 6.I.2007, M. Arnone leg., A. Carapezza coll.); d. same, ventral view (Photo: A. Carapezza)

to pale brown in colour, mottled in brown, with a reddish tinge in the apical portion of corium, small metallic green spots and an intense black puncturation. Segments of connexivum black or metallic green with a central, trapezoid yellow spot; membrane with longitudinal dark spots on veins. Antennae with the first three segments yellow to brown with more or less intense black dots; third segment black with both base and apex black, fourth segment black with yellow base. Ventral side of body and legs yellowish brown with more or less intense black punctures. Head before eyes generally rectangular in outline with lateral margins distinctly concave, anterior angle of pronotum distinctly thorn-like, scent gland openings not continued outward by a groove.

Larvae of *H. balys* (Fig. 3) are easily distinguished from larvae of other European Pentatomidae by a long spiny process along margin of head anteriorad of eye.



Fig. 3 — 5th instar larva of *Halyomorpha balys* (Stål, 1855) (Italy, Lombardia, Milano, X.2013, P. Dioli leg.) (Photo: A. Carapezza)

Among European Pentatomidae *H. halys* is most similar to *Rhaphigaster nebulosa* (Poda, 1761) (Figs 2c, 2d), member of tribe Pentatomini, which can be separated from it by the presence of a robust spine at the base of the abdomen projecting forward, the triangular anterior part of the head, the absence of a pale ring at the apex of the fourth antennal segment, the rectangular central pale spots of segments of connexivum, the rounded dark spots on the membrane and the more convex body.

BIOLOGY AND ECONOMIC IMPORTANCE

The pest feeds on a wide range (more than 300 species) of cultivated and wild plant (LESKEY *et al.*, 2012a; LEE *et al.*, 2013; RICE *et al.*, 2014; BAKKEN *et al.*, 2015; BERGMANN *et al.*, 2016). Genetic investigations on the potential pathways of origin of *H. halys* populations in Italy allowed to assess that the haplotype present in northern Italy is the same detected in most of the USA and in the Beijing area, while in Switzerland a different haplotype was found (CESARI *et al.*, 2015).

In Europe, the biology of *H. halys* was investigated in northern Switzerland (HAYE *et al.*, 2014) and Italy (COSTI *et al.*, 2017), showing clear differences in the number of generations/year. In fact, the insect performs only one generation/year in Switzerland and two in Italy (COSTI *et al.*, 2017), as found in southern USA and in China (ZHANG *et al.*, 1993; LESKEY *et al.*, 2012b; BAKKEN *et al.*, 2015).

In the first 2-3 years after its introduction in Italy, fruit injuries were reported mainly in small untreated orchards, but in 2015 serious economic damage was recorded in commercial pear orchards, in which more than 50% of fruits were infested (COSTI *et al.*, 2017; MAISTRELLO *et al.*, 2017).

In order to limit *H. halys* populations and fruit injury, chemical control is widely applied by repeated insecticide treatments, causing disruption of previous integrated pest management programs (LESKEY *et al.*, 2012b). The potential of classical biological control is currently investigated (RICE *et al.*, 2014), even if the parasitoid species known to attack *H. halys* eggs in its native range, belonging to the genera *Trissolcus*, *Telenomus*, *Ooencyrtus* (Hymenoptera Platygasteridae) and *Anastatus* (Hymenoptera Eupelmidae), are usually not specific (HAYE *et al.*, 2014, 2015). In order to prevent negative ecological side effects derived from the introduction of not-specific natural enemies, the possibility to use local parasitoids in augmentative biological control programs is being investigated (HAYE *et al.*, 2015).

Acknowledgements — We express our gratitude to Angelo Troia and Giovanna Tusciano who reported us the presence of the insects in Palermo.

REFERENCES

- ARNOLD K., 2009. *Halyomorpha halys* (Stål, 1855), eine für die europäische Fauna neu nachgewiesene Wanzenart (Insecta: Heteroptera, Pentatomidae, Pentatominae, Cappaeini). *Mitt. Thüringer Ent.*, 16: 19.
- BAKKEN A.J., SCHOOF S.C., BICKERTON M., KAMMINGA K.L., JENRETTE J.C., MALONE S., ABNEY M.A., HERBERT D.A., REISIG D., KUCHAR T.P. & WALGENBACH, J.F., 2015. Occurrence of brown marmorated stink bug (Hemiptera: Pentatomidae) on wild hosts in non managed woodlands and soybean fields in North Carolina and Virginia. *Environ Entomol.*, 44: 1011–1021.
- BARISELLI M., BUGIANI R. & MAISTRELLO L., 2016. Distribution and damage caused by *Halyomorpha halys* in Italy. *EPPO Bulletin*, 46(2), 332–334.
- BERGMANN E.J., VENUGOPAL P.D., MARTINSON H.M., RAUPP, M.J. & SHREWSBURY, P.M., 2016. Host plant use by the invasive *Halyomorpha halys* (Stål) on woody ornamental trees and shrubs. *PLoS one*, 11(2), e0149975.
- CALLOT H. & BRUA C., 2013. *Halyomorpha halys* (Stål, 1855) la Punaise diabolique, nouvelle espèce pour la faune de France (Heteroptera Pentatomidae). *Entomologiste*, 69: 69–71.
- CESARI M., MAISTRELLO L., GANZERLI F., DIOLI P., REBECCHI L. & GUIDETTI R., 2015. A pest alien invasion in progress: potential pathways of origin of the brown marmorated stink bug *Halyomorpha halys* populations in Italy. *J. Pest Science*, 88(1): 1–7.
- COSTI E., HAYE T. & MAISTRELLO L., 2017. Biological parameters of the invasive brown marmorated stink bug, *Halyomorpha halys*, in southern Europe. *J. Pest Science*, 90(4): 1059–1067.
- DIOLI P., LEO P. & MAISTRELLO L., 2016. Prime segnalazioni in Spagna e in Sardegna della specie aliena *Halyomorpha halys* (Stål, 1855) e note sulla sua distribuzione in Europa (Hemiptera, Pentatomidae). *Rev. gaditana Entomol.*, 7(1): 539–548.
- EPPO Reporting Service no. 05 - 2013 Num. article: 2013/109
- FOGAIN R. & GRAFF S., 2011. First records of the invasive pest, *Halyomorpha halys* (Hemiptera: Pentatomidae), in Ontario and Quebec. *J. ent. Soc. Ontario*, 142: 45–48.
- HAYE T., ABDALLAH S., GARIEPY T., & WYNIGER D., 2014. Phenology, life table analysis and temperature requirements of the invasive brown marmorated stink bug, *Halyomorpha halys*, in Europe. *J. Pest Science*, 87(3): 407–418.
- HAYE T., FISCHER S., ZHANG J. & GARIEPY T., 2015. Can native egg parasitoids adopt the invasive brown marmorated stink bug, *Halyomorpha halys* (Heteroptera: Pentatomidae), in Europe? *J. Pest Science*, 88(4): 693–705.
- HECKMANN R., 2012. Erster Nachweis von *Halyomorpha halys* (Stål, 1855) (Heteroptera: Pentatomidae) für Deutschland. *Heteropteron*, 36: 17–18.
- LEE D.H., SHORT, B.D., JOSEPH, S.V., BERGH, J.C. & LESKEY T.C., 2013. Review of the biology, ecology, and management of *Halyomorpha halys* (Hemiptera: Pentatomidae) in China, Japan, and the Republic of Korea. *Env. Entom.*, 42(4): 627–641.
- LESKEY T.C., HAMILTON G.C., NIELSEN A.L., POLK D.F., RODRIGUEZ-SAONA C., BERGH J.C., AMES HERBERT D., KUCHAR T.P., PFEIFFER D., DIVELY G.P., HOOKS C.R.R., RAUPP M.J., SHREWSBURY P.M., KRAWCZYK G., SHEARER P.W., WHALEN J., KOPLINKA-LOEHR C., MYERS E., INKLEY D., HOELMER K.A., LEE D.H. & WRIGHT S.E., 2012a. Pest status of the brown marmorated stink bug, *Halyomorpha halys* in the USA. *Outlooks Pest Manag.*, 23(5): 218–226.
- LESKEY T. C., WRIGHT S. E., SHORT B. D. & KHRIMIAN A., 2012. Development of behaviorally-based monitoring tools for the brown marmorated stink bug (Heteroptera: Pentatomidae) in commercial tree fruit orchards. *J. ent. Science*, 47(1): 76–85.
- MACAVEI L.I., BĂETAN R., OLTEAN I., FLORIAN T., VARGA M., COSTI E. & MAISTRELLO L., 2015. First

- detection of *Halyomorpha halys* Stål, a new invasive species with a high potential of damage on agricultural crops in Romania. *Lucrări științifice, Universitatea de Științe Agricole și Medicină Veterinară "Ion Ionescu de la Brad" Iași, Seria Agronomie*, 58(1): 105-108.
- MAISTRELLO L., DIOLI P. & BARISELLI M., 2013. Trovata una cimice esotica dannosa per i frutteti. *Agricoltura*, 6: 67-68.
- MAISTRELLO L., DIOLI P., BARISELLI M., MAZZOLI G. L. & GIACALONE-FORINI I., 2016. Citizen science and early detection of invasive species: phenology of first occurrences of *Halyomorpha halys* in Southern Europe. *Biol. Invasions*, 18(11): 3109-3116.
- MAISTRELLO L., DIOLI P., VACCARI G., NANNINI R., BORTOLOTTI P., CARUSO S., COSTI E., MONTERMINI A., CASOLI L. & BARISELLI M., 2014. Primi rinvenimenti in Italia della cimice esotica *Halyomorpha halys*, una nuova minaccia per la frutticoltura. *Atti Giornate Fitopatologiche CLUEB*, Bologna: 283-288.
- MAISTRELLO L., VACCARI G., CARUSO S., COSTI E., BORTOLINI S., MACAVEI L., FOCA G., ULRICI A., BORTOLOTTI P.P., NANNINI R., CASOLI L., FORNACIARI M., MAZZOLI G.L. & DIOLI P., 2017. Monitoring of the invasive *Halyomorpha halys*, a new key pest of fruit orchards in northern Italy. *J. Pest Science*, 90(4): 1231-1244.
- MILONAS P.G. & PARTSINEVELOU G.K., 2014. First report of brown marmorated stink bug *Halyomorpha halys* Stål (Hemiptera: Pentatomidae) in Greece. *OEPP/EPPO Bull.*, 44: 183-186.
- MITYUSHEV I.M., 2016. First record of marmorated bug detection in Russia. *Zashchita i Karantin Rastenii*, 3: 48 (in Russian).
- PANSA M.G., ASTEGGIANO L., COSTAMAGNA C., VITTONI G. & TAVELLA L., 2013. First discovery of *Halyomorpha halys* in peach orchards in Piedmont. *Inf. re agr.*, 69: 60-61.
- RICE K., BERGH C., BERGMAN E., BIDDINGER D., DIECKHOFF C., DIVELY G., FRASER H., GARIEPY T., HAMILTON G., HAYE T., HERBERT A., HOELMER K., HOOKS C., JONES A., KRAWCZYK G., KUCHAR T., MITCHELL W., NIELSEN A., PFEIFFER D., RAUPP M., RODRIGUEZ-SAONA C., SHEARER P., SHREWSBURY P., VENUGOPAL P.D., WHALEN J., WIMAN N., LESKEY T. & TOOKER J., 2014. Biology, ecology, and management of brown semistriatus marmorated stink bug (*Halyomorpha halys*). *J. Integr. Pest Manag.*, 5(3): A1-A13.
- ŠEAT J., 2015. *Halyomorpha halys* (Stål, 1855) (Heteroptera: Pentatomidae) a new invasive species in Serbia. *Acta entom. serbica*, 20: 167-171.
- SIMOV N., 2016. The invasive brown marmorated stink bug, *Halyomorpha halys* (Stål, 1855) (Heteroptera: Pentatomidae) already in Bulgaria. *Ecol. montenegrina*, 9: 51-53.
- USDA-NIFA SCRI (2017) Stop BMSB. Biology, ecology, and management of brown marmorated stink bug in specialty crop. In: Stop BMSB. <http://www.stopbmsb.org/where-is-bmsb/>. Accessed 20 November 2017.
- WERMELINGER B., WYNIGER D. & FORSTER B., 2008. First records of an invasive bug in Europe: *Halyomorpha halys* Stål (Heteroptera: Pentatomidae), a new pest on woody ornamentals and fruit trees? *Mitt. Schweiz Entomol. Ges.*, 81: 1-8.
- ZHANG C.T., LI D., SU H. & XU G., 1993. A study on the biological characteristics of *Halyomorpha picus* and *Erthesina fullo*. *For Res. China*, 6: 271-275.
- ZHU G., BU W., GAO Y. & LIU G., 2012. Potential geographic distribution of brown marmorated stink bug invasion (*Halyomorpha halys*). *PLoS One*, 7(2), E31246.

Authors' addresses — A. CARAPEZZA, via Sandro Botticelli, 15 – 90144 Palermo; email: attilio.carapezza@unipa.it; G. LO VERDE, Department of Agricultural, Food and Forest Science, University of Palermo, Viale delle Scienze Ed. 5A - 90128 Palermo (I); email: gabriella.loverde@unipa.it