

SEPARATION ANXIETY IN PEDIATRIC MIGRAINE WITHOUT AURA: A PILOT STUDY

MARGHERITA SALERNO², AGATA MALTESE¹, GABRIELETRIPPI^{3,4}, PALMIRA ROMANO⁵, ANNABELLA DI FOLCO¹, TERESA DI FILIPPO¹, LUCIA PARISI¹

¹Department of Psychological, Pedagogical and Educational Sciences, University of Palermo, Italy - ²Sciences for Mother and Child Health Promotion, University of Palermo, Italy - ³Department PROSAMI, University of Palermo, Italy - ⁴Childhood Psychiatric Service for Neurodevelopmental Disorders, CH Chinon, France - ⁵Clinic of Child and Adolescent Neuropsychiatry, Department of Mental Health and Physical and Preventive Medicine, Università degli Studi della Campania-Luigi Vanvitelli, Italy

ABSTRACT

Background: Separation anxiety (SA) can be defined as the fear reaction and protest manifested by children when the main caregivers move away from him/her or in front of unfamiliar person. SA near eight months may be considered as an important and normal phase of the correct and typical social neurodevelopment. Aims of the present pilot study is assessing the prevalence of separation anxiety in a population of patients with migraine without aura (MwA).

Materials and methods: 119 children (69 males) suffering from MwA (mean age $1.78 \pm 7:59$) were consecutively recruited. The control population consisted of 231 (114 males) healthy subjects similar for age ($7.64 \pm 1:34$; $p = 0.768$) and gender ($p=0.987$)

The Screen for child anxiety related emotional disorders (SCARED) test was used to assess the prevalence of separation anxiety among MwA children.

Results: Healthy individuals are on average less affected (mean $4.72 \pm 0:32$) of the Separation Anxiety Disorder respect of MwA children (mean 6.83 ± 0.97 ; $p < 0.001$) (Figure 1).

Conclusion. MwA presents many psychiatric comorbidities and among ones separation anxiety may be considered in the clinical and therapeutic management of pediatric primary headache.

Keywords: Migraine without aura, Mw, separation anxiety, SCARED.

DOI: 10.19193/0393-6384_2017_4_092

Received November 30, 2016; Accepted March 20, 2017

Introduction

Separation anxiety (SA) can be defined as the fear reaction and protest manifested by children when the main caregivers move away from him/her or in front of unfamiliar person. SA near eight months may be considered as an important and normal phase of the correct and typical social neurodevelopment. The Austrian psychoanalyst Renè Spitz described the stage at which SA occurs as the stranger's anguish or distress of the eighth month, in

which the child learns to differentiate between the various human faces and react positively and negatively to those familiar to him those strangers to him. In this light, SA is an important achievement in terms of development of its social and interpersonal skills and, as such, is a necessary step but transitional. Many clinical conditions may elicit and perpetuate the natural SA, particularly algic conditions such as migraine headache⁽¹⁻¹²⁾. Migraine without aura (MwA) has many comorbidities also psychiatric ones. In this light, aim of the present pilot study is

assessing the prevalence of separation anxiety in a population of patients with migraine without aura (MwA).

Materials and methods

119 children (69 males) suffering from MwA (mean age $1.78 \pm 7:59$) were consecutively recruited. MwA diagnosis was made according to ICHD-3 criteria.

Exclusion criteria were the following: overweight ($z\text{-BMI} > 85$ pc) and obesity ($z\text{-BMI} > 95$ pc), cognitive disability ($IQ < 70$), neurological disorders (ie headaches, epilepsy), chromosomal syndromes (eg. Down, Prader-Willi, Crouzon, Pierre-Robin, trisomy 18), psychiatric illness (ie. mood disorders, anxiety disorders, psychosis) and specific neuropsychological disorders (13-38).

The control population consisted of 231 (114 males) healthy subjects similar for age ($7.64 \pm 1:34$; $p = 0.768$) and gender ($p = 0.987$)

Screen for child anxiety related emotional disorders (SCARED)

MwA children filled in the Italian version of the 41-item SCARED questionnaire, and were asked to rate the frequency with which they experienced each symptom on a 3-point likert scale (0 = "almost never," 1 = "sometimes," 2 = "often"). Via principal component factor analysis, five subscales were identified: Panic/Somatic Anxiety (PD), General Anxiety (GAD), Separation Anxiety (SAD), Social Phobia (SOC), and School Phobia (SCH). With reference to the psychometric properties, the tool showed good internal consistency, test-retest reliability, discriminative validity. When a cut-off point of 25 was applied to the total score endorsed by subjects across these five factors, data showed good sensitivity (70%) and good ability to discriminate between children with AD versus those without AD (specificity: 67%), and between children with AD versus those with depression, or disruptive disorders: 61 and 71%, respectively.

Specifically, we take into account only the Separation anxiety scale with a cut-off of 5 points.

Statistical analysis

STATISTICA 6.0 software (StatSoft, Inc.); was used to perform the statistical comparison with t-test and chi-squared when appropriated. p values ≤ 0.005 were used as significant.

Results

Healthy individuals are on average less affected (mean $4.72 \pm 0:32$) of the Separation Anxiety Disorder respect of MwA children (mean 6.83 ± 0.97 ; $p < 0.001$) (Figure 1).




Figure 1. shows the higher prevalence of separation anxiety among MwA suffering children than controls.

Discussion

According to the bio-psychosocial perspective, the separation anxiety disorder may be linked to genetic factors, such as an inhibited temperament that would make the child more anxious and dependent on the mother, and negative life events and stressful experiences during childhood. In this light, caregiving styles have responsibilities: overprotective parents, anxious, unresolved, obsessive, boost the child's discomfort and increase your anxiety levels. The disorder, if not resolved, can persist into adulthood and manifest itself in a different form, eg. through a disorder, panic attacks, agoraphobia, generalized anxiety. All these psychiatric manifestations are prevalent in children and adults suffering from MwA⁽³⁹⁻⁵⁰⁾.

The separation anxiety disorder seems, therefore, arises more frequently in migraine sufferers than in control subjects, probably this could be the result from the changes in the educational pipeline implemented by the parents in front of the management of painful episodes of the child, however, this argument should be confirmed by performing additional psychodiagnostic investigation⁽⁴⁰⁻⁴⁵⁾.

Although the limit of this study is the reduced statistical sample ($d = -2921$), our findings suggest, in agreement with the common clinical practice, the need for an approach to migraine pathology based on total care of the patient and his family, in order to allow more equitable management of development psychology of the subject.

References

- 1) Carotenuto M, Esposito M. Nutraceuticals safety and efficacy in migraine without aura in a population of children affected by neurofibromatosis type I. *Neurol Sci.* 2013 Nov; 34(11): 1905-9. doi: 10.1007/s10072-013-1403-z.
- 2) Perillo L, Esposito M, Caprioglio A, Attanasio S, Santini AC, Carotenuto M. Orthodontic treatment need for adolescents in the Campania region: the malocclusion impact on self-concept. *Patient Prefer Adherence.* 2014 Mar 19;8:353-9. doi: 10.2147/PPA.S58971.
- 3) Carotenuto M, Gimigliano F, Fiordelisi G, Ruberto M, Esposito M. Positional abnormalities during sleep in children affected by obstructive sleep apnea syndrome: the putative role of kinetic muscular chains. *Med Hypotheses.* 2013 Aug; 81(2): 306-8. doi: 10.1016/j.mehy.2013.04.023.
- 4) Esposito M, Carotenuto M. Intellectual disabilities and power spectra analysis during sleep: a new perspective on borderline intellectual functioning. *J Intellect Disabil Res.* 2014 May; 58(5):421-9. doi: 10.1111/jir.12036.
- 5) Esposito M, Parisi P, Miano S, Carotenuto M. Migraine and periodic limb movement disorders in sleep in children: a preliminary case-control study. *J Headache Pain.* 2013 Jul 1; 14:57. doi: 10.1186/1129-2377-14-57
- 6) Carotenuto M, Esposito M, Pascotto A. Migraine and enuresis in children: An unusual correlation? *Med Hypotheses.* 2010 Jul; 75(1): 120-2. doi: 10.1016/j.mehy.2010.02.004.
- 7) Carotenuto M, Gallai B, Parisi L, Roccella M, Esposito M. Acupressure therapy for insomnia in adolescents: a polysomnographic study. *Neuropsychiatr Dis Treat.* 2013; 9: 157-62. doi: 10.2147/NDT.S41892.
- 8) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Gritti A, Mazzotta G, Carotenuto M. Maternal stress and childhood migraine: a new perspective on management. *Neuropsychiatr Dis Treat.* 2013; 9: 351-5. doi: 10.2147/NDT.S42818.
- 9) Carotenuto M, Esposito M, Precenzano F, Castaldo L, Roccella M. Cosleeping in childhood migraine. *Minerva Pediatr.* 2011 Apr ;63(2): 105-9.
- 10) Esposito M, Carotenuto M, Roccella M. Primary nocturnal enuresis and learning disability. *Minerva Pediatr.* 2011 Apr; 63(2): 99-104.
- 11) Esposito M, Roccella M, Parisi L, Gallai B, Carotenuto M. Hypersomnia in children affected by migraine without aura: a questionnaire-based case-control study. *Neuropsychiatr Dis Treat.* 2013;9:289-94. doi: 10.2147/NDT.S42182.
- 12) Esposito M, Pascotto A, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Gritti A, Mazzotta G, Carotenuto M. Can headache impair intellectual abilities in children? An observational study. *Neuropsychiatr Dis Treat.* 2012;8:509-13. doi: 10.2147/NDT.S36863.
- 13) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Mazzotta G, Patriciello G, Precenzano F, Carotenuto M. Visuomotor competencies and primary monosymptomatic nocturnal enuresis in prepubertal aged children. *Neuropsychiatr Dis Treat.* 2013; 9:921-6. doi: 10.2147/NDT.S46772.
- 14) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, Vagliano C, Messina G, Di Folco A, Di Filippo T, Michele Roccella. Executive functioning in preschool children affected by autism spectrum disorder: a pilot study. *Acta Medica Mediterranea,* 2017, 33: 35-39; DOI: 10.19193/0393-6384_2017_1_005.
- 15) Precenzano F, Lombardi P, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Della Valle I, Magliulo RM, Messina G, Roccella M. Internalizing symptoms in children affected by childhood absence epilepsy: a preliminary study. *Acta Medica Mediterranea,* 2016, 32: 1749-1753; DOI: 10.19193/0393-6384_2016_6_158.
- 16) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Della Valle I, Visco G, Magliulo RM, Messina G, Roccella M. ADHD-like symptoms in children affected by obstructive sleep apnea syndrome: a case-control study. *Acta Medica Mediterranea,* 2016, 32: 1755-1759; DOI: 10.19193/0393-6384_2016_6_159.
- 17) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Grappa MF, Magliulo RM, Messina G, Roccella M. Borderline intellectual functioning and parental stress: an italian case-control study. *Acta Medica Mediterranea,* 2016, 32: 1761-1765; DOI: 10.19193/0393-6384_2016_6_160.
- 18) Ruberto M, Precenzano F, Parisi L, Salerno M, Maltese A, Messina G, Roccella M. Visuomotor integration skills in children affected by obstructive sleep apnea syndrome: a case-control study. *Acta Medica Mediterranea,* 2016, 32: 1659; DOI: 10.19193/0393-6384_2016_5_146.
- 19) Parisi L, Ruberto M, Precenzano F, Di Filippo T, Russotto C, Maltese A, Salerno M, Roccella M. The quality of life in children with cerebral palsy. *Acta Medica Mediterranea,* 2016, 32: 1665; DOI: 10.19193/0393-6384_2016_5_147.
- 20) Epifanio, M.S., Genna, V., De Luca, C., Roccella, M., La Grutta, S. Paternal and maternal transition to parenthood: The risk of postpartum depression and parenting stress (2015) *Pediatric Reports,* 7 (2), pp. 38-44.
- 21) Parisi, L., Di Filippo, T., Roccella, M. The child with Autism Spectrum Disorders (ASDS): Behavioral and neurobiological aspects. *Acta Medica Mediterranea,* 2015, 31 (6), pp. 1187-1194.
- 22) Vecchio, D., Salzano, E., Vecchio, A., Di Filippo, T., Roccella, M. A case of femoral-facial syndrome in a patient with autism spectrum disorders. *Minerva Pediatrica,* 2011, 63 (4), pp. 341-344.
- 23) Parisi, L., Di Filippo, T., Roccella, M. Hypomelanosis of Ito: Neurological and psychiatric pictures in developmental age. *Minerva Pediatrica,* 2012, 64 (1), pp. 65-70.
- 24) Di Filippo, T., Parisi, L., Roccella, M. Psychological aspects in children affected by duchenne de boulogne muscular dystrophy. *Mental Illness,* 2012, 4 (1), pp. 21-24.
- 25) Epifanio MS, Genna V, Vitello MG, Roccella M, La Grutta S. Parenting stress and impact of illness in parents of children with coeliac disease. *Pediatr Rep.* 2013 Dec 19; 5(4): e19. doi: 10.4081/pr.2013.e19.
- 26) Esposito M, Parisi L, Gallai B, Marotta R, Di Dona A, Lavano SM, Roccella M, Carotenuto M. Attachment styles in children affected by migraine without aura. *Neuropsychiatr Dis Treat.* 2013; 9: 1513-9. doi:

- 10.2147/NDT.S52716.
- 27) Di Filippo T, Orlando MF, Concialdi G, La Grutta S, Lo Baido R, Epifanio MS, Esposito M, Carotenuto M, Parisi L, Roccella M. The quality of life in developing age children with celiac disease. *Minerva Pediatr.* 2013 Dec; 65(6): 599-608.
- 28) Maltese A, Pepi A, Scifo L, Roccella M. Referential communication skills in children with Down Syndrome. *Minerva Pediatr.* 2014 Feb; 66(1): 7-16.
- 29) Esposito M, Marotta R, Roccella M, Gallai B, Parisi L, Lavano SM, Carotenuto M. Pediatric neurofibromatosis 1 and parental stress: a multicenter study. *Neuropsychiatr Dis Treat.* 2014 Jan 22;10:141-6. doi: 10.2147/NDT.S55518;
- 30) Alesi M, Battaglia G, Roccella M, Testa D, Palma A, Pepi A. Improvement of gross motor and cognitive abilities by an exercise training program: three case reports. *Neuropsychiatr Dis Treat.* 2014 Mar 14;10:479-85. doi: 10.2147/NDT.S58455;
- 31) Panico A, Messina G, Lupoli GA, Lupoli R, Cacciapuoti M, Moscatelli F, Esposito T, Villano I, Valenzano A, Monda V, Messina A, Precenzano F, Cibelli G, Monda M, Lupoli G. Quality of life in overweight (obese) and normal-weight women with polycystic ovary syndrome. *Patient Prefer Adherence.* 2017 Mar 2; 11: 423-429.
- 32) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, Gallai B, Marotta R, Lavano SM, Lavano F, Roccella M. Visual-spatial training efficacy in children affected by migraine without aura: a multicenter study. *Neuropsychiatr Dis Treat.* 2017 Jan 27; 13:253-258. doi: 10.2147/NDT.S119648.
- 33) Epifanio, M.F., Genna, V., Di Marco, S., Furnari, M.L., Pardo, F., Collura, M., Roccella, M., La Grutta, S. Quality of life, affect regulation and resilience in adult patients with cystic fibrosis. *Gazzetta Medica Italiana Archivio per le Scienze Mediche*, 2013 172 (9), pp. 705-711.
- 34) Parisi L, Di Filippo T, La Grutta S, Lo Baido R, Epifanio MS, Esposito M, Carotenuto M, Roccella M. Sturge-weber syndrome: a report of 14 cases. *Ment Illn.* 2013 Jun 3; 5(1): e7. doi: 10.4081/mi.2013.e7.
- 35) Chieffi S, Messina G, Villano I, Messina A, Esposito M, Monda V, Valenzano A, Moscatelli F, Esposito T, Carotenuto M, Viggiano A, Cibelli G, Monda M. Exercise Influence on Hippocampal Function: Possible Involvement of Orexin-A. *Front Physiol.* 2017 Feb 14; 8: 85. doi: 10.3389/fphys.2017.00085.
- 36) Villano I, Messina A, Valenzano A, Moscatelli F, Esposito T, Monda V, Esposito M, Precenzano F, Carotenuto M, Viggiano A, Chieffi S, Cibelli G, Monda M, Messina G. Basal Forebrain Cholinergic System and Orexin Neurons: Effects on Attention. *Front Behav Neurosci.* 2017 Jan 31; 11: 10. doi: 10.3389/fnbeh.2017.00010.
- 37) Matricardi S, Spalice A, Salpietro V, Di Rosa G, Balistreri MC, Grosso S, Parisi P, Elia M, Striano P, Accorsi P, Cusmai R, Specchio N, Coppola G, Savasta S, Carotenuto M, Tozzi E, Ferrara P, Ruggieri M, Verrotti A. Epilepsy in the setting of full trisomy 18: A multicenter study on 18 affected children with and without structural brain abnormalities. *Am J Med Genet C Semin Med Genet.* 2016 Sep; 172(3): 288-95. doi: 10.1002/ajmg.c.31513;
- 38) Messina A, De Fusco C, Monda V, Esposito M, Moscatelli F, Valenzano A, Carotenuto M, Viggiano E, Chieffi S, De Luca V, Cibelli G, Monda M, Messina G. Role of the Orexin System on the Hypothalamus-Pituitary-Thyroid Axis. *Front Neural Circuits.* 2016 Aug 25; 10: 66. doi: 10.3389/fncir.2016.00066;
- 39) Moscatelli F, Valenzano A, Petito A, Triggiani AI, Ciliberti MAP, Luongo L, Carotenuto M, Esposito M, Messina A, Monda V, Monda M, Capranica L, Messina G, Cibelli G. Relationship between blood lactate and cortical excitability between taekwondo athletes and non-athletes after hand-grip exercise. *Somatosens Mot Res.* 2016 Jun;33(2):137-44. doi: 10.1080/08990220.2016.1203305.
- 40) Carotenuto M, Esposito M, Cortese S, Laino D, Verrotti A. Children with developmental dyslexia showed greater sleep disturbances than controls, including problems initiating and maintaining sleep. *Acta Paediatr.* 2016 Sep; 105(9): 1079-82. doi: 10.1111/apa.13472.
- 41) Pasquali D, Carotenuto M, Leporati P, Esposito M, Antinolfi L, Esposito D, Accardo G, Carella C, Chiovato L, Rotondi M. Maternal hypothyroidism and subsequent neuropsychological outcome of the progeny: a family portrait. *Endocrine.* 2015 Dec; 50(3): 797-801. doi: 10.1007/s12020-015-0564-3.
- 42) Morandi A, Bonnefond A, Lobbens S, Carotenuto M, Del Giudice EM, Froguel P, Maffei C. A girl with incomplete Prader-Willi syndrome and negative MS-PCR, found to have mosaic maternal UPD-15 at SNP array. *Am J Med Genet A.* 2015 Nov; 167A(11): 2720-6. doi: 10.1002/ajmg.a.37222.
- 43) Verrotti A, Carotenuto M, Altieri L, Parisi P, Tozzi E, Belcastro V, Esposito M, Guastaferrò N, Ciuti A, Mohn A, Chiarelli F, Agostinelli S. Migraine and obesity: metabolic parameters and response to a weight loss programme. *Pediatr Obes.* 2015 Jun; 10(3): 220-5. doi: 10.1111/ijpo.245.
- 44) Esposito M, Precenzano F, Sorrentino M, Avolio D, Carotenuto M. A Medical Food Formulation of Griffonia simplicifolia/Magnesium for Childhood Periodic Syndrome Therapy: An Open-Label Study on Motion Sickness. *J Med Food.* 2015 Aug; 18(8): 916-20. doi: 10.1089/jmf.2014.0113.
- 45) Franzoni E, Matricardi S, Di Pisa V, Capovilla G, Romeo A, Tozzi E, Pruna D, Salerno GG, Zamponi N, Accorsi P, Giordano L, Coppola G, Cerminara C, Curatolo P, Nicita F, Spalice A, Grosso S, Pavone P, Striano P, Parisi P, Boni A, Gobbi G, Carotenuto M, Esposito M, Cottone C, Verrotti A. Refractory absence seizures: An Italian multicenter retrospective study. *Eur J Paediatr Neurol.* 2015 Nov;19(6): 660-4. doi: 10.1016/j.ejpn.2015.07.008;
- 46) Esposito M, Gallai B, Roccella M, Marotta R, Lavano F, Lavano SM, Mazzotta G, Bove D, Sorrentino M, Precenzano F, Carotenuto M. Anxiety and depression levels in prepubertal obese children: a case-control study. *Neuropsychiatr Dis Treat.* 2014 Oct 3; 10:1897-902. doi: 10.2147/NDT.S69795;
- 47) Carotenuto M, Esposito M, Pascotto A. Facial patterns and primary nocturnal enuresis in children. *Sleep Breath.* 2011 May; 15(2): 221-7. doi: 10.1007/s11325-010-0388-6;
- 48) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R,

- Lavano SM, Mazzotta G, Carotenuto M. Primary nocturnal enuresis as a risk factor for sleep disorders: an observational questionnaire-based multicenter study. *Neuropsychiatr Dis Treat*. 2013;9:437-43. doi: 10.2147/NDT.S43673;
- 49) Santamaria F, Esposito M, Montella S, Cantone E, Mollica C, De Stefano S, Mirra V, Carotenuto M. Sleep disordered breathing and airway disease in primary ciliary dyskinesia. *Respirology*. 2014 May;19(4):570-5. doi: 10.1111/resp.12273;
- 50) Carotenuto M, Esposito M, D'Aniello A, Rippa CD, Precenzano F, Pascotto A, Bravaccio C, Elia M. Polysomnographic findings in Rett syndrome: a case-control study. *Sleep Breath*. 2013 Mar; 17(1): 93-8. doi: 10.1007/s11325-012-0654-x

Corresponding author

MARGHERITA SALERNO; MD;
Sciences for Mother and Child Health Promotion
University of Palermo
(Italy)