



# The Effects of 8-Week Mindfulness-Based Stress Reduction Program on Cognitive Control: an EEG Study

Francesca Incagli<sup>1</sup> · Vincenza Tarantino<sup>2</sup> · Cristiano Crescentini<sup>3</sup> · Antonino Vallesi<sup>1,4</sup> 

© Springer Science+Business Media, LLC, part of Springer Nature 2019

## Abstract

**Objectives** Mindfulness practice can enhance different aspects of attentional functions, such as the ability to sustain the attentional focus over time. However, it is still unclear whether this practice might indeed impact higher cognitive functions, such as control mechanisms that allow the appropriate and flexible allocation of attentional resources. In this longitudinal study, changes associated with a mindfulness-based stress reduction (MBSR) program were investigated, with a focus on proactive and reactive cognitive control mechanisms, namely, the ability to maintain task-relevant information and to prepare in advance the response, and the ability to promptly adjust overlearned behaviors in response to conflicting stimuli.

**Methods** Two groups of participants took part in the study: 26 participants who completed a formal MBSR training (mean age = 43 years, females = 21) and 23 participants who performed a control training (mean age = 47.2 years, females = 20). They were tested on a modified AX-continuous performance task (AX-CPT), before and after eight weeks of training. The electroencephalographic (EEG) signal was recorded during task execution, and amplitude modulations of event-related potentials (ERPs) associated with cues and probes were examined.

**Results** After the training, the MBSR group exhibited a significant reduction of errors on high conflicting trials. Concurrently, the Contingent Negative Variation (CNV), an ERP index of anticipatory processes elicited by task cues, became more pronounced in the post-training session in the MBSR group only. In addition, an attenuated probe-locked N2 and an increased P3a component emerged.

**Conclusions** Taken together, the behavioral and electrophysiological results suggest that the mindfulness practice enhanced the ability to implement both proactive and reactive cognitive control processes.

**Keywords** Meditation · Cognitive control · AX-CPT · Event-related potentials (ERP) · Mindfulness-Based Stress Reduction program (MBSR) · Proactive control

---

✉ Antonino Vallesi  
antonino.vallesi@unipd.it

<sup>1</sup> Department of Neuroscience & Padova Neuroscience Center, University of Padova, Via Giustiniani 5, 35128 Padua, Italy

<sup>2</sup> Department of Psychology, Educational Science and Human Movement, University of Palermo, Viale delle Scienze Ed. 15, 90128 Palermo, Italy

<sup>3</sup> Department of Languages and Literatures, Communication, Education and Society, University of Udine, Via Margreth 3, 33100 Udine, Italy

<sup>4</sup> Brain Imaging and Neural Dynamics Research Group, IRCCS San Camillo Hospital, Venice, Italy

Mindfulness has been conceptualized as an attentional regulatory training, which aims to calm and stabilize mind while cultivating an open and nonelaborating attitude toward one's own experience (Lutz et al. 2008; Malinowski 2013). In recent years, the study of the cognitive effects of mindfulness is receiving a growing interest. However, there is no clear consensus about the exact cognitive processes that are involved in and shaped by this meditation practice. Recently, some interesting attempts have been made to create a coherent theoretical framework regarding the neurocognitive processes underlying mindfulness (e.g., Raffone et al. 2019). A large body of research has shown mindfulness-related effects on attentional processes, such as selective and sustained attention, as well as on executive functions, such as inhibition (for reviews, see Chiesa et al. 2011; Gallant 2016; Lutz et al. 2008).

However, little research has been devoted to investigate the neural dynamics underpinning these changes (see Tang et al. 2015 for a review).

A functional magnetic resonance study showed an increase of the hemodynamic brain activity in the orbitofrontal cortex during a body mental rotation task, after an 8-week mindfulness training (Tomasino et al. 2016). Another longitudinal study found a cortical thickness increase in the right insula and in the somatosensory cortex after an 8-week Mindfulness-Based Stress Reduction (MBSR) training (Santarnecchi et al. 2014). Interestingly, this increase correlated with a decrease in psychological indices associated to alexithymia, worry, anxiety, and depression. Group differences in the right insula, and specifically a stronger functional coupling of this area with the lateral prefrontal cortex, were also found in a cross-sectional study comparing MBSR practitioners to a naive group (Farb et al. 2007). This effect was interpreted as a change in self-referential processes. Overall, neuroimaging findings have suggested that mindfulness training might impact brain structures and their functioning.

Similarly, electroencephalographic (EEG) evidence has accounted for meditation-related effects on brain activity (see Cahn and Polich 2006 for an extensive review). For instance, Moore et al. (2012) compared the EEG markers of executive functions of a group of participants in an 8-week MBSR program with a waiting list control group. To this end, the classical Color-Word Stroop task was administered, which requires to inhibit a dominant automatized process (e.g., reading a word) in favor of a task-relevant but weaker one (e.g., naming the font color of the word). At the behavioral level, no differential changes were found between the meditation and the control group. In contrast, a significant effect of mindfulness practice emerged in two event-related potential (ERP) components, that is, the N2 and P3, over occipito-parietal and parietal scalp sites, respectively. Interestingly, the P3 amplitude decreased when evoked by incongruent stimuli. This finding was interpreted as less demanding allocation of attentional resources to monitor conflicting stimulus–response (S–R) associations, in the meditation group only (Moore et al. 2012). A meditation-related effect on a more anterior P3 component (i.e., the P3a) was also observed in a cross-sectional study employing a passive three-stimulus oddball task (Cahn and Polich 2009). The smaller P3a amplitude elicited by novel stimuli was interpreted as a relative disengagement of the exogenous/stimulus-driven orienting system, since the system became more internally directed and less responsive to unexpected task-unrelated changes. A meditation-related benefit on the ability to allocate attentional resources was also observed in a study by Slagter et al. (2007), which employed the attentional blink paradigm to assess the effects of a 3-month meditation retreat. In this study, the group of meditators exhibited an improved detection of the target stimulus when the P3 evoked by the preceding stimulus was attenuated. This evidence suggested that meditation

training resulted in more efficient deployment of attentional resource allocation over time, in accordance with task requirements (Slagter et al. 2007).

Another ERP component that was found to be modulated by meditation practice is the Contingent Negative Variation (CNV; Walter 1964). This potential is typically observed over fronto-central scalp sites and shows a negative going waveform and a slow development. Its peculiarity is that it shows up in anticipation of the onset of target stimuli and its amplitude increases (i.e., becomes more negative) when these targets are task-relevant. Specifically, the CNV is evoked by cue stimuli that announce upcoming probe stimuli and reflects expectancy and preparatory processes (see Mento 2013 for an extensive review). Two studies, comparing meditators with different length of experience in transcendental meditation (TM) and different frequency of self-reported transcendental experiences, reported more negative CNV amplitude over central scalp sites in more experienced meditators (Travis et al. 2000, 2002). Interestingly, when a distracting task was presented (i.e., a short-term memory task), the disruption of the typical CNV development (distraction effect) was higher in less experienced meditators, compared to more experienced ones (Travis et al. 2000). Unfortunately, so far, collected evidence concerning the CNV index is limited to cross-sectional designs; longitudinal evidence is needed in order to make more compelling inferences about the relation between meditation practice and observed changes in preparatory processes between meditators and nonmeditators (Moore et al. 2012).

Taken together, the ERP findings suggest that meditation practice influences cognitive control functions. Namely, the abilities to allocate attentional resources to task-relevant information over time, to suppress task-unrelated information, and to engage preparatory processes are affected. According to the Dual Mechanism of Cognitive Control (DMC), theorized by Braver et al. (2007), cognitive control includes a set of attentional and executive processes that allow individuals to flexibly adapt responses to task demands, emphasizing goal-directed behaviors over habituation and dominant response. Specifically, an efficient cognitive control results from a dynamic interaction between two qualitatively distinct mechanisms, that is, proactive and reactive cognitive control mechanisms (Braver et al. 2007; Braver 2012). Proactive cognitive control mechanisms includes future-oriented “early-selection” processes, which allow to actively maintain contextual information prior to the occurrence of the demanding events, in order to prepare the response. These preparatory processes bias successive cognitive processing and affect task performance (e.g., memory, perception, and action). Reactive cognitive control is instead intended as a past-oriented, “late-correction” process, activated *just-in-time* by target stimuli, to detect and solve interference, when required.

Recently, cognitive control mechanisms have been investigated in people who participated to a MBSR program

(Kabat-Zinn 1990), a structured and standardized intervention. In a study (Li et al. 2018), the AX-Continuous Performance Task (AX-CPT) paradigm was used. This paradigm tests the dynamic interaction between proactive and reactive cognitive control mechanisms (e.g., Dias et al. 2003; Gonthier et al. 2016; Morales et al. 2015; Richmond et al. 2015). The task requires to respond to probe stimulus according to the preceding cue stimulus. The study by Li et al. (2018) showed a shift toward the use of more proactive control processes in a group engaged in a Mindfulness-Based Cognitive Therapy (MBCT) program. However, no significant interactions emerged when comparing the MBCT with a waiting list control group. Moreover, a follow-up assessment of participants who completed the mindfulness intervention was missing. Therefore, the study was not robust enough to draw firm conclusions about the effectiveness of mindfulness training on cognitive control mechanisms. Another recent study documented an association between either higher mindfulness traits, as tested by a self-report questionnaire, or a brief mindfulness intervention, and a better balance between the proactive and reactive control systems (Chang et al. 2017). Unfortunately, in this study only behavioral measures have been examined. Thus, there is a need to complement behavioral data with electrophysiological measures in order to elucidate the degree of change of these two interacting control processes, both behaviorally and neurally, after a structured mindfulness practice.

The choice of a standardized structured intervention overcomes a widely discussed limitation of the meditation literature, which is the great heterogeneity of the meditation practices included in the same investigation (Chiesa and Malinowski 2011). Indeed, both ancient Buddhist meditation, such as Vipassana, Zen, and transcendental meditation, and modern mindfulness-based intervention, such as the MBSR (Kabat-Zinn 1990) and mindfulness-based cognitive therapy (MBCT; Segal et al. 2002), are commonly referred to with the general term of “mindfulness meditation practice”; however, they largely differ in the way mindfulness is conceptualized and practiced (Chiesa and Malinowski 2011). The use of the MBSR program clearly defines the generalizability of the findings, facilitates replication, and makes the comparison among studies more reliable.

Pilates trainees were selected as control participants in order to assess non-MBSR-specific learning effects. The involvement of an active group allowed us to address another important issue, which concerns the choice of an adequate control group. In the meditation literature, the use of an active control group is usually preferred against the widely used passive or waiting list control group, in order to attenuate group biases related to motivational levels (MacCoon et al. 2014). Our control participants were engaged in a regular physical training, that is, the practice of Pilates, during 8 weeks. This training is similar to Hatha Yoga in the

emphasis that is given on performing correct and harmonious movements, which must be healthy and enjoyable, in contrast with body overtraining (Hoffman and Gabel 2015). However, there is not any explicit instruction about achieving a mindful state, characterized by self-regulation of attention on present moment experience, typical of the MBSR training (Kabat-Zinn 1990). Given previous findings, we expected that the MBSR practice, but not the Pilates one, would induce significant changes on behavioral performance and, more importantly, on the associated neural processes. We hypothesized the mindfulness training to modulate some key ERP components, namely, the CNV elicited by cue stimuli, which is linked with an enhanced implementation of proactive cognitive control processes, and the N2 and P3a components elicited by probe stimuli, which are linked with an enhanced implementation of reactive control mechanisms.

## Methods

### Participants

Overall, 29 participants were recruited from an 8-week MBSR training course promoted by a local center conducted by an instructor certified by the Center for Mindfulness in Medicine, Health Care and Society of the University of Massachusetts (Medical School). Twenty-six participants were included in the final statistical analyses, since they completed the training and both pre- and post-training experimental sessions (mean age = 43.6 years, SD = 11.3, range = 26–60 years; mean education = 16.6 years, SD = 2.9; females = 21). Based on the Edinburgh Handedness Inventory (EHI; Oldfield 1971), 23 were right handed (EHI range = 55–100), 2 were mixed right handed (EHI score = 15 and 45, respectively), and 1 was left handed (EHI score = -70).

For the Pilates control group, 25 participants were recruited from training classes conducted by local centers specialized in physical well-being and health care. Twenty-three of them regularly attended the Pilates training for 8 weeks, completed both pre- and post-training experimental sessions, and were included in the final statistical analyses (mean age = 47.2 years, SD = 11.9, range = 28–67; mean education = 15.5, SD = 3.7; females = 20). Based on the EHI questionnaire, 19 were right handed (EHI range = 60–100), 2 were mixed right handed (EHI score = 45 and 50, respectively), and 2 were left handed (EHI score = -70 and -80, respectively). The two groups did not differ in terms of age ( $t(47) = 1.58, p = 0.12$ ), education level ( $t(47) = 1.04, p = 0.30$ ), and handedness ( $t(47) = .05, p = 0.96$ ).

All participants signed consent forms prior to testing and received a reimbursement of 10 euros soon after each experimental session. The study was approved by the local Ethical Committee in agreement with Helsinki declaration.

## Procedure

Both groups underwent the same procedure. Specifically, all participants performed two identical experimental sessions about 8 weeks apart. At the beginning of each experimental session, a 64-channel EEG cap was mounted and participants were instructed about the task. At the end of each session, they were invited to fill in self-report mindfulness questionnaires.

## Measures

Three self-report questionnaires were administered to assess daily subjective experience of mindful states. The questionnaires allowed us to evaluate the level of dispositional mindfulness. We expected an increase of scores on all questionnaires in the MBSR group after the training.

### Mindfulness Attention and Awareness Scale

The Mindfulness Attention and Awareness Scale (MAAS) is a self-report questionnaire developed to assess individual differences in the frequency of mindful states (Brown and Ryan 2003; Veneziani and Voci 2014). The scale is based on a single-factor structure and consists of 15 items that measure the tendency to be attentive and aware of present experience in day-to-day circumstances.

### Cognitive Affective Mindfulness Scale-Revised

The Cognitive and Affective Mindfulness Scale-Revised (CAMS-R) is a self-report questionnaire (12 items) developed to assess “attitudes and approaches toward internal experiences of emotions and thoughts” in the general population (Feldman et al. 2007; Veneziani and Voci 2015). The CAMS-R is based on a broad conceptualization of mindfulness that captures not only a particular type of attention and awareness, such as the MAAS questionnaire, but also attitudinal components related to mindfulness, such as acceptance and nonjudgmental attitude.

### Five Facets Mindfulness Questionnaire

The Five Facets Mindfulness Questionnaire (FFMQ) consists of 39 items that assess five scales of mindfulness: (1) nonreactivity to inner experience (Nonreacting), (2) observing/attending/noticing to sensations/feelings/thoughts (Observe), (3) acting with awareness/automatic pilot/concentration/nondistracted (Acting with awareness), (4) describing/labeling with words (Describe), and (5) nonjudging of experience (Nonjudging) (Baer et al. 2006; Giovannini et al. 2014).

## The AX-Continuous Performance Task

The task consisted in the presentation of pairs of letters (i.e., a cue and a probe). The letters were delivered one at a time. Participants were asked to respond to each letter by pressing one of two buttons (“z” or “m”), with the index finger of either hand. They were instructed that the target pair was “AX” and that the response to this target requires to press the left button with the left index finger at the onset of the first letter (i.e., the “A”), and to press the right button with the right index finger at the onset of the second letter (i.e., the “X”). To any other pair of letters, they were asked to respond by pressing the left button with the left index finger at the onset of the first letter and by pressing again the left button at the onset of the second letter. Left and right response keys were counterbalanced across participants. Overall, four trial types (i.e., pairs of letters) were included in the task: the “AX” trials, the A–nonX trials (“AY”), the nonA–X trials (“BX”), and nonA–nonX trials (“BY”). In contrast to traditional AX-CPT, an equal number of A trials and B trials were delivered (Richmond et al. 2015). Specifically, the proportion of trials was as follows: 37.5% AX, 12.5% AY, 12.5% BX, and 37.5% BY. Within a block, the frequency of A- and B-cues was equated (50/50 vs. 80/20 in the traditional version), allowing us to control for cue validity across trials (here the 75% of A-cues anticipated an X-probe, and 75% of B-cues anticipated a Y-probe) and for potential confounding factors related to the infrequency/novelty of B-cues. As a consequence, the frequency of each probe type was the same as well (i.e., 50% of trials contained an X-probe, and 50% of trials contained a Y-probe). In that way, a greater emphasis was placed on cue information (see Richmond et al. 2015). No information about probability of AX pairs was provided; therefore, participants were not induced to implement a specific proactive strategy.

All stimuli were black (about  $2 \times 2.3$  cm) and centrally displayed on a silver background, on a 19-in. screen (1024  $\times$  768 pixel resolution). The task was performed in an acoustically and electrically isolated booth for EEG signal acquisition. The cue letter could be either the letter A (A-cue) or the letters C, D, F, H, M, N, P, U, or O (B-cue); the probe letter could be either the letter X (X-probe) or the letters B, E, G, L, Q, R, S, V, J, or W (Y-probe).

Participants completed a total of 6 blocks that lasted about 5 min each. Each block contained a sequence of 64 trials, comprising 24 AX, 8 AY, 8 BX, and 24 BY trials presented in random order. The stimulus timing was as follows: both the cue and the probe letter were on the screen for 300 ms and a fixation cross was displayed for 1700 ms between the two. From the probe offset to the successive cue, a jittered intertrial interval (ITI) randomly changing from 2000 to 3000 ms was inserted, during which a string of three fixation crosses was displayed. Rest breaks were offered at the end of each block.

Before starting the experimental task, participants performed a short practice session (15 trials) in order to familiarize with it. If less than 11 responses were correct, they repeated the practice (overall, this was the case only for two participants that repeated the practice once). Stimulus presentation and response collection were controlled by E-Prime 2.0 software (Schneider et al. 2002).

### EEG Recording and Preprocessing

The EEG signal was recorded using an EEG portable system (Brain Products, Munich, Germany), connected to 64 sintered Ag/AgCl ring electrodes mounted on an elastic cap (EASYCAP GmbH, Germany), according to the extended 10–20 system (Jasper, 1958). Reference and ground channels were located over FCz and AFz, respectively. Impedance values were kept below 5 k $\Omega$ . Raw data were band-pass-filtered between 0.016 and 250 Hz and digitized at a 500-Hz sampling rate.

The data preprocessing was performed using EEGLAB (v. 13.4.4b, Delorme and Makeig 2004), an open source toolbox for EEG analysis implemented in MATLAB (v. R2017b). Firstly, a low-pass filter was applied, using the windowed sinc FIR filter method (Kaiser window type), with a cutoff frequency of 35 Hz and a transition bandwidth of 10 Hz (Widmann et al. 2015). Then, a high-pass filter was applied (cutoff frequency = 0.125 Hz, transition bandwidth = 0.25 Hz), in order to remove slow drifts. The continuous EEG signal was initially segmented in epochs ranging from 1500 ms before to 5300 ms after cue onset. An independent component analysis (ICA) was run, based on an extended Infomax algorithm. The resulting ICA components were visually scrutinized to identify and remove ocular movements and muscle artifacts. For the cue-locked ERP analysis, epochs were segmented from 500 ms before to 3000 ms after cue onset and the signal was baseline-corrected using the average amplitude in the 500-ms time window before cue onset. For the probe-locked ERP analysis, epochs were segmented from 200 ms before to 1000 after probe onset and the signal was baseline-corrected using the 200-ms time window before probe onset.

The epochs associated with missing and incorrect responses were removed. An automatic detection of artifactual and/or outlier epochs was performed, by applying four criteria (see Delorme et al. 2007): (i)  $-100/+100$   $\mu$ V as extreme amplitude values threshold; (ii) drifts larger than  $\pm 50$   $\mu$ V and  $R^2 > 0.3$  for the linear trend test; (iii)  $SD > 7$  (for each channel) and  $SD > 4$  (for all channels) from the mean probability distribution of amplitude values across epochs for the improbability test; and (iv)  $SD > 7$  (for each channel) and  $SD > 4$  (for all channels) from the kurtosis of the probability distribution. Epochs containing data points exceeding these criteria were excluded from further analyses. Table 1 summarizes the

number of epochs included in the analyses, for each condition. Finally, data were re-referenced to the average of all electrodes, with the exception of the vertical EOG channel.

### Data Analysis

#### Questionnaire Data

A 2 (session: pre- vs. post-training)  $\times$  2 (group: MBSR vs. control) mixed ANOVA was performed for each questionnaire score. Least significant difference (LSD) post hoc tests were used in order to interpret significant effects and partial  $\eta^2$  to estimate effect size.

#### Behavioral Data

Error rates and response times (RTs) were examined. Trials containing omissions or anticipations ( $RTs \leq 100$  ms), either to the cue or to the probe, or incorrect responses to the cue, were removed. Error rates were computed in terms of incorrect responses to probes (i.e., false alarms). In order to adjust for floor effects, a correction was applied to error rates equal to 0, which were replaced by  $0.5/(\text{total number of trials} + 1)$ . The total number of trials was  $n = 144$  for the AX and BY trials and  $n = 48$  for the AY and BX trials. Raw RTs were log-transformed to improve normality. Then, RTs exceeding 3 standard deviations above the mean, computed for each participant and condition separately, were removed. A 2 (session: pre- vs. post-training, within-subjects factor)  $\times$  4 (trial type: AX, AY, BX, and BY, within-subjects factor)  $\times$  2 (group: MBSR vs. control, between-subjects factor) mixed ANOVA was performed on RTs. LSD post hoc tests were used in order to clarify significant effects and partial  $\eta^2$  to estimate effect size.

To estimate the amount of proactive control, additional behavioral indices were computed by comparing performance across trial types, specifically,  $d'$ -context, A-cue bias, and proactive behavioral index (PBI; Barch et al. 2001; Braver et al. 2009; Gonthier et al. 2016; Richmond et al. 2015). The first two indices,  $d'$ -context and A-cue bias, correspond to  $d'$  and  $c$  criterion, respectively, as described in the Signal Detection Theory (Stanislaw and Todorov 1999). The  $d'$ -context index was calculated as the difference between hits on AX trials and false alarms on BX trials. Hits were calculated as the proportion of correct trials (correct response to both cue and probe). A correction was applied when hits matched total trials, as follows:  $(\text{total number of trials} - 0.5)/(\text{total number of trials} + 1)$ . Hit and false alarm values were then  $z$ -transformed. Therefore, the  $d'$ -context was computed as follows:  $Z$  (hits on AX trials)  $- Z$  (false alarms on BX trials). This measure reflects the ability of participants to use contextual information from the cue to drive the response to the probe. The A-cue bias was calculated on hits on AX trials and false alarms on AY trials, as follows:  $Z$  (correct hits on AX trials)  $+ Z$  (errors

**Table 1** Mean and standard deviation (in brackets) of the number of event-related potential (ERP) epochs after artifact rejection and removal of missing and incorrect responses

	MBSR group				Control group			
	AX tot = 144	AY tot = 48	BX tot = 48	BY tot = 144	AX tot = 144	AY tot = 48	BX tot = 48	BY tot = 144
Pre-training	136.6 (4.9)	44.3 (2.8)	45.0 (2.6)	137.9 (5.7)	140.3 (5.1)	46.7 (1.5)	46.6 (1.9)	141.8 (5.6)
Post-training	137.6 (4.2)	45.7 (2.2)	45.6 (1.7)	139.2 (2.8)	138.8 (2.7)	46.0 (55.0)	45.4 (1.4)	139.2 (1.4)

on AY trials)/2. This index represents the tendency to make a target response after an A-cue regardless of the probe identity. The PBI was calculated both in terms of error rates and RTs, as  $(AY - BX)/(AY + BX)$ . A positive PBI is given by more errors/longer RTs on AY trials compared to BX trials and reflects a higher proactive control engagement; a negative PBI value is given by more errors/longer RTs on BX trials compared to AY trials and reflects a higher reactive control engagement.

The statistical analysis on error rates and proactive control indices was performed by means of nonparametric tests, since data distribution did not meet the normality assumption, as tested via the Shapiro–Wilk normality test (all  $ps < 0.05$ ). The Wilcoxon signed rank test for paired data was applied on each trial type in order to assess the pre-/post-training effects in each group and session, separately. The Mann–Whitney tests for unpaired data were performed to assess group differences in each session and trial type, separately. Since multiple tests were performed, false discovery rate (FDR) correction was applied to correct  $p$  values (Benjamini and Hochberg 1995).

### Event-Related Potential Data

One participant from each group was excluded from the ERP analyses due to insufficient data quality (excessive eye movements and muscular artifacts, which could not be reliably removed by the ICA algorithm and the automatic artifact rejection procedure). The final sample consisted of 25 participants in the MBSR group and 22 participants in the control group.

First of all, the amplitude of potentials evoked by the cue letters (cue-locked ERPs) was analyzed to investigate pre-/post-training effects. Specifically, the CNV amplitude was examined in a 500-ms time window immediately preceding the probe onset, over midline fronto-central electrode sites (FCz and Fz collapsed), where it was expected to reach the maximum amplitude (Boksem et al. 2006; Dias et al. 2003; Ng et al. 2011; Tarantino et al. 2010). The mean amplitude over a time window of 500 ms preceding the probe onset was extracted for each trial. Trials containing A-cue (AX, AY) and trials containing B-cue (BX, BY) were then averaged. The mean CNV amplitude values were submitted to a 2 (session:

pre- vs. post-training)  $\times$  2 (cue type: A-cue vs. B-cue)  $\times$  2 (group: MBSR vs. control) mixed ANOVA.

Secondly, the amplitude of potentials evoked by the probe letters (probe-locked ERPs) was analyzed. Specifically, the anterior N2 and P3a components were examined. The N2 is a fronto-central negative potential, peaking around 200–300 ms after a target stimulus; it reflects conflict monitoring and is especially pronounced in tasks requiring high cognitive control costs, such as the Stroop task (Donkers and Van Boxtel 2004; Eimer 1993; Folstein and Van Petten 2008; Larson et al. 2014). The P3a is a positive deflection characterized by a fronto-central scalp distribution, peaking around 300–500 ms post-stimulus; it is mainly involved in context-updating and attentional resource allocation (Hämmerer et al. 2010; Polich 2007). We extracted mean ERP amplitude over a 200–300-ms time window after probe onset and over a 300–400-ms time window after probe onset, in order to examine the anterior N2 and P3a components, respectively. Both the N2 and the P3a amplitudes were extracted from midline centro-frontal electrode sites (FCz, Cz), where they are maximally expressed. Mean amplitude values were submitted to a 4 (trial type: AX, AY, BX, BY)  $\times$  2 (session: pre- vs. post-training)  $\times$  2 (group: MBSR vs. control) mixed ANOVA.

The Spearman correlation analysis was also performed, on the whole data sample, between ERP component's amplitude, behavioral data, AX-CPT indices, and questionnaire scores in order to explore possible relationships between these measures. The FDR correction was applied to correct for multiple comparisons.

All statistical analyses were performed using the IBM SPSS software (release 23).

## Results

### Mindfulness Questionnaires

Tables 2 and 3 show mean scores for each self-report mindfulness questionnaire and the relative inferential statistics testing the effect of experimental session and group. All questionnaire scores but the CAMS showed a significant group  $\times$  session interaction. The post hoc tests revealed significantly higher MAAS and FFMQ scores in the post- compared to the

**Table 2** Mean and standard deviation (in brackets) of the mean scores on the Mindfulness Attention Awareness Scale (MAAS), on the Cognitive and Affective Mindfulness Scale-Revised (CAMS-R), on the Five Facets Mindfulness Questionnaire (FFMQ), and on its five subscales: Nonreactivity (FFMQ-NR), Observing (FFMQ-O), Acting with awareness (FFMQ-A), Describing (FFMQ-D), and Nonjudging (FFMQ-NJ)

	MBSR group		Control group	
	Pre-training	Post-training	Pre-training	Post-training
MAAS	3.46 (0.86)	4.03 (0.67)	4.20 (0.53)	4.17 (0.51)
CAMS-R	2.54 (0.29)	2.65 (0.31)	2.76 (0.31)	2.76 (0.29)
FFMQ-total	3.06 (0.66)	3.53 (0.44)	3.54 (0.40)	3.54 (0.46)
FFMQ-NR	2.66 (0.73)	3.18 (0.48)	3.00 (0.60)	2.92 (0.58)
FFMQ-O	3.00 (0.82)	3.60 (0.50)	3.57 (0.70)	3.64 (0.67)
FFMQ-A	2.87 (0.67)	3.40 (0.62)	3.78 (0.68)	3.86 (0.52)
FFMQ-D	3.33 (0.73)	3.63 (0.84)	3.73 (0.88)	3.68 (0.94)
FFMQ-NJ	3.24 (0.92)	3.83 (0.72)	3.44 (0.80)	3.55 (0.79)

pre-training session for the MBSR group [ $t(24) = 4.40$ ,  $p < 0.001$ , Cohen's  $d = 0.87$ , and  $t(24) = 5.92$ ,  $p < 0.001$ , Cohen's  $d = 0.94$ , respectively]. The analysis of each FFMQ subscale revealed a significant score increase in the MBSR group only, for all dimensions. No training effects were observed in the CAMS-R scores (all  $ps > 0.11$ ).

## Behavioral Results

Figure 1 illustrates the percentage of error rates for each trial type before and after the training. Based on the Wilcoxon signed rank test for paired data, only participants who underwent the MBSR training exhibited significantly fewer errors on AY trials [ $\text{mean}_{\text{pre}} = 4.82\%$  (SD = 4.07),  $\text{mean}_{\text{post}} = 2.67\%$  (SD = 2.21);  $Z = -3.16$ ,  $p = 0.002$ ], and on BX trials [ $\text{mean}_{\text{pre}} = 2.59\%$  (SD =

2.16),  $\text{mean}_{\text{post}} = 1.31\%$  (SD = 0.69);  $Z = -2.77$ ,  $p = 0.006$ ] after the training. No training effect was observed in the control group. However, based on the Mann–Whitney tests, the control group exhibited significantly lower error rates on AY trials both before and after the training compared to the MBSR group ( $Z = 2.78$ ,  $p = 0.005$  and  $Z = 2.54$ ,  $p = 0.011$ , respectively).

Figure 2 illustrates mean RTs for each trial type, before and after the training. As expected, a main effect of trial type [ $F(1, 47) = 109.06$ ,  $p < 0.0001$ , partial  $\eta^2 = 0.70$ ] was found, which revealed that RTs significantly differed across trial types ( $\text{RT}_{\text{AY}} > \text{RT}_{\text{AX}} > \text{RT}_{\text{BX}} > \text{RT}_{\text{BY}}$ , all  $ps < 0.03$ ). Moreover, significant session  $\times$  group [ $F(1, 47) = 5.93$ ,  $p = 0.018$ , partial  $\eta^2 = 0.11$ ] and session  $\times$  trial type interactions [ $F(1, 47) = 4.32$ ,  $p = 0.006$ , partial  $\eta^2 = 0.08$ ] emerged. Post hoc tests showed shorter RTs in the post- compared to pre-training session for the control group ( $p = 0.015$ ). Additionally, irrespective of group, participants were faster in responding to trials containing B-cues in post- compared to pre-training session (all  $ps < 0.0001$ ), whereas no session effect was found for the A-cue trials (all  $ps > 0.077$ ).

Wilcoxon signed rank tests were performed on behavioral indices, namely, the  $d'$ -context, the A-cue bias, and the PBI (both on errors and on log-RTs). Table 4 summarizes the mean scores for each behavioral index, as a function of group and experimental session. First of all, the effect of training was assessed in each group separately. Statistical analyses showed more positive PBI-RT values in the post-training session for the control group ( $Z = 2.98$ ,  $p = 0.003$ ). All other comparisons did not survive the FDR correction. Group differences on indices, in both pre- and post-training sessions, were submitted to multiple Mann–Whitney tests. No comparison survived the significance correction.

In order to explore possible relationships between behavioral data and questionnaire scores, nonparametric Spearman correlation analyses were performed, regardless of the group.


**Table 3** Inferential statistics overview for each self-report mindfulness questionnaire score (average). We report the effects for session factor (pre- vs. post-8-week training), group factor (MBSR vs. control group),

and session by group interaction. Only the significant post hoc tests were reported. They showed an increase in the questionnaire scores (namely a training effect) in the MBSR group

	Session			Group			Session $\times$ group interaction			Post hoc tests
	$F$	$p$	Partial $\eta^2$	$F$	$p$	Partial $\eta^2$	$F$	$p$	Partial $\eta^2$	
MAAS	8.63	0.005	0.17	6.06	0.018	0.13	18.22	< 0.001	0.30	MBSR <sub>post</sub> > MBSR <sub>pre</sub>
CAMS-R	2.22	ns	0.05	2.66	ns	0.06	0.89	ns	0.02	–
FFMQ-total	12.71	< 0.001	0.23	2.86	ns	0.06	12.32	< 0.001	0.22	MBSR <sub>post</sub> > MBSR <sub>pre</sub>
FFMQ-NR	6.65	0.013	0.14	0.08	ns	0.001	9.1	0.004	0.18	MBSR <sub>post</sub> > MBSR <sub>pre</sub>
FFMQ-O	10.46	0.002	0.20	2.52	ns	0.06	11.54	0.001	0.21	MBSR <sub>post</sub> > MBSR <sub>pre</sub>
FFMQ-A	17.7*	< 0.001	0.31	13.70	< 0.001	0.24	7.30	0.009	0.15	MBSR <sub>post</sub> > MBSR <sub>pre</sub>
FFMQ-D	4.68	0.036	0.10	1.74	ns	0.04	6.24	0.016	0.13	MBSR <sub>post</sub> > MBSR <sub>pre</sub>
FFMQ-NJ	12.33	0.001	0.23	0.09	ns	0.002	5.00	0.03	0.11	MBSR <sub>post</sub> > MBSR <sub>pre</sub>

ns =  $p > 0.05$

**Fig. 1** Mean percentage of error rates for each trial type, depending on the pre-training (dark gray) and post-training (light gray) sessions. Error bars represent standard errors of the mean. Asterisks indicate significant differences ( $p < 0.05$ ). The MBSR group is represented on the left panel and the control group on the right one



In terms of error rates, in the pre-training session, a negative correlation between MAAS score and errors on AY trials [ $\rho(47) = -0.399$ ,  $p = 0.005$ ] and a negative correlation between CAMS score and error on AY trials [ $\rho(47) = -0.423$ ,  $p = 0.002$ ] emerged. This means that participants with higher values of “mindfulness,” as measured by the self-report questionnaires, exhibited a higher accuracy on AY trials. Neither the analysis on RTs nor the analysis on behavioral indices yielded significant results (according to the FDR correction). In the post-training session, the negative correlation between the MAAS score and the AY error rate was maintained [ $\rho(47) = -0.535$ ,  $p < 0.001$ ]. Moreover, a negative correlation between the MAAS score and the PBI, both in terms of accuracy [ $\rho(47) = -0.536$ ,  $p < 0.001$ ] and RTs [ $\rho(47) = -0.450$ ,  $p < 0.002$ ] emerged. This means that participants with a higher “mindfulness” score on the self-report questionnaire exhibited higher accuracy on AY trials both in terms of errors and RTs; in other words, in the post-training session, AY trials exerted less interference in individuals with higher mindfulness.


### ERP Data: Cue-Locked Analysis

We first computed cue-locked analysis in order to evaluate mindfulness-related effects on proactive control processes. Figure 3 shows ERP waveforms locked to A-cue (a) and B-cue (b) The amplitude maps represent the topographical distribution of the post-pre amplitude difference during the 500-

ms interval preceding the probe onset, in the MBSR (c) and in the control (d) group. As expected, a slow negative potential developed after the cue onset (CNV), which increased in amplitude throughout the cue-to-probe delay, in anticipation of the probe, and reached maximal (negative) amplitude values over fronto-central sites. As one can see, the CNV evoked after A-cues was steeper in post- compared to pre-training session, in the MBSR group only. This effect was not observed in the control group.

The 2 (cue type)  $\times$  2 (session)  $\times$  2 (group) ANOVA contrasting the CNV amplitude in A-cue and B-cue trials during the 500-ms time window preceding probe onset, over Fz and FCz electrodes, yielded a significant main effect of cue type [ $F(1, 45) = 66.47$ ,  $p < 0.0001$ , partial  $\eta^2 = 0.59$ ]. This effect demonstrated that the CNV amplitude after A-cue was larger than after B-cue, irrespective of group and session. Neither a main effect of session nor a main effect of group was observed [ $F(1, 45) = 2.59$ ,  $p = 0.115$ ;  $F(1, 045) = 0.825$ ,  $p = 0.369$ , respectively]. A significant session  $\times$  group interaction [ $F(1, 45) = 7.11$ ,  $p = 0.011$ , partial  $\eta^2 = 0.14$ ] and a significant cue type  $\times$  session interaction [ $F(1, 45) = 6.64$ ,  $p = 0.013$ , partial  $\eta^2 = 0.13$ ] were found. The post hoc analysis of the session  $\times$  group interaction revealed that only the MBSR group showed a training effect, namely, a more negative CNV amplitude in the post-training session [mean<sub>pre</sub> =  $-1.28$   $\mu\text{V}$  (SD = 1.50); mean<sub>post</sub> =  $-1.60$   $\mu\text{V}$  (SD = 2.15);  $p = 0.003$ ], whereas the control group did not [mean<sub>pre</sub> =  $-1.25$   $\mu\text{V}$  (SD = 1.09);

**Fig. 2** Mean response times (RTs, in ms) for each trial type, depending on the pre-training (dark gray) and post-training (light gray) sessions. Error bars represent standard errors of the mean. The MBSR group is represented on the left panel and the control group on the right one






**Table 4** Mean and standard deviation (in brackets) for the AX-CPT measures, as a function of group and experimental session

	MBSR group		Control group	
	Pre-training	Post-training	Pre-training	Post-training
$d'$ -context	4.19 (0.47)	4.43 (0.32)	4.37 (0.32)	4.46 (0.38)
A-cue bias	0.16 (0.15)	0.09 (0.18)	0.05 (0.14)	0.040 (0.10)
PBI-errors	0.22 (0.47)	0.19 (0.38)	0.09 (0.39)	0.01 (0.31)
PBI-RTs	0.02 (0.02)	0.02 (0.02)	0.02 (0.16)	0.03 (0.02)

mean<sub>post</sub> = -1.17  $\mu$ V (SD = 1.10);  $p = 0.472$ ]. The post hoc analyses of the cue type  $\times$  session interaction revealed that only the CNV amplitude in the A-cue trials was affected by the training. Namely, only the CNV evoked by the A-cue was

larger (more negative) in the post-training session. A marginally significant three-way interaction between cue type  $\times$  group  $\times$  session [ $F(1, 45) = 4.03$ ,  $p = 0.051$ , partial  $\eta^2 = 0.08$ ] suggested that the MBSR training affected CNV amplitude in A-cue trials only, namely, the amplitude of the CNV showed more negative values in post- compared to pre-training session in the MBSR group in A-cue trials ( $p = 0.001$ ), whereas it did not differ in all other comparisons (all  $ps > 0.159$ ).

In order to understand the behavioral counterpart of the CNV, Spearman correlation analyses with behavioral data were performed. Specifically, the CNV amplitude was correlated with error rates, RTs, behavioral indices, and questionnaire scores, on the whole participants' sample, for each trial type and for pre- and post-training sessions, separately. In the pre-training session, the CNV amplitude evoked by A-cues



**Fig. 3** In the left panels, grand average event-related potential (ERP) waveforms in A-cue (a) and B-cue (b) trials are depicted. ERPs are time-locked to the cue onset (0 ms); probe onset is at 2000 ms. FCz and Fz electrodes are collapsed. A negative component (the contingent negative variation, CNV) emerges in anticipation of probe onset, in A-cue trials only. The shaded area represents the standard error of the mean. In

the right panels, topographical maps of the post- vs. pre-training difference in ERP amplitude ( $\mu$ V) in the MBSR group (c) and the control group (d) are represented. This amplitude difference was extracted from 1500 to 2000 ms relative to A-cues onset. The fronto-central distribution of the post-pre amplitude difference can be appreciated in the MBSR group only

was positively correlated with RTs on all trial types [AX:  $\rho(45) = 0.404$ ,  $p = 0.005$ ; AY:  $\rho(45) = 0.382$ ,  $p = 0.008$ ; BX:  $\rho(45) = 0.430$ ,  $p = 0.003$ ; BY:  $\rho(45) = 0.410$ ,  $p = 0.004$ ]. To be noted that the CNV is a negative component, thus the positive correlation means that a more negative CNV amplitude is associated to shorter RTs. Significant correlations did not emerge between the CNV on B-cue trials and RTs ( $p = 0.286$ ). Interestingly, a significant negative correlation between A-cue-locked CNV and the A-cue bias emerged [ $\rho(45) = -0.405$ ,  $p = 0.005$ ]. Furthermore, the CNV amplitude on A trials negatively correlated with PBI index, both in terms of error rates [ $\rho(45) = -0.419$ ,  $p = 0.003$ ] and RTs [ $\rho(45) = -0.351$ ,  $p = 0.016$ ].

Similarly, in the post-training session, the CNV amplitude evoked by A-cues was positively correlated with RTs on AX trials [ $\rho(45) = 0.405$ ,  $p = 0.005$ ], BX trials [ $\rho(45) = 0.388$ ,  $p = 0.007$ ], and BY trials [ $\rho(45) = 0.407$ ,  $p = 0.005$ ]. Negative correlations were also found with PBI index [PBI-errors:  $\rho(45) = -0.550$ ,  $p < 0.001$ ; PBI-RTs:  $\rho(45) = -0.373$ ,  $p = 0.010$ ]. All these correlations survived FDR correction for multiple comparisons. No significant correlations were found with questionnaire scores.

### ERP Data: Probe-Locked Analysis

Figure 4 represents the grand average of ERPs evoked by the probe letter in each trial type, over FCz–Cz electrodes (collapsed). Mean amplitude was extracted in the 200–300-ms time window for the analysis of the N2 component and in the 300–400-ms time window for the analysis of the P3a component. The 4 (trial: AX, AY, BX, BY)  $\times$  2 (session: pre-, post-training)  $\times$  2 (group: MBSR, control) mixed ANOVA revealed a main effect of trial type on N2 amplitude [ $F(3, 45) = 50.07$ ,  $p < 0.0001$ , partial  $\eta^2 = 0.52$ ], which was less pronounced (i.e., more positive) on AX compared to AY, BX, and BY trials (all  $ps < 0.0001$ ), and on AY and BX trials compared to BY trials (both  $ps < 0.001$ ). We did not observe either a main effect of group [ $F(1, 45) = 0.12$ ,  $p = 0.733$ ] or a main effect of session [ $F(1, 45) = 0.34$ ,  $p = 0.563$ ]. A significant session  $\times$  group interaction [ $F(1, 45) = 9.45$ ,  $p = 0.003$ , partial  $\eta^2 = 0.17$ ] revealed less pronounced N2 amplitude in the post- compared to the pre-training session in the MBSR group [mean<sub>pre</sub> = 0.16  $\mu$ V (SD = 1.26); mean<sub>post</sub> = 0.59  $\mu$ V (SD = 1.48);  $p = 0.010$ ] and not in the control group [mean<sub>pre</sub> = 0.33  $\mu$ V (SD = 0.92); mean<sub>post</sub> = 0.04  $\mu$ V (SD = 0.98);  $p = 0.094$ ].


The 4 (trial: AX, AY, BX, BY)  $\times$  2 (session: pre-, post-training)  $\times$  2 (group: MBSR, control) mixed ANOVA on P3a amplitude revealed a main effect of trial [ $F(3, 45) = 73.43$ ,  $p < 0.0001$ , partial  $\eta^2 = 0.62$ ], which reflected a larger amplitude on AY trials compared to AX, BX, and BY trials (all  $ps < 0.011$ ) and on AX trials compared to BX and BY trials (all  $ps < 0.0001$ ). Moreover, a significant main effect of session was found [ $F(1, 45) = 6.87$ ,  $p = 0.012$ , partial  $\eta^2 = 0.13$ ].

Irrespective of group and trial type, the P3a component in the post-training session exhibited larger amplitude compared to the pre-training session [mean<sub>pre</sub> = 1.62  $\mu$ V (SD = 1.81); mean<sub>post</sub> = 2.07  $\mu$ V (SD = 2.13)]. On the other hand, a main effect of group did not emerge [ $F(1, 45) = 3.31$ ,  $p = 0.075$ ]. A marginally significant session  $\times$  group interaction [ $F(1, 45) = 4.05$ ,  $p = 0.050$ , partial  $\eta^2 = 0.08$ ] denoted a more positive P3a amplitude in the post- compared to pre-training session in the MBSR group [mean<sub>pre</sub> = 2.05  $\mu$ V (SD = 1.90); mean<sub>post</sub> = 2.84  $\mu$ V (SD = 2.26);  $p = 0.001$ ], but not in the control group [mean<sub>pre</sub> = 1.19  $\mu$ V (SD = 1.89); mean<sub>post</sub> = 1.29  $\mu$ V (SD = 1.98);  $p = 0.677$ ].

The correlation analysis performed between N2 amplitude and behavioral and questionnaire indices did not yield significant results. On the other hand, the P3a amplitude negatively correlated with RTs, in all trial types and in both pre- [AX:  $\rho(45) = -0.434$ ,  $p = 0.002$ ; AY:  $\rho(45) = -0.308$ ,  $p = 0.035$ ; BX:  $\rho(45) = -0.381$ ,  $p = 0.008$ ; BY:  $\rho(45) = -0.453$ ,  $p = 0.001$ ] and post-training sessions [AX:  $\rho(45) = -0.366$ ,  $p = 0.011$ ; AY:  $\rho(45) = -0.303$ ,  $p = 0.038$ ; BX:  $\rho(45) = -0.414$ ,  $p = 0.004$ ; BY:  $\rho(45) = -0.416$ ,  $p = 0.004$ ]. This is in line with the traditional literature (Polich 2007). Interestingly, the P3a amplitude in BX trials positively correlated with the PBI index in pre- [ $\rho(45) = 0.440$ ,  $p = 0.002$ ] and post-training sessions [ $\rho(45) = 0.425$ ,  $p = 0.003$ ]. Moreover, the P3a amplitude in AX trials significantly correlated with the A-cue bias [ $\rho(45) = 0.347$ ,  $p = 0.017$ ]. This means that also probe-related neural activity was associated with behavioral indices that have been usually interpreted as proactive.

### Discussion

The present study was aimed to elucidate the effects of mindfulness-based meditation training (MBSR) on cognitive control processes, within the theoretical framework of the Dual Mechanism of Cognitive Control (Braver et al. 2007). In particular, we investigated the potential effects of an 8-week training on the dynamic interplay between two functionally distinct control systems, the proactive and the reactive system, by complementing behavioral data with electrophysiological measures. Based on previous evidence on cognitive benefits of meditation practice (e.g., Lutz et al. 2008, 2009; MacLean et al. 2010; Moore et al. 2012; Slagter et al. 2007), we hypothesized that the training would enhance the participants' ability to implement proactive cognitive control, which allows to actively maintain contextual information prior to the occurrence of target events and to strategically allocate attention in order to improve the target response. Additionally, we explored the training effect on reactive cognitive control, which is instead intended as a past-oriented mechanism, activated *just-in-time* by target stimuli to detect and solve interference in unexpected stimuli. To this aim, a modified version of the traditional AX-CPT paradigm was



**Fig. 4** Grand average of the ERPs time-locked to the probe (0 ms). All trial types are represented, namely clockwise from the top-left: AX (a), AY (b), BX (c), and BY (d) trials. The standard error of the mean is represented as shaded error bars. ERPs in the two groups and the two

sessions are represented in different colors. FCz and Cz electrodes are collapsed. The N2 mean amplitude was extracted from the 200- to 300-ms post-probe time window, whereas the P3a mean amplitude was extracted from the 300- to 400-ms post-probe time window

administered to a group of MBSR trainees and to a control group of Pilates trainees (Gonthier et al. 2016; Richmond et al. 2015). Both behavioral and ERP data were examined. The main findings consisted in a reduction of errors specifically on more conflicting trials (i.e., AY and BX trials) and in an amplitude increase of anticipatory and preparatory ERP correlates (i.e., the CNV), in the post-training session of the MBSR group only, as discussed in detail below. An amplitude modulation of reactive ERP components (i.e., the probe-locked N2 and P3a) was also observed in the post-training session of the MBSR group.

Compared to pre-training session, after the 8-week mindfulness training, participants exhibited a significant reduction of error percentages, both on AY and BX trial types. This specific accuracy enhancement might be interpreted as reflecting a more efficient balance between proactive and reactive control processes (Gonthier et al. 2016). Indeed, relative to other trial types, AY and BX trials are associated with higher proactive and reactive control costs. Given the

probability distribution of trials, the occurrence of a Y-probe after an A-cue (AY trials) is unexpected; therefore, it requires a prompt correction of an already prepared response. On the other hand, the occurrence of an X-probe after a B-cue (BX trials) might activate a wrong target response if participants had not prepared in advance the response and had not maintained the cue context. Therefore, higher accuracy on AY and BX trials reflects an efficient adaptation of proactive and reactive control processes. In brief, from a behavioral point of view, the MBSR group showed a more efficient balance between the proactive and reactive control processes post-training, which facilitates a flexible adaptation to environmental requirements.

Control participants exhibited an overall reduction of RTs in the post-training session. When examining the PBI index calculated on RTs, a significant improvement in the post-training session relative to pre-training session emerged in the control group. A closer look at the data revealed that this

effect was driven by a significant RT decrease on BX trials, rather than on AY trials. This effect might be interpreted as reflecting a diminished interference of task-relevant but non-target stimuli, that is, improved reactive control processes in the Pilates group.

In order to better understand the neurofunctional origin of this pattern of behavioral results, we analyzed the simultaneous brain activity monitored through EEG. A significantly larger negative potential was evoked by the cue stimuli (i.e., more negative CNV amplitude) in the MBSR group during post-training session relative to pre-training session (see Fig. 3a). The control group did not show CNV changes after the training. As already discussed in the first part of the article, the CNV is a negative deflection that develops between a cue stimulus and an upcoming target; it reflects central processes of anticipatory attention and response preparation (Boksem et al. 2006; Gaillard 1977; Gomez et al. 2003). The more pronounced CNV amplitude exhibited by MBSR participants likely reflected an increased employment of proactive cognitive control processes after the 8-week MBSR training compared to the control group. This result is consistent with the findings reported in the cross-sectional study of Travis et al. (2000, 2002), in which a larger CNV was elicited after a cue in more trained meditators compared to a control group. The authors interpreted the CNV effect as an increase of the arousal level and a better allocation of attentional resources with increasing frequency of meditation practice (Travis et al. 2000).

During mindfulness meditation, one needs to monitor moment-by-moment the focus of attention on an intended object (e.g., the breathing), detecting distractors (e.g., interfering thoughts, sensations, emotions), disengaging attention from them, and (re)directing attention to the original object (Lutz et al. 2008; Malinowski 2013). The present behavioral and EEG findings suggest that this extensive training on focusing and sustaining attention might have contributed in improving higher cognitive control processes, such as proactive control. Moreover, they suggest that attentional control abilities trained during mindfulness training might have generalized to other contexts and tasks, such as in the experimental setting of our study.

The positive correlation observed between A-locked CNV amplitude and RTs revealed that faster responses to probes were preceded by larger CNV. This effect is consistent with a vast literature showing that increased CNV amplitude is associated with performance improvement on speed (e.g., Boksem et al. 2006; Travis et al. 2000). Interestingly, the A-cue-evoked CNV amplitude correlated with RTs on both A trials and B trials. This finding suggests that individuals who relied on anticipatory control strategies on A trials boosted performance on B trials as well. The negative correlation observed between A-locked CNV amplitude and both A-cue bias and PBI (in terms of errors as well as RTs) further supported the evidence that CNV amplitude reflects the preferential implementation of proactive cognitive control processes.

The probe-locked data revealed additional interesting results about mindfulness-related effects on more transient and reactive processes, activated by the probe. In the AX-CPT paradigm, AY trials are expected to exhibit higher N2 amplitude compared to AX, BX, and BY trials, since whenever an A-cue appeared participants were prompted to make a target response (*Go Local Context*), and if a Y-probe is then displayed, a response conflict arises (Dias et al. 2003; Hämmerer et al. 2010; Morales et al. 2015). In the present study, we observed a reduction of the frontal N2 (more positivity) at probe onset in post- compared to pre-training session in the MBSR group only. This N2 amplitude reduction might be the result of facilitated implementation of probe-related control processes and suggests an increased ability to manage response conflict. The N2 amplitude reduction that we observed seems in contrast with the findings of a previous study also investigating the longitudinal effects of MBSR training on attentional functions (Moore et al. 2012). Indeed, the authors found a more negative N2 amplitude after the 8-week training. There are, however, noteworthy differences with respect to our study. In the study of Moore et al. (2012), the verbal Stroop task was used, and the main finding was a post-training increase of posterior N2 amplitude (more negativity) irrespective of stimulus congruency. Therefore, it could be the case that this component did not reflect the implementation of control mechanisms, but rather an increased activation of task-relevant cortical areas (namely, enhanced stimulus processing when attending to the color of the word), as suggested by the authors.

The probe-locked analysis revealed an increase in P3a amplitude in the post- compared to pre-training session, in all trial types. The marginally significant session  $\times$  group interaction suggests that this effect was specific for the MBSR group. Apparently, this result contrasts previous findings in a passive oddball task, showing diminished P3a amplitude (Cahn and Polich 2009). Nevertheless, in AX-CPT paradigms, a larger probe-locked P3a denotes a better inhibition of prepotent responses (Morales et al. 2015). Furthermore, a larger probe-locked P3 amplitude in similar cue-probe paradigms (such as task-switching) has been interpreted as reflecting the reactivation of task context and the resolution of the conflict induced by overlearned S-R mapping (Brydges and Barceló 2018; Tarantino et al. 2016). Taken together, the modulation of the N2 and P3a components in post- compared to pre-training session suggests that the MBSR group implemented more efficient reactive cognitive control mechanisms.

Remarkably, participants in the MBSR training group only exhibited significantly higher scores on the self-report questionnaires in the post-training session, namely on the total FFMQ score and on all the five relative subscales (i.e., Observing, Describing, Act with awareness, Nonjudging, and Nonreacting). The FFMQ increase after the MBSR training suggests a stronger openness to experience, higher emotional intelligence, and self-compassion, as well as a decrease in neuroticism, thought suppression, and alexithymia (Baer

et al. 2006). Additionally, we observed a significant improvement of the total MAAS score, which captures the tendency to be attentive and aware in the day-to-day circumstances. Previous studies found that the MAAS score positively correlates with emotional intelligence and openness to experience and with a number of well-being measures, such as the PANAS (Positive Affect Negative Affect Scale), that is a subjective emotional well-being scale (Brown and Ryan 2003).

Correlational analyses between behavioral indices and questionnaire scores revealed that individuals that reported subjective experience of mindfulness more frequently exhibited lower error rates on AY trials. In addition, a negative correlation between MAAS score and PBI (calculated on both errors and RTs) showed decreased interference of AY trials compared to BX ones in “more mindful” individuals. Overall, in the post-training session, participants with higher self-reported mindfulness scores exhibited a more efficient use of contextual information, with lower costs on more demanding trials (e.g., AY).

It is worth noting the reduction of RTs on BX trials in the Pilates group and the main effect of session on N2 and P3a amplitudes. This evidence suggests that the control group benefitted from training as well, but only in the reactive components of control processes. This is in line with studies showing the favorable influence of regular physical activity on cognitive processing (e.g., Voss et al. 2013).

One of the strengths of this study is that neural changes were examined together with behavioral data, in a highly controlled experimental setting. Furthermore, a structured mindfulness intervention (MBSR) was investigated, characterized by a standard program, conducted worldwide in the same way (see Kabat-Zinn 1990 for an extensive description of the training). This choice was important to reduce the enormous heterogeneity of meditation practices and to improve study replicability. An additional strength of our study consists in the choice of an active control group. This aspect guaranteed that pre- vs. post-training differences could be selectively attributed to mindfulness practice itself, rather than to general training effect, and contributed to overcome a main limitation in the meditation literature, which concerns the inclusion of passive or waiting list control groups (MacCoon et al. 2014; Marcus et al. 2014). Lastly, the inclusion of a group of participants actively engaged and motivated in the Pilates training allowed us to control for possible biases due to differences in expectancies between groups. Indeed, both groups were characterized by (i) the expectation that the training will succeed, (ii) experiential exercises made in group (sharing of experience), and (iii) the emphasis that is given on the benefits of the training on physical and psychological well-being. Importantly, all participants were informed that the aim of the study was to explore the cognitive benefits related to practices addressed to individual psychological and physical well-being, which applies to both trainings. On the other hand, we

should cautiously consider that the Pilates training was not as structured as the MBSR training.

In summary, the study shows that the abilities trained during mindfulness meditation practice (such as sustaining the attentional focus on a chosen object and reporting, whenever necessary, attention back to the object) generalize to higher cognitive functions. Specifically, the results provide behavioral as well as neural evidence that the MBSR training improves the implementation of both proactive and reactive cognitive control processes. This type of training influences anticipatory and sustained processes, which act in preparation to future predictable events. Moreover, it boosts the ability to promptly adjust planned responses in reaction to unexpected events. Overall, the findings suggest that the training may act on neural plasticity and modulate cortical functioning.

## Limitations and Future Directions

Among the limits of our study, we may point out the fact that this was a quasi-experiment, as we did not randomly assign the participants to the two groups (mindfulness vs. Pilates) starting from the same general population, but they were already self-selected for the type of training of their choice before volunteering. This makes it difficult to totally discard the possibility that intrinsic pre-existing differences between the two groups (e.g., personality aspects, motivation, expectations, etc.) might at least partially have influenced the results. Moreover, the relatively low sample sizes and the differences in structure and daily schedule between the two trainings could have confounded the comparison between groups. Future studies should try to overcome these shortcomings.

A better understanding of the cognitive benefits associated with the mindfulness practice would help to further improve its clinical applicability, especially in those populations characterized by control impairments, including some developmental disorders.

**Acknowledgments** We wish to thank all participants for their extremely kind collaboration. We would particularly thank Dr. Franco Cucchio, the mindfulness trainer of *Motus Mundi Centre for Mindfulness*. Motus Mundi is an Italian center membership CommonGood™ of the CFM-Centre of Mindfulness in Medicine, Health Care and Society in Medical School, Massachusetts University (<https://www.motusmundi.it/home>). We also want to thank Ettore Ambrosini for his precious support and his suggestions during the data analysis.

**Author Contributions** FI and VT: designed and executed the study, performed the data analysis, and wrote the paper. CC: collaborated with the design and editing of the final manuscript. AV: supervised all research steps and wrote and revised the manuscript. Francesca Incagli and Vincenza Tarantino share first author credit.

Francesca Incagli and Vincenza Tarantino contributed equally to this work.

**Funding** This work was funded by the European Research Council under the European Union's 7th Framework Program (FP7/2007–2013), Grant Agreement n. 313692 awarded to AV.

## Compliance with Ethical Standards

**Ethical Approval** This study was approved by the Bioethical Committee of the Azienda Ospedaliera di Padova (Prot. N. 2758P) and was conducted according to the guidelines of the Declaration of Helsinki (World Medical Association 2013).

**Conflict of Interest** The authors declare that they have no conflict of interests.

**Informed Consent** Informed consent was obtained from all individual participants included in the study.

## References

- Baer, R. A., Smith, G. T., Hopkins, J., Krietemeyer, J., & Toney, L. (2006). Using self-report assessment methods to explore facets of mindfulness. *Assessment, 13*(1), 27–45. <https://doi.org/10.1177/1073191105283504>.
- Barch, D. M., Carter, C. S., Braver, T. S., Sabb, F. W., MacDonald, A., Noll, D. C., & Cohen, J. D. (2001). Selective deficits in prefrontal cortex function in medication-naïve patients with schizophrenia. *Archives of General Psychiatry, 58*(3), 280. <https://doi.org/10.1001/archpsyc.58.3.280>.
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological), 57*(1), 289–300.
- Boksem, M. A., Meijman, T. F., & Lorist, M. M. (2006). Mental fatigue, motivation and action monitoring. *Biological Psychology, 72*(2), 123–132.
- Braver, T. S., Gray, J. R., & Burgess, G. C. (2007). Explaining the many varieties of working memory variation: dual mechanisms of cognitive control. In A. Conway et al. (Eds.), *Variation in working memory* (pp. 76–106). Oxford: Oxford University Press.
- Braver, T. S., Paxton, J. L., Locke, H. S., & Barch, D. M. (2009). Flexible neural mechanisms of cognitive control within human prefrontal cortex. *Proceedings of the National Academy of Sciences, 106*(18), 7351–7356.
- Braver, T. S. (2012). The variable nature of cognitive control: a dual-mechanisms framework. *Trends in Cognitive Sciences, 16*(2), 106–113. <https://doi.org/10.1016/j.tics.2011.12.010>.
- Brydges, C. R., & Barceló, F. (2018). Functional dissociation of latency-variable, stimulus- and response-locked target P3 sub-components in task-switching. *Frontiers in Human Neuroscience, 12*, 60.
- Brown, K. W., & Ryan, R. M. (2003). The benefits of being present: mindfulness and its role in psychological well-being. *Journal of Personality and Social Psychology, 84*(4), 822–848. <https://doi.org/10.1037/0022-3514.84.4.822>.
- Cahn, B. R., & Polich, J. (2006). Meditation states and traits: EEG, ERP, and neuroimaging studies. *Psychological Bulletin, 132*(2), 180–211. <https://doi.org/10.1037/0033-2909.132.2.180>.
- Cahn, B. R., & Polich, J. (2009). Meditation (Vipassana) and the P3a event-related brain potential. *International Journal of Psychophysiology, 72*(1), 51–60. <https://doi.org/10.1016/j.ijpsycho.2008.03.013>.
- Chang, J.-H., Kuo, C.-Y., Huang, C.-L., & Lin, Y.-C. (2017). The flexible effect of mindfulness on cognitive control. *Mindfulness, 9*(3), 792–800. <https://doi.org/10.1007/s12671-017-0816-9>.
- Chiesa, A., Calati, R., & Serretti, A. (2011). Does mindfulness training improve cognitive abilities? A systematic review of neuropsychological findings. *Clinical Psychology Review, 31*(3), 449–464. <https://doi.org/10.1016/j.cpr.2010.11.003>.
- Chiesa, A., & Malinowski, P. (2011). Mindfulness-based approaches: are they all the same? *Journal of Clinical Psychology, 67*(4), 404–424. <https://doi.org/10.1002/jclp.20776>.
- Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods, 134*(1), 9–21. <https://doi.org/10.1016/j.jneumeth.2003.10.009>.
- Delorme, A., Sejnowski, T., & Makeig, S. (2007). Enhanced detection of artifacts in EEG data using higher-order statistics and independent component analysis. *Neuroimage, 34*(4), 1443–1449.
- Dias, E. C., Foxe, J. J., & Javitt, D. C. (2003). Changing plans: a high density electrical mapping study of cortical control. *Cerebral Cortex, 13*(7), 701–715. <https://doi.org/10.1093/cercor/13.7.701>.
- Donkers, F. C. L., & Van Boxtel, G. J. M. (2004). The N2 in go/no-go tasks reflects conflict monitoring not response inhibition. *Brain and Cognition, 56*(2 SPEC. ISS), 165–176. <https://doi.org/10.1016/j.bandc.2004.04.005>.
- Eimer, M. (1993). Effects of attention and stimulus probability on ERPs in a Go/Nogo task. *Biological Psychology, 35*(2), 123–138. [https://doi.org/10.1016/0301-0511\(93\)90009-W](https://doi.org/10.1016/0301-0511(93)90009-W).
- Farb, N. A. S., Segal, Z. V., Mayberg, H., Bean, J., McKeon, D., Fatima, Z., & Anderson, A. K. (2007). Attending to the present: mindfulness meditation reveals distinct neural modes of self-reference. *Social Cognition and Affective Neuroscience, 2*(4), 313–322. <https://doi.org/10.1093/scan/nsm030>.
- Feldman, G., Hayes, A., Kumar, S., Greeson, J., & Laurenceau, J. P. (2007). Mindfulness and emotion regulation: the development and initial validation of the Cognitive and Affective Mindfulness Scale-Revised (CAMS-R). *Journal of Psychopathology and Behavioral Assessment, 29*(3), 177–190. <https://doi.org/10.1007/s10862-006-9035-8>.
- Folstein, J. R., & Van Petten, C. (2008). Influence of cognitive control and mismatch on the N2 component of the ERP: a review. *Psychophysiology, 45*(1), 152–170. <https://doi.org/10.1111/j.1469-8986.2007.00602.x>.
- Gallant, S. N. (2016). Mindfulness meditation practice and executive functioning: breaking down the benefit. *Consciousness and Cognition, 40*, 116–130. <https://doi.org/10.1016/j.concog.2016.01.005>.
- Giovannini, C., Giromini, L., Bonalume, L., Tagini, A., Lang, M., & Amadei, G. (2014). The Italian five facet mindfulness questionnaire: a contribution to its validity and reliability. *Journal of Psychopathology and Behavioral Assessment, 36*(3), 415–423.
- Gonthier, C., Macnamara, B. N., Chow, M., Conway, A. R. A., & Braver, T. S. (2016). Inducing proactive control shifts in the AX-CPT. *Frontiers in Psychology, 7*(NOV), 1–14. <https://doi.org/10.3389/fpsyg.2016.01822>.
- Hämmerer, D., Li, S. C., Müller, V., & Lindenberger, U. (2010). An electrophysiological study of response conflict processing across the lifespan: assessing the roles of conflict monitoring, cue utilization, response anticipation, and response suppression. *Neuropsychologia, 48*(11), 3305–3316. <https://doi.org/10.1016/j.neuropsychologia.2010.07.014>.
- Hoffman, J., & Gabel, C. P. (2015). The origins of Western mind-body exercise methods. *Physical Therapy Reviews, 20*(5–6), 315–324. <https://doi.org/10.1080/10833196.2015.1125587>.
- Jasper, H. H. (1958). The 10-20 electrode system of the International Federation. *Electroencephalography and Clinical Neurophysiology, 10*, 371–375.

- Kabat-Zinn, J. (1990). *Full catastrophe living: using the wisdom of your body and mind to face stress, pain and illness*. New York: Dell Publishing.
- Larson, M. J., Clayson, P. E., & Clawson, A. (2014). Making sense of all the conflict: a theoretical review and critique of conflict-related ERPs. *International Journal of Psychophysiology*, *93*(3), 283–297.
- Li, Y., Liu, F., Zhang, Q., Liu, X., & Wei, P. (2018). The effect of mindfulness training on proactive and reactive cognitive control. *Frontiers in Psychology*, *9*, 1002. <https://doi.org/10.3389/fpsyg.2018.01002>.
- Lutz, A., Slagter, H. A., Dunne, J. D., & Davidson, R. J. (2008). Attention regulation and monitoring in meditation. *Trends in Cognitive Sciences*, *12*(4), 163–169. <https://doi.org/10.1016/j.tics.2008.01.005>.
- Lutz, A., Slagter, H. A., Rawlings, N. B., Francis, A. D., Greischar, L. L., & Davidson, R. J. (2009). Mental training enhances attentional stability: neural and behavioral evidence. *Journal of Neuroscience*, *29*(42), 13418–13427. <https://doi.org/10.1523/JNEUROSCI.1614-09.2009>.
- Gaillard, A. W. (1977). The late CNV wave: preparation versus expectancy. *Psychophysiology*, *14*, 563–568.
- Gomez, C. M., Marco, J., & Grau, C. (2003). Preparatory visuo-motor cortical network of the contingent negative variation estimated by current density. *Neuroimage*, *20*(1), 216–224.
- MacCoon, D. G., MacLean, K. A., Davidson, R. J., Saron, C. D., & Lutz, A. (2014). No sustained attention differences in a longitudinal randomized trial comparing mindfulness based stress reduction versus active control. *PLoS One*, *9*(6). <https://doi.org/10.1371/journal.pone.0097551>.
- MacLean, K. A., Ferrer, E., Aichele, S. R., Bridwell, D. A., Zanesco, A. P., Jacobs, T. L., et al. (2010). Intensive meditation training improves perceptual discrimination and sustained attention. *Psychological Science: A Journal of the American Psychological Society / APS*, *21*(6), 829–839. <https://doi.org/10.1177/0956797610371339>.
- Malinowski, P. (2013). Neural mechanisms of attentional control in mindfulness meditation. *Frontiers in Neuroscience*, *7*, 1–11. <https://doi.org/10.3389/fnins.2013.00008>.
- Marcus, D. K., O'Connell, D., Norris, A. L., & Sawaqdeh, A. (2014). Is the Dodo bird endangered in the 21st century? A meta-analysis of treatment comparison studies. *Clinical Psychology Review*, *34*(7), 519–530.
- Mento, G. (2013). The passive CNV: carving out the contribution of task-related processes to expectancy. *Frontiers in Human Neuroscience*, *7*, 827.
- Moore, A., Gruber, T., Derose, J., & Malinowski, P. (2012). Regular, brief mindfulness meditation practice improves electrophysiological markers of attentional control. *Frontiers in Human Neuroscience*, *6*, 1–15. <https://doi.org/10.3389/fnhum.2012.00018>.
- Morales, J., Yudes, C., Gómez-Ariza, C. J., & Bajo, M. T. (2015). Bilingualism modulates dual mechanisms of cognitive control: evidence from ERPs. *Neuropsychologia*, *66*, 157–169. <https://doi.org/10.1016/j.neuropsychologia.2014.11.014>.
- Ng, K. K., Tobin, S., & Penney, T. B. (2011). Temporal accumulation and decision processes in the duration bisection task revealed by contingent negative variation. *Frontiers in Integrative Neuroscience*, *5*, 1–10. <https://doi.org/10.3389/fnint.2011.00077>.
- Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia*, *9*(1), 97–113. [https://doi.org/10.1016/0028-3932\(71\)90067-4](https://doi.org/10.1016/0028-3932(71)90067-4).
- Polich, J. (2007). Updating P300: an integrative theory of P3a and P3b. *Clinical Neurophysiology*, *118*(10), 2128–2148. <https://doi.org/10.1016/j.clinph.2007.04.019>.
- Richmond, L. L., Redick, T. S., & Braver, T. S. (2015). Remembering to prepare: the benefits (and costs) of high working memory capacity. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *41*(6), 1764–1777.
- Raffone, A., Marzetti, L., Del Gratta, C., Perrucci, M. G., Romani, G. L., & Pizzella, V. (2019). Toward a brain theory of meditation. *Progress in Brain Research*, *244*, 207–232. <https://doi.org/10.1016/bs.pbr.2018.10.028>.
- Santarnecchi, E., D'Arista, S., Egiziano, E., Gardi, C., Petrosino, R., Vatti, G., et al. (2014). Interaction between neuroanatomical and psychological changes after mindfulness-based training. *PLoS One*, *9*(10), 1–9. <https://doi.org/10.1371/journal.pone.0108359>.
- Schneider, W., Eschman, A., & Zuccolotto, A. (2002). *E-Prime 2.0 software*. Pittsburgh: Psychology SoftwareTools Inc..
- Segal, Z. V., Williams, J. M. G., & Teasdale, J. D. (2002). *Mindfulness-based cognitive therapy for depression*. Guilford.
- Slagter, H. A., Lutz, A., Greischar, L. L., Francis, A. D., Nieuwenhuis, S., Davis, J. M., & Davidson, R. J. (2007). Mental training affects distribution of limited brain resources. *PLoS Biology*, *5*(6), 1228–1235. <https://doi.org/10.1371/journal.pbio.0050138>.
- Stanislaw, H., & Todorov, N. (1999). Calculating of signal detection theory measures. *Behavior Research Methods, Instruments, & Computers*, *31*(1), 137–149.
- Tang, Y., Hölzel, B. K., Posner, M. I., & Ogfkvckqp, C. O. (2015). The neuroscience of mindfulness meditation. *Nature Reviews Neuroscience*, *16*, 213–225. <https://doi.org/10.1038/nrn3916>.
- Tarantino, V., Ehlis, A. C., Baehne, C., Boreatti-Huemmer, A., Jacob, C., Bisiacchi, P., & Fallgatter, A. J. (2010). The time course of temporal discrimination: an ERP study. *Clinical Neurophysiology*, *121*(1), 43–52.
- Tarantino, V., Mazzonetto, I., & Vallesi, A. (2016). Electrophysiological correlates of the cognitive control processes underpinning mixing and switching costs. *Brain Research*, *1646*, 160–173. <https://doi.org/10.1016/j.brainres.2016.05.048>.
- Tomasino, B., Campanella, F., & Fabbro, F. (2016). Medial orbital gyrus modulation during spatial perspective changes: pre- vs. post-8weeks mindfulness meditation. *Consciousness and Cognition*, *40*, 147–158. <https://doi.org/10.1016/j.concog.2016.01.006>.
- Travis, F., Tecce, J., Arenander, A., & Wallace, R. K. (2002). Patterns of EEG coherence, power, and contingent negative variation characterize the integration of transcendental and waking states. *Biological Psychology*, *61*(3), 293–319. [https://doi.org/10.1016/S0301-0511\(02\)00048-0](https://doi.org/10.1016/S0301-0511(02)00048-0).
- Travis, F., Tecce, J. J., & Guttman, J. (2000). Cortical plasticity, contingent negative variation, and transcendent experiences during practice of the transcendental meditation technique. *Biological Psychology*, *55*(1), 41–55. [https://doi.org/10.1016/S0301-0511\(00\)00063-6](https://doi.org/10.1016/S0301-0511(00)00063-6).
- Veneziani, C. A., & Voci, A. (2015). The Italian adaptation of the cognitive and affective mindfulness scale-revised. *TPM - Testing, Psychometrics, Methodology in Applied Psychology*, *22*(1), 43–52. <https://doi.org/10.4473/TPM22.1.4>.
- Veneziani, C. A., & Voci, A. (2014). The Italian adaptation of the Mindful Awareness Attention Scale and its relation with individual differences and quality of life indexes. *Mindfulness*, *6*(2), 373–381.
- Voss, M. W., Heo, S., Prakash, R. S., Erickson, K. I., Alves, H., Chaddock, L., et al. (2013). The influence of aerobic fitness on cerebral white matter integrity and cognitive function in older adults: results of a one-year exercise intervention. *Human Brain Mapping*, *34*(11), 2972–2985.
- Walter, W. G. (1964). Contingent negative variation: an electric sign of sensori-motor association and expectancy in the human brain. *Nature*, *230*, 380–384.
- Widmann, A., Schröger, E., & Maess, B. (2015). Digital filter design for electrophysiological data—a practical approach. *Journal of Neuroscience Methods*, *250*, 34–46. <https://doi.org/10.1016/j.jneumeth.2014.08.002>.