

E. Di Gristina, D. Varshanidze & G. Domina

Seed germination data of three *Hieracium* and one *Pilosella* (*Asteraceae*) from Sicily and southern Italy

Abstract

Di Gristina, E., Varshanidze, D. & Domina, G.: Seed germination data of three *Hieracium* and one *Pilosella* (*Asteraceae*) from Sicily and southern Italy [In Magrini, S. & Salmeri, C. (eds), *Mediterranean plant germination reports* – 2]. *Fl. Medit.* 30: 431-437. 2020. <http://dx.doi.org/10.7320/FIMedit30.431>

This work reports the first seed germination data of three *Hieracium* and one *Pilosella* (*Asteraceae*) taxa from Sicily and southern Italy. The studied taxa are: *Hieracium hypchoeroides* subsp. *cilentanum*, endemic to Campania, *H. schmidtii* subsp. *nebrodense*, endemic to Sicily, *H. terraccianoii*, endemic to Calabria and *Pilosella hoppeana* subsp. *sicula*, endemic to Sicily. The obtained results showed that the investigated taxa produce non-dormant seeds, with a high percentage of germination at all tested thermoperiods (15°C to 25°C and the alternating temperature of 25/15°C). Germination tests carried out in gibberellic acid (GA₃) provided similar germination percentages. The data presented in this paper provides useful information for the conservation of these endemic taxa.

Key words: apomixis, conservation, endemics, sexuality, taxonomy.

Introduction

Hieracium L. s. str. and *Pilosella* Vaill. (*Asteraceae*) are two of the largest angiosperm genera notorious for their taxonomic complexity caused by widespread interspecific hybridization, polyploidy, apomixis and sexuality (see, e.g., Krahulcová & al. 2000; Mráz & al. 2019). Many *Hieracium* and *Pilosella* taxa have a very restricted distribution and are punctual endemics. In southern Europe there are only local populations and most of them seem to be relict (Mráz & al. 2019). Therefore, their biological, phytogeographical and taxonomical relevance requires specific actions of *ex situ* and *in situ* protection. To this regard, the knowledge of the seed germination behaviour is an important contribution to the conservation of these taxa.

The taxonomy and distribution of *Hieracium* and *Pilosella* in southern Italy are currently under review. Recent studies resulted in the description of new endemic taxa with very local distributions (Brullo & al. 2001; Raimondo & Di Gristina 2004, 2007; Caldarella & al. 2014; Gottschlich & al. 2013, 2015; Di Gristina & al. 2013a, 2014, 2015a, 2015b, 2016a, 2016b, 2018, 2019). Also, little-known taxa were rediscovered (Di Gristina & al. 2016c; Gottschlich & al. 2017b) or reported as new to the Italian flora (Gottschlich & al. 2017a).

In this study, we present the seed germination data of four endemic hawkweed from Sicily and southern Italy: *Hieracium hypochoeroides* subsp. *cilentanum* Di Grist., Gottschl. & Raimondo, *H. schmidtii* subsp. *nebrodense* (Tineo ex Lojac.) Di Grist., Gottschl. & Raimondo, *H. terraccianoii* Di Grist., Gottschl. & Raimondo, and *Pilosella hoppeana* subsp. *sicula* Di Grist., Gottschl. & Raimondo

42. *Hieracium hypochoeroides* subsp. *cilentanum* Di Grist., Gottschl. & Raimondo (*Asteraceae*)

Accession data

It: Campania. Sanza (Salerno), Mt Cervati (Parco Nazionale del Cilento e Vallo di Diano) (WGS84: 40.290336°N, 15.477447°E), carbonate rocks, 1.860 m a.s.l., 09 Jul 2015, *E. Di Gristina* (PAL 102652).

Germination data

Pre-treatments: No treatment. Only manual removal of pappus.

Germination medium: Petri dishes with 2 sheets of sterilized Whatman 40 filter papers, imbibed in sterilized distilled water or in 10^{-3} M gibberellic acid (GA_3) water solution⁽¹⁾.

Sample size: 80 seeds for each test (20×4 replicates).

Germination	Thermoperiod	Photoperiod [light/dark]	T ₁ [d]	T ₅₀ [d]	T _{max} [d]	MTG [d]
94%	constant 20°C	0/24h	6.0	12.4	21.0	12.8
95%⁽¹⁾	constant 20°C	0/24h	5.0	11.8	22.0	12.8

Observations

Hieracium hypochoeroides subsp. *cilentanum* is a chasmophytic hawkweed endemic to Mt Cervati (Campania, S Italy) (Di Gristina & al. 2016a). The collective species *H. hypochoeroides* s. l. is a young aggregate of many apomictic microtaxa which have evolved probably during the post-glacial period (Di Gristina & al. 2015a). Many of the taxa so far described have a very restricted distribution and are punctual endemics (Di Gristina & al. 2013a, 2015b). In southern Europe there are only local populations, and most of them seem to be relict (Di Gristina & al. 2015b). *H. hypochoeroides* subsp. *cilentanum* matures and disperses seeds in July (Di Gristina & al. 2016a).

Germination tests were carried out 3 months after seed harvesting, using the constant temperatures of 15°C, 20°C, 25°C, and the alternating temperature of 25/15°C (16h/8h), in continuous darkness. The obtained results showed that *H. hypochoeroides* subsp. *cilentanum* produces non-dormant seeds, with a high percentage of germination at all tested thermoperiods. This can be due to a strategy aimed at allowing populations living in unfavourable ecological conditions, such as chasmophytic environments, to increase the probability of dispersal and affirmation of the progeny. The optimal ger-

mination temperature was 20°C (94%). Germination tests carried out in 10⁻³ M gibberellic acid (GA₃) provided similar germination percentages. No germination data are present in literature for this taxon nor for the *H. hypochoeroides* aggregate.

43. *Hieracium schmidtii* subsp. *nebrodense* (Tineo ex Lojac.) Di Grist., Gottschl. & Raimondo

Accession data

Si: Polizzi Generosa (Palermo), Monte Cavallo (Monti Madonie) (WGS84: 37.831973°N, 14.037852 E), quartzarenitic rocks, 1.400 m a.s.l., 28 Jun 2015, E. Di Gristina (PAL 102631).

Germination data

Pre-treatments: No treatment. Only manual removal of pappus.

Germination medium: Petri dishes with 2 sheets of sterilized Whatman 40 filter papers, imbibed in sterilized distilled water or in 10⁻³ M gibberellic acid (GA₃) water solution⁽¹⁾.

Sample size: 80 seeds for each test (20 × 4 replicates).

Germination	Thermoperiod	Photoperiod [light/dark]	T ₁ [d]	T ₅₀ [d]	T _{max} [d]	MTG [d]
90%	constant 20°C	0/24h	6.0	11.5	21.0	13.1
95% ⁽¹⁾	constant 20°C	0/24h	5.0	12.5	20.0	11.7

Observations

Hieracium schmidtii subsp. *nebrodense* is a chasmophytic hawkweed endemic to the Madonie Mountains (Sicily, Italy) (Di Gristina & al. 2016c). This taxon has been recently rediscovered after almost two centuries since its first and single collection made in 1830 and reclassified as a subspecies of the collective species *H. schmidtii* (Di Gristina & al. 2016c). This aggregate in Sicily is also represented by another taxon endemic to the Madonie Mountains, *H. schmidtii* subsp. *madoniense* (Raimondo & Di Grist.) Greuter (syn. *H. madoniense*, see Raimondo & Di Gristina 2007). *H. schmidtii* subsp. *nebrodense* matures and disperses seeds in June and July (Di Gristina & al. 2016c).

Germination tests were carried out 3 months after seed harvesting, using the constant temperatures of 15°C, 20°C, 25°C and the alternating temperature of 25/15°C (16h/8h), in continuous dark. In water, high percentage of germination was obtained at all tested thermoperiods, the highest (90%) and with high germination speed (T₅₀: 11.5, MTG: 13.1) at the constant temperature of 20°C. Germination tests carried out in 10⁻³ M gibberellic acid (GA₃) provided a bit higher germination percentage (95%). Here we report the first germination data for this taxon and for the *H. schmidtii* aggregate.

44. *Hieracium terracciano* Di Grist., Gottschl. & Raimondo (*Asteraceae*)**Accession data**

It: Calabria, Morano Calabro (CS), Scala di Gaudolino (Parco Nazionale del Pollino) (WGS84: 39.900303° N, 16.169194°E), carbonate rocky slopes, 1.350 m a.s.l., 12 Jul 2014, *E. Di Gristina* (PAL 101239).

Germination data

Pre-treatments: No treatment. Only manual removal of pappus.

Germination medium: Petri dishes with 2 sheets of sterilized Whatman 40 filter papers, imbibed in sterilized distilled water or in 10^{-3} M gibberellic acid (GA_3) water solution⁽¹⁾.

Sample size: 80 seeds for each test (20×4 replicates).

Germination	Thermoperiod	Photoperiod [light/dark]	T ₁ [d]	T ₅₀ [d]	T _{max} [d]	MTG [d]
90%	constant 15°C	0/24h	7.0	14.9	19.0	15.0
90% ⁽¹⁾	constant 15°C	0/24h	6.0	13.4	19.0	14.0

Observations

Hieracium terracciano is a pseudorosulate hemicryptophytic hawkweed endemic to the Pollino National Park (S Italy) (Di Gristina & al. 2014). This species belongs to a complex of similar morphotypes which have been grouped together in *H. sect. Grovesiana* (Gottschlich & al. 2013). *H. terracciano* matures and disperses the seeds in June and July (Di Gristina & al. 2014).

Germination tests were carried out 3 months after seed harvesting, using the constant temperatures of 15°C, 20°C, 25°C and the alternating temperature of 25/15°C (16h/8h), in continuous dark. The seeds of *H. terracciano* showed no dormancy and the highest germination (90%) at the temperature of 15°C. High germination was also obtained at 20°C reaching a value of 84%, while the alternating temperature of 25/15°C only reached a maximum of 65%. These results suggest that the seeds of *H. terracciano* prefer constant, cool or medium temperatures, for germination. Tests carried out in 10^{-3} M gibberellic acid (GA_3) provided similar germination percentages. Our germination tests are the first for this species and for the *H. sect. Grovesiana*.

45. *Pilosella hoppeana* subsp. *sicula* Di Grist., Gottschl. & Raimondo (*Asteraceae*)**Accession data**

Si: Polizzi Generosa (Palermo), Monte Scalone (Monti Madonie) (WGS84: 37.841030°N 14.018628°E), quartzarenitic rocky slopes, 1.510 m a.s.l., 15 Jun 2011, *E. Di Gristina* (PAL 101238).

Si: Nicosia (Enna), Monte Sambughetti, (Monti Nebrodi) (WGS84: 37.820000°N 14.385000°E), quartzarenitic rocky slopes, 1.420 m a.s.l., 20 Jun 2011, *E. Di Gristina* (PAL 100473).

Germination data

Pre-treatments: No treatment. Only manual removal of pappus.

Germination medium: Petri dishes with 2 sheets of sterilized Whatman 40 filter papers, imbibed in sterilized distilled water or in 10⁻³ M gibberellic acid (GA₃) water solution⁽¹⁾.

Sample size: 80 seeds for each test (20 × 4 replicates).

Germination	Thermoperiod	Photoperiod [light/dark]	T ₁ [d]	T ₅₀ [d]	T _{max} [d]	MTG [d]	Herbarium Id no.
85%	constant 20°C	0/24h	5.0	9.2	18.0	10.0	PAL101238
87% ⁽¹⁾	constant 20°C	0/24h	5.0	9.1	17.0	9.9	PAL101238

Observations

Pilosella hoppeana subsp. *sicula* is a rosulate hemicryptophytic hawkweed endemic to the Madonie and Nebrodi Mountains (Sicily, Italy) (Di Gristina & al. 2016b). The *P. hoppeana* (Schult.) F.W. Schultz & Sch. Bip. aggregate includes widespread taxa and also very localized ones (Di Gristina & al. 2013b). In Sicily the group is represented by the widely spread *P. hoppeana* subsp. *macrantha* (Ten.) S. Bräut. & Greuter and *P. hoppeana* subsp. *sicula*, recently described on the basis of an integrated morphological, karyological and isoenzymatic study (Di Gristina & al. 2013b). The two populations of *P. hoppeana* subsp. *sicula* from the Madonie and Nebrodi Mountains were investigated. This taxon matures and disperses seeds in June and July (Di Gristina & al. 2016b).

Germination tests were carried out 3 months after seed harvesting, using the constant temperatures of 15°C, 20°C, 25°C and the alternating temperature of 25/15°C (16h/8h), in continuous dark. The examined populations of *P. hoppeana* subsp. *sicula* showed no seeds dormancy and 20°C as optimal germination temperature. However, the germination rates were different between the two populations: the Madonie one showed the highest percentage (85%) with a high germination speed (T₅₀: 9.25 MTG: 10), while the seed germination value of the Nebrodi population only reached 75%, with lower speed (T₅₀: 14.5 MTG: 15.4). Tests carried out in 10⁻³ M gibberellic acid (GA₃) provided similar or slightly higher germination percentages. These are the first germination data for this taxon and for the *P. hoppeana* aggregate.

References

Brullo, S., Scelsi, F. & Spampinato, G. 2001: La Vegetazione dell'Aspromonte. Studio fitosociologico. – Reggio Calabria.

Caldarella, O., Gianguzzi, L. & Gottschlich, G. (2014) *Hieracium busambarense*, a new species of the sect. *Grovesiana* (Asteraceae) from Sicily (Italy). – Pl. Biosyst. **148**: 439-443. <https://doi.org/10.1080/11263504.2013.778352>

- Di Gristina, E., Gottschlich, G. & Raimondo, F. M. 2014: *Hieracium terraccianoi* (Asteraceae), a new species endemic to the Pollino National Park (Southern Italy). – *Phytotaxa* **188**(1): 55-60. <https://doi.org/10.11646/phytotaxa.188.1.8>
- , — & — 2015a: *Hieracium hypochoeroides* subsp. *peracutisquamum* (Asteraceae), a new taxon from Basilicata, southern Italy. – *Ann. Bot. Fennici* **52**: 376-380. <http://dx.doi.org/10.5735/085.052.0519>
- , — & — 2015b: Taxonomic remarks on *Hieracium sartorianum* var. *lucanicum* (Asteraceae), a little known taxon of Cilento (Campania, southern Italy). – *Nordic J. Bot.* **33**: 465-468. <http://dx.doi.org/10.1111/njb.00755>
- , — & — 2016a: *Hieracium hypochoeroides* subsp. *cilentanum* (Asteraceae), a new taxon from S Italy. – *Phytotaxa* **246**(3): 192-197. <https://doi.org/10.11646/phytotaxa.246.3.3>
- , — & — 2016b: *Pilosella hoppeana* subsp. *sicula* Di Grist., Gottschl. & Raimondo (Asteraceae), a new endemic subspecies from Sicily (Italy). – *Candollea* **71**(1): 7-12. <https://doi.org/10.15553/c2016v711a2>
- , — & — 2016c: Rediscovery of *Hieracium nebrodense* (Asteraceae), a little-known endemic of Sicily (Italy). – *Phytotaxa* **265**(1): 59-66. <https://doi.org/10.11646/phytotaxa.265.1.5>
- , — & Scafidi, F. 2018: *Hieracium jurassicum* subsp. *serrapretense* (Asteraceae), a new hawkweed taxon from the Pollino National Park (southern Italy). – *Phytotaxa* **340**(2): 186-190. <https://doi.org/10.11646/phytotaxa.340.2.8>
- , Domina, G., Gottschlich, G., Maturo, F. & Scafidi, F. 2019: *Hieracium racemosum* subsp. *lucanum* (Asteraceae), a new hawkweed from southern Italy. – *Phytotaxa* **425**(5): 297-300. <https://doi.org/10.11646/phytotaxa.425.5.5>
- , —, P. Mazzola & Geraci, A. 2013b: Morphological and genetic diversity within *Pilosella hoppeana* aggr. (Asteraceae) in Italy and taxonomic implications. – *Pl. Biosyst.* **147**(3): 788-799. <https://doi.org/10.15553/c2016v711a2>
- , Gottschlich, G., Galesi, R., Raimondo, F. M. & Cristaudo, A. 2013a: *Hieracium hypochoeroides* subsp. *montis-scuderii* (Asteraceae), a new endemic subspecies from Sicily (Italy). – *Fl. Medit.* **23**: 49-55. <http://dx.doi.org/10.7320/FIMedit23.049>
- Gottschlich, G., Domina, G. & Di Gristina, E. 2017a: *Hieracium umbrosum* subsp. *abietinum* (Asteraceae), a further example of amphi-Adriatic disjunction. – *Pl. Biosyst.* **151**: 792-794. <https://doi.org/10.1080/11263504.2017.1341439>
- , Raimondo, F. M. & Di Gristina, E. 2013: *Hieracium pallidum* subsp. *aetnense* (Asteraceae), a new subspecies from Sicily (Italy), with notes on the taxonomy of *H. pallidum* Biv. – *Pl. Biosyst.* **147**(3): 826-831. <https://doi.org/10.1080/11263504.2013.829882>
- , Scafidi, F. & Di Gristina, E. 2017b: *Hieracium pollinense* (Asteraceae), an endemic species to the Pollino National Park (Southern Italy) rediscovered. – *Acta Bot. Croat.* **76**(1): 95-97. <https://doi.org/10.1515/botcro-2016-0047>
- , Raimondo, F.M., Greuter, W. & Di Gristina, E. 2015: *Hieracium barrelieri*, a new hawkweed species from S Italy, with notes on Tenore's *Hieracium murorum* var. *barrelieri* (Asteraceae). – *Phytotaxa* **208**(1): 70-74. <http://dx.doi.org/10.11646/phytotaxa.208.1.7>
- Krahulcová, A., Krahulec, F. & Chapman, H. M. 2000: Variation in *Hieracium* subgen. *Pilosella* (Asteraceae): What do we know about its sources? – *Folia Geobot.* **35**: 319-338.
- Mráz, P., Filipas, L., Bărbos, M. I., Kadlecová, L., Belyayev, A. & Fehrer, J. 2019: An unexpected new diploid *Hieracium* from Europe: Integrative taxonomic approach with a phylogeny of diploid *Hieracium* taxa. – *Taxon* **68**(6): 1258-1277. <https://doi.org/10.1002/tax.12149>
- Raimondo, F. M. & Di Gristina, E. 2004: *Hieracium pignattianum* (Asteraceae), a new species from the Madonie Mountains (N-Sicily). – *Bocconea* **17**: 313-324.

— & — 2007: *Hieracium madoniense* (Asteraceae), a new species from Sicily. – Pl. Biosyst. **141(1)**: 86-92. <https://doi.org/10.1080/11263500601153982>

Addresses of the authors:

Emilio Di Gristina^{1*}, Dali Varshanidze² & Gianniantonio Domina¹,

¹ Department of Agricultural, Food and Forest Sciences (SAAF), University of Palermo, Viale delle Scienze, bld. 4, 90128 Palermo, Italy. E-mail: emilio.digristina@unipa.it; gianniantonio.domina@unipa.it;

² Batumi Botanical Garden, Green Cape, Batumi, 6411, Georgia. E-mail: dvarshanidze@bbg.ge

*Corresponding author