

PALAEARCTIC GRASSLANDS

Journal of the Eurasian Dry Grassland Group



Research Article

DOI: 10.21570/EDGG.PG.62.28-33

(Re)discovery of four annual vascular plant species in dry grasslands in the Southwestern Alps (NW Italy)

Riccardo Guarino^{1*}, Denys Vynokurov^{2,3,4}, Jürgen Dengler⁵, Asun Berastegi⁶, Idoia Biurrun⁷, Itziar Garcia-Mijangos⁷, Monika Janišová⁸, Michele Lonati⁹, Robin Nikolei¹⁰, Andrea Mainetti¹¹, Olena Miskova⁴, Sara Sánchez-Carmona⁷, Oksana Tyshchenko¹² & Ginevra Nota⁹

¹Department of Biological, Chemical and Pharmaceutical Sciences and Technologies, University of Palermo, Palermo, Italy; riccardo.guarino@unipa.it

²Institute of Biology/Geobotany and Botanical Garden, Martin Luther University Halle-Wittenberg, Halle, Germany; denys.vynokurov@gmail.com

³German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany

⁴M.G. Kholodny Institute of Botany, National Academy of Sciences of Ukraine, Kyiv, Ukraine

⁵Vegetation Ecology Research Group, Institute of Natural Resource Sciences, Zurich University of Applied Sciences (ZHAW), Wädenswil, Switzerland; dr.juergen.dengler@gmail.com

⁶Navarra Environmental Management (GAN-NIK), Pamplona-Iruña, Spain; aberastg@gan-nik.es

⁷Department of Plant Biology and Ecology, Faculty of Science and Technology, University of the Basque Country UPV/EHU, Bilbao, Spain; idoia.biurrun@ehu.eus (I. Biurrun), itziar.garcia@ehu.eus (I. García-Mijangos), sara.sanchez@ehu.eus (Sara Sánchez-Carmona)

⁸Institute of Botany, Plant Science and Biodiversity Centre, Slovak Academy of Sciences, Bratislava, Slovakia; monika.janisova@gmail.com

⁹Department of Agriculture, Forest and Food Sciences, University of Torino, Grugliasco (TO), Italy; michele.lonati@unito.it (M. Lonati), ginevra.nota@unito.it (G. Nota)

¹⁰Ober dem Pfarracker 1, 63826 Geiselbach, Germany; robin@nikolei.org

¹¹Gran Paradiso National Park, Cogne (AO), Italy; andrea.mainetti@pngp.it

¹²Plant Biology Department, Taras Shevchenko National University of Kyiv, Kyiv, Ukraine; oksana_tyshchenko@knu.ua

*) Corresponding author

Palaeoartctic Grasslands 62 (2024): 28-33

Abstract: The Eurasian Dry Grassland Group (EDGG) has developed a rigorous sampling methodology aimed at comprehensively documenting plant diversity across non-forest habitats in the Palaeoartctic realm. During the 18th EDGG Field Workshop devoted to the dry grasslands in the Southwestern Alps, three annual vascular plant species not observed in the Aosta Valley for over one century were re-discovered: *Bupleurum baldense*, *Hypochaeris glabra*, and *Trifolium dubium*. Moreover, the annual grass species *Brachypodium distachyon* was recorded for the first time from the Piedmont Region in the Susa valley. We provide here details on the identification and distribution of these four species, along with some environmental features of the discovery sites. Our findings do not only enrich the regional flora but also underscore the importance of historical records and accurate floristic exploration for biodiversity assessment.

Keywords: Aosta Valley; *Brachypodium distachyon*; *Bupleurum baldense*; *Hypochaeris glabra*; inneralpine dry valley; Italy; new record; Piedmont; regional flora; *Trifolium dubium*.

Nomenclature: Bartolucci et al. (2024) for vascular plants; Nimis et al. (2018) for lichens; Mucina et al. (2016) for syntaxa.

Abbreviations: EDGG = Eurasian Dry Grassland Group.

Submitted: 15 October 2024; Accepted: 31 October 2024

Scientific Editor: Alireza Naqinezhad

Introduction

The [Eurasian Dry Grassland Group](#) (EDGG) has developed a thorough sampling methodology to ensure high-quality data collection on plant diversity from non-forest habitats across the Palaeoarctic realm. This approach includes precisely delimited plots, standardized plot sizes, and meticulous sampling techniques that pay equal attention to vascular plants, terricolous bryophytes, and lichens, alongside measurements of a range of environmental variables (Dengler et al. 2016).

Such accurate exploration often leads to significant discoveries, including new regional or national records (Dengler & Boch 2007; Moysiyanenko et al. 2022; Boch et al. 2024; Dengler et al. 2024), and even species new to science (Khodosovtsev et al. 2024). This highlights the critical importance of comprehensive floristic exploration for biodiversity assessment, which is essential for conservation efforts and advancing our understanding of ecological communities.

During sampling of semi-natural grasslands with the EDGG methodology in the Southwestern Alps (Abdulahak et al. 2024), an annual grass not recorded for the Piedmont region before was found in the Susa valley. Moreover, three annual species not observed in Aosta Valley for over a century and thus requiring confirmation for the regional flora, were rediscovered. We thus provide here a detailed overview of the species identification, their overall distribution and some environmental features of the discovery sites.

Methods

The 18th EDGG Field Workshop took place in the inner-alpine dry valleys of the Southwestern Alps, from June 1st to 11th 2024 (Abdulahak et al. 2024). This research expedition focused on three valleys with a continental climate: Aosta and Susa in Italy, and Durance in France. In total, 43 nested plots (nested-plot series of 0.0001–100 m², see Dengler et al. 2016) and 155 single plots, 10 m² each, were sampled, resulting in 241 single 10 m² plots if the 10 m² subplots from the nested-plot series are counted. Plots were distributed across semi-natural grassland types of the three mentioned valleys, between 296 and 1880 m a.s.l., with the aim of portraying most of the variability of the dry grassland types occurring there.

Three species targeted in this floristic report were collected in the lower south-eastern reaches of Aosta Valley (NW Italy), where sub-Mediterranean species are more frequent. The fourth species was collected in the furthest downstream part of the Susa Valley (Piedmont region), on the lower slopes of Mount Musinè. The climate in the studied area is typical of the inner valleys of the Alps, with annual precipitation ranging from 460 to 700 mm (Mercalli 2003).

The collected specimens were examined using a stereomicroscope, identified with the Flora of Italy (Pignatti et al. 2017–2019), pressed, dried, and deposited at the Herbarium AO or at the Herbarium TO (all herbarium codes men-

tioned here refer to Thiers 2022+). Information on regional historical records of the re-discovered species proceeds from the Vascular Flora of Valle d'Aosta (Bovio 2018+).

Results

Brachypodium distachyon (L.) P.Beauv. was sampled in the plot SDA01x on 2 June 2024. This plot was located in the municipality of Almese, on the south-facing slope of Mount Musinè (45.10392° N, 7.44489° E). It grew in a xeric grassland at 420 m a.s.l. (Fig. 1). Based on a preliminary phytosociological classification, the stand belonged to the *Diplachnion serotinae*. The herb layer covered 67% and the cryptogam layer 6%. The total plant species richness in 100 m² was 26 (23 vascular plants, 2 bryophytes and 1 lichen). The most dominant species were *Heteropogon contortus* (36.5%), *Chrysopogon gryllus* (19%), *Thymus oenipontanus* (5.5%), *Fumana procumbens* (3.8%), *Cleistogenes serotina* (3.5%) in the herb layer, and *Weissia* sp. (6.5%) in the cryptogam layer.

Brachypodium distachyon, a new model plant for functional genomics of grasses and monocots (Garvin et al. 2008; Catalán et al. 2012), has a native distribution range from the Mediterranean basin to Central Asia. The species has also been introduced to Central Europe, North and South America, Australia and South Africa. It grows across a broad range of elevations, from the sea level to about 2000 m a.s.l. (Garvin et al. 2008; Catalán et al. 2012). In Europe, the species gives the name to the alliance *Trachynion distachyae* (*Stipo-Trachynietea distachyae*), distributed in the Western Mediterranean. Before the present discovery, the species was reported to occur in most of the Italian territory, except for Piedmont and Aosta Valley. Today, *Brachypodium distachyon* has been proposed as a complex of three distinct species having cytotypes of 2n = 10, 20, and 30 chromosomes (Catalán et al. 2012) and characterized by different environmental niches (López-Alvarez et al. 2015). Ecological niche models suggest that our specimen probably refers to *Brachypodium distachyon* s.str. growing in colder, wetter and higher-elevated conditions than the other taxa of the complex.

Bupleurum baldense Turra was sampled in the plot SDAR205 on 8 June 2024. This plot was located in the municipality of Montjovet (Aosta Valley), next to the south-facing walls of the Chenal Castle (45.73007° N, 7.67184° E). It grew in a xeric grassland topping a sheepback rock at 692 m a.s.l. (Fig. 2). Based on a preliminary phytosociological classification, the stand belonged to the alliance *Stipo-Poion xerophilae*. The herb layer covered 94% and the cryptogam layer 7%, and the total plant species richness in 10 m² was 45 (40 vascular plants, 2 bryophytes and 3 lichens). The most dominant species were *Bromopsis erecta* (27%), *Trifolium arvense* (15%); *Aira caryophyllea*, *Festuca* cf. *laevigata* and *Sempervivum tectorum* (all 12%) in the herb layer and *Cladonia foliacea* (3%) and *C. rangiformis* (2%) in the cryptogam layer.

Bupleurum baldense (incl. *B. veronense* Turra) is endemic to

Southern and Western Europe, ranging from Southern Britain and France to the Eastern Iberian Peninsula, Balearic Islands, Sardinia, Corsica, Italian Peninsula, Maltese Archipelago, Sicily, and Western Balkan Peninsula (Snogerup & Snogerup 2001). In Italy, it has been recorded in all regions except for Trentino-Alto Adige and Molise, from the sea level to 1500 m a.s.l. (Bartolucci et al. 2024). However, only two historical records were available from Aosta Valley, both of which repeatedly reported in recent floras as putatively disappeared from the region. The first record is an herbarium specimen collected by Charles Ferina (sine die) from Montjovet and verified and cited by Lino Vaccari (1904–1911), who also collected the species in Montjovet on the hills of Plout, a village located at about 3 km south of the site of the present note. Our finding matches also very well the location and altitude of the old herbarium sheet collected by an unknown collector and stored in AO-S.SFV (sub *Bupleurum odontites* L., Montjovet, 600 m a.s.l.), which could perhaps be the otherwise lost Ferina's specimen cited by Vaccari (Bovio 2014; 2018+).

The specimens studied by us correspond to the var. *aristatum* (Bartl.) Briq., typically found in xeric grasslands on heterogeneous, limestone-rich substrates (also on recent alluvial depositions and screes), in continental valleys of the lower mountain belt of NW Alps, where it was indicated by Braun-Blanquet (1961) as a typical taxon of the *Stipo-Poion xerophilae* (*Festucetalia valesiacae*).

Hypochaeris glabra L. and *Trifolium dubium* Sibth. were sampled in the plot SDAR020 on 8 June 2024. This plot was located 60 m WNW of Chenal Castle, in the municipality of Montjovet (Aosta Valley, 45.73049° N, 7.67084° E). It grew in a mesoxeric grassland at 627 m a.s.l. (Fig. 3). Based on a preliminary phytosociological classification, the stand belonged to the *Brachypodietalia pinnati*. The herb layer covered 85% and the cryptogam layer 5%, and the total plant species richness in 10 m² was 43 (38 vascular plants, 4 bryophytes and 1 lichen). The most dominant species were *Carex humilis* (30%), *Bromopsis erecta* (13%), *Filipendula vulgaris* (9%), and *Peucedanum oreoselinum* (6%) in the herb layer and *Weissia* sp. (4%) in the cryptogam layer.

Hypochaeris glabra has a Mediterranean-temperate distribution, encompassing Europe, North Africa, and warm-temperate western Asia. It is also widely naturalised across Australia, New Zealand, South Africa, South America and North America (Stroh 2015). In Italy, it has been recorded in all regions except for Trentino-Alto Adige and Molise, from the sea level to 1500 m a.s.l.; however, the only record of this species from Aosta Valley relates to a historical specimen stored in TO, collected by Antonio Carestia at Albard from Donnas in 1871. This occurrence was mentioned by Vaccari (1904–1911) and reported as no longer found by Bovio (2014, 2018+). Our finding is located much further into the Aosta Valley than Carestia's find, about 20 km WNW from Albard di Donnas.

Trifolium dubium is an allotetraploid native to most of Europe and the Caucasus region, resulting from the hybridiza-

tion of *T. campestre* and *T. micranthum*. It has naturalised as an adventive pasture species and as a weed of wasteland and roadsides in North America and many countries of the southern hemisphere (Ansari et al. 2008). It occurs in all regions of the Italian Peninsula, between 200 and 1600 m a.s.l., but it has never been observed in Sicily and Sardinia. As for Aosta Valley, there was only historical bibliographic data, not supported by recent findings. Favre (1874) reported it below Saint-Rhémy, at 1620 m a.s.l. Vaccari (1904–1911) added the following personal observations to Favre's record: Cignai, Bard, Arnad, lower Valley of Champorcher, Valley of Cogne above Vieyes, and between Châtillon and Valtournenche. However, considering the many localities mentioned in Vaccari's Catalogue, the lack of specimens in his herbarium leads to suppose that his reports of *T. dubium* could be due to misidentified exsiccata, which were later corrected and attributed to other species. For this reason, and the lack of recent findings, Bovio (2014, 2018+) suggested that the occurrence of this species in Aosta Valley needed to be confirmed.

Discussion

The finding of *Brachypodium distachyon* in Susa Valley and the rediscovery of *Bupleurum baldense*, *Hypochaeris glabra*, and *Trifolium dubium* in Aosta Valley after more than one century since the last observation emphasizes the value of comprehensive and accurate floristic surveys in semi-natural grasslands. The application of the EDGG sampling methodology, which incorporates multi-taxon approaches and standardized plot sizes, along with particularly accurate sampling, has proven instrumental in detecting these elusive species.

Interestingly, all four species are annuals, which in general have higher dispersal ability. However, the sites in which we found them were neither recently disturbed nor close to typical entry paths of (regional) neophytes, such as cities and traffic networks. Rather than a recent (re-)introduction, this suggests that the species had strived in the regions for a longer period of time and just due to their small stature and low number of individuals had been overlooked by botanists.

Our study confirms that xeric and semi-natural grasslands in the inner-alpine dry valleys are highly valuable for preserving plant species with fragmented and isolated distributions, otherwise absent or rarely found in the Alpine Region (Dengler et al. 2020). The climate uniqueness of these valleys, hosting relict populations of many plant species that reached these isolated outposts during Pleistocene climatic fluctuations, makes the continental xerothermic islands of the inner Alps a priority target for the conservation of European biodiversity. Our findings also underscore the importance of botanical collections and historical records to reconstruct the history, dynamics, and current distribution of plant species (Buldrini et al. 2023), as well as the need for continued fieldwork to integrate historical botanical records with contemporary data.

Author contributions

Most authors were involved in the field sampling of the three plots. D.V. and J.D. are coordinating the EDGG Field Workshops, M.L., A.M. and G.N. co-organized the 18th Field Workshop; R.G. planned and drafted the article, G.N. drafted the section on *Brachypodium distachyon*, all other authors revised and commented on the text.

Acknowledgements

We thank Maurizio Bovio for revising the herbarium specimens and Alireza Naqinezhad as well as one anonymous reviewer for the very fast and careful peer review.

References

- Abdulkhak, S., Lonati, M., Mainetti, A., Marengo, G., Nota, G., Dengler, J. & Vynokurov, D. 2024. 18th EDGG Field Workshop – Dry grasslands of South-Western Alps – Aosta, Susa and Durance valleys, Italy and France, 1–11 June 2024, Second Call. *Palaeoartctic Grasslands* 60: 9–15.
- Ansari, H.A., Ellison, N.W. & Williams, W.M. 2008. Molecular and cytogenetic evidence for an allotetraploid origin of *Trifolium dubium* (Leguminosae). *Chromosoma* 117: 159–167.
- Bartolucci, F., Peruzzi, L., Galasso, G., Alessandrini, A., Ardenghi, N.M.G., Bacchetta, G., Banfi, E., Barberis, G., Bernardo, L., (...) & Conti, F. 2024. A second update to the checklist of the vascular flora native to Italy. *Plant Biosystems* 158: 219–296.
- Boch, S., Dembic, I., Keller, C. & Dengler, J. 2024. *Cladonia homosekikaica* Nuno new to Sweden: a cup lichen found in the grey dunes of Gotland. *Palaeoartctic Grasslands* 61: 14–17.
- Bovio, M. 2014. *Flora Vascolare della Valle d'Aosta*. Testolin editore, Sarre, AO, IT.
- Bovio, M. 2018+. *Flora Vascolare della Valle d'Aosta*. URL: <http://www.floravda.it/it/> [accessed 16 August 2024].
- Braun-Blanquet, J. 1961. Die inneralpine Trockenvegetation. Von der Provence bis zur Steiermark. *Geobotanica Selecta* 1: 1–273.
- Buldrini, F., Pezzi, G., Barbero, M., Alessandrini, A., Amadei, L., Andreatta, S., Ardengui, N.M.G., Armiraglio, S., Bagella, S., (...) & Lambertini, C. 2023. The invasion history of *Elodea canadensis* and *E. nuttallii* (Hydrocharitaceae) in Italy from herbarium accessions, field records and historical literature. *Biological Invasions* 25: 827–846.
- Catalán, P., Müller, J., Hasterok, R., Jenkins, G., Mur, L.A.J., Langdon, T., Betekhtin, A., Siwinska, D., Pimentel, M. & López-Alvarez, D. 2012. Evolution and taxonomic split of the model grass *Brachypodium distachyon*. *Annals of Botany* 109: 385–405.



Figure 1. “Biodiversity Plot” SDA01x in a stand of the *Diplachnion serotinae* on Mount Musinè, Susa valley, where *Brachypodium distachyon* was recorded for the first time in the Region of Piedmont. Photo: J. Dengler.

- Dengler, J. & Boch, S. 2007. Taxonomy, ecology, and distribution of six remarkable plant taxa (Spermatophyta, Bryophyta, and Lichenes) of the Estonian dry grassland flora. *Phyton Annales Rei Botanici* 47: 47–71.
- Dengler, J., Boch, S., Filibeck, G., Chiarucci, A., Dembicz, I., Guarino, R., Henneberg, B., Janišová, M., Marcenò, C., (...) & Biurrun, I. 2016. Assessing plant diversity and composition in grasslands across spatial scales: the standardised EDGG sampling methodology. *Bulletin of the Eurasian Dry Grassland Group* 32: 13–30.
- Dengler, J., Guarino, R., Moysiyenko, I., Vynokurov, D., Boch, S., Cykowska-Marzencka, B., Babbi, M., Catalano, C., Eggenberg, S., (...) & Dembicz, I. 2020. On the trails of Josias Braun-Blanquet II: First results from the 12th EDGG Field Workshop studying the dry grasslands of the inneralpine dry valleys of Switzerland. *Palaeartic Grasslands* 45: 59–88.
- Dengler, J., Barandun, M., Borovyk, D., Dembicz, I., Frey, D. & Vynokurov, D. 2024. The overlooked diversity of dry grasslands in Ticino (Switzerland) – with impressions from the 19th EDGG Field Workshop. *Palaeartic Grasslands* 62: 54–69.
- Favre, E. 1874. Supplément au Guide du botaniste sur le Grand Saint-Bernard de M. le Chanoine Tissière. *Bulletin des travaux de la Murithienne* 4: 41–78.
- Garvin, D.F., Gu, Y.-Q., Hasterok, R., Hazen, S.P., Jenkins, G., Mockler, T.C., Mur, L.A.J. & Vogel J.P. 2008. Development of genetic and genomic research resources for *Brachypodium distachyon*, a new model system for grass crop research. *Crop Science* 48: 69–84.
- Khodosovtsev, A., Darmostuk, V., Dembicz, I., Dengler, J., Moysiyenko, I. & Kuzemko, A. 2024. *Circinaria ucrainica* sp. nov., a new species from sand dunes of the Lower Dnipro valley (Ukraine). *The Lichenologist* 56: 159–167.
- López-Alvarez, D., Manzaneda, A.J., Rey, P.J., Giraldo, P., Benavente, E., Allainguillaume J., Mur, L., Caicedo, A.L., Hazen, S.P., (...) & Catalán, P. 2015. Environmental niche variation and evolutionary diversification of the *Brachypodium distachyon* grass complex species in their native circum-Mediterranean range. *American Journal of Botany* 102: 1073–1088.
- Mercalli, L. 2003. Atlante climatico della Valle d'Aosta. [Climatic atlas of the Aosta Valley]. Regione Autonoma Valle d'Aosta, Direzione Protezione civile, Ufficio Meteorologico, AO, IT.
- Moysiyenko, I., Umanets, O.Y., Dengler, J., Guarino, R., Dembicz, I., Kucher, O.O., Skobel, N.O. & Bednarska, I.O. 2022. *Torilis pseu-*



Figure 2. Vegetation plot SDAR205 in a stand probably belonging to the *Stipo-Poion xerophilae* at the Chenal castle, where *Bupleurum baldense* was rediscovered for the Aosta Valley. Photo: I. Biurrun.

- donodosa* Bianca (Apiaceae): new species for the flora of Ukraine. *Chornomorski Botanical Journal* 17: 331–338.
- Mucina, L., Bültmann, H., Dierßen, K., Theurillat, J.-P., Raus, T., Čarni, A., Šumberová, K., Willner, W., Dengler, J., (...) & Tichý, L. 2016. Vegetation of Europe: Hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities. *Applied Vegetation Science* 19 (Suppl. 1): 3–264.
- Nimis, P.L., Hafellner, J., Roux, C., Clerc, P., Mayrhofer, H., Martellos, S. & Bilovitz, P.O. 2018. The lichens of the Alps – an annotated checklist. *MycKeys* 31: 1–634.
- Pignatti, S., Guarino, R. & La Rosa, M. 2017–2019. *Flora d'Italia, volumes 1–4*. Edagricole, Edizioni Agricole di Newbusiness Media, Bologna, IT.
- Snogerup, S. & Snogerup, B. 2001. *Bupleurum* L. (Umbelliferae) in Europe—1. The annuals, B. sect. *Bupleurum* and sect. *Aristata*. *Willdenowia* 31(2): 205–308.
- Stroh, P.A. 2015. *Hypochaeris glabra* L., Smooth Cat's-ear. Species Account. Botanical Society of Britain and Ireland. URL: https://bsbi.org/wp-content/uploads/dlm_uploads/Hypochaeris_glabra_species_account.pdf [accessed 16 August 2024].
- Thiers, B. 2022+. *Index herbariorum: A global directory of public herbaria and associated staff*. New York Botanical Garden's Virtual Herbarium. URL: <http://sweetgum.nybg.org/ih/> [accessed 16 August 2024].
- Vaccari, L. 1904–1911. *Catalogue raisonné des plantes vasculaires de la Vallée d'Aoste. Volume I. Thalamiflores et Calyciflores*. Impr. Catholique, VIII, Aoste, IT.



Figure 3. Vegetation plot SDAR020 in a stand probably belonging to the mesoxeric order *Brachypodietalia pinnati* at Chenal castle, where *Hypochaeris glabra* and *Trifolium dubium* were rediscovered for Aosta Valley. Photo: M. Janišová.