

Magmatic Activity in Incipient Continental Break-Up as Revealed by Coupling Melt and Fluid Inclusions

G. BOUDOIRE^{1,2,*}, N. BOBROWSKI^{3,4}, P.-Y. BURGI⁵, S. CALABRESE^{2,6}, L. FRANCE^{7,8}, G. GIUFFRIDA³, F. GRASSA², K. KARUME⁹, J.-C. KAZADI MWEPU⁹, J. KUHN¹⁰, R. MORITZ¹¹, O. MUNGUIKO MUNYAMAHORO^{1,9}, A. L. RIZZO^{12,13} and D. TEDESCO^{14,15,16}

¹Laboratoire Magmas et Volcans, OPGC, CNRS, IRD, Université Clermont Auvergne, 6 Avenue Blaise Pascal, 63178 Aubière, France

²Istituto Nazionale di Geofisica e Vulcanologia, Sezione di Palermo, Via Ugo La Malfa 153, 90146 Palermo, Italy

³Istituto Nazionale di Geofisica e Vulcanologia, Sezione di Catania, Piazza Roma 2, 95125 Catania, Italy

⁴Institut für Umweltpophysik, Ruprecht Karls University, Im Neuenheimer Field 229, D-69120 Heidelberg, Germany

⁵IT Department, University of Geneva, 24 rue du Général-Dufour, 1211 Geneva, Switzerland

⁶DiSTeM, Università degli Studi di Palermo, Via Archirafi 22, 90123 Palermo, Italy

⁷CNRS, CRPG, Université de Lorraine, 15 rue Notre Dame des Pauvres, 54500 Vandoeuvre-lès-Nancy, France

⁸Institut Universitaire de France, 1 rue Descartes, 75231 Paris, France

⁹Observatoire Volcanologique de Goma, 142 Av. du Rond-Point, Q. Le Volcan, Goma, Democratic Republic of Congo

¹⁰Department of Atmospheric and Oceanic Sciences, University of California, 520 Portala Plaza, Los Angeles, CA 90095, USA

¹¹Department of Earth Science, University of Geneva, Rue des Maraichers 13, CH-1205, Geneva, Switzerland

¹²Dipartimento di Scienze dell'Ambiente e della Terra, Università degli Studi di Milano-Bicocca, Piazza della Scienza 4, 20126 Milan, Italy

¹³Istituto Nazionale di Geofisica e Vulcanologia, Sezione di Milano, Via Bassini 15, 20133 Milano, Italy

¹⁴DiSTABiF, Università degli Studi della Campania "Luigi Vanvitelli", Via Vivaldi 43, 81100 Caserta, Italy

¹⁵Osservatorio Vesuviano, Istituto Nazionale di Geofisica e Vulcanologia, Via Diocleziano 328, 80124 Napoli, Italy

¹⁶United Nations-MONUSCO, 27 Av. de la Révolution, Goma, Republic Democratic of Congo

*Corresponding author. Telephone: +33 (0)473346723; E-mail: guillaume.bouidoire@gmail.com

Deciphering deep magmatic processes driving the onset of continental break-up is fundamental to constrain our understanding of plate tectonics. The East African Rift System (EARS) represents the only opportunity to study a currently active system on Earth undergoing distinct stages of rift evolution. We present a coupled analysis of melt and fluid inclusions in the Virunga Volcanic Province (VVP) offering unprecedented insight into the dynamics of incipient rifting and its evolution. Our study highlights that melting of distinct metasomes in the deep lithosphere is a common feature of immature rifts. In the VVP, it leads to the emission of nephelinitic and basanitic melts at Nyiragongo and Nyamulagira volcanoes, respectively. Additionally, the chemical composition of melt and fluid inclusions supports the identification of another magmatic series in the area. We suggest that the related alkali basaltic melts were produced by contemporary melting of a less enriched domain in the upper lithosphere, a process that is more commonly documented in mature rifts. Various extents of mixing and crystallization of these three distinct magmatic series occur in the lower crust beneath the VVP where the barometric estimates are consistent with the presence of a thick seismic low velocity zone (LVZ). The involvement of alkali basaltic melts in the regional magma production would be also detected in the spread of gas emissions in the rift valley and in the fumaroles of the main active volcanoes. Melting of the corresponding mantle domain is an added source of gas release that may largely contribute to CO₂ emissions along the EARS.

Key words: Continental rift; melt and fluid inclusions; degassing

INTRODUCTION

Magmatic activity in continental rifting handles some of the most remarkable volcanic manifestations on Earth, including emission of hyperalkaline undersaturated and carbonatitic volcanism (Rooney, 2020a), and massive CO₂ degassing (Wong *et al.*, 2019). However, the origin and evolution of the involved magmas as well as the architecture of associated magmatic systems (especially the depth of magma ponding zones) are still poorly understood with respect to other geodynamical contexts (Biggs *et al.*, 2021).

The East African Rift System (EARS) is regarded as the current largest example of continental break-up on Earth with magmatotectonic events starting about 45 Myr on the Eastern branch (Michon *et al.*, 2022 and references therein). It evolved from trap-scale to plate-scale rifting under the combined effect of an overall complex mantle upwelling dynamics and of extensive stress

acting on the African lithosphere (Michon *et al.*, 2022). The origin of the large chemical variability of associated lavas erupted is still a matter of debate, and several models have been formulated over the years, including various extents of mixing of distinct components (Rooney, 2020a), such as (1) a convecting mantle with different mantle plumes or thermal anomalies rooted in the African Large Low Shear Velocity Province, together with a depleted mantle endmember (DMM) and (2) a sub-continental lithospheric mantle (SCLM) with both a common lithospheric mantle (CLM) component partially metasomatized by the agents extracted from the convecting mantle, and a Pan-African (PA) lithospheric domain that is enriched in metasomes likely deriving from previous subduction events.

Magmatic activity on the Western branch started about 25 Myr ago and concentrated in four main volcanic provinces, which

are from North to South (Michon *et al.* (2022) and references therein): Toro Ankole, Virunga, Kivu, and Rungwe. Carbonatitic, silica undersaturated hyperalkaline and tholeiitic lavas are predominant with a prominent increase in SiO₂ and depletion in K₂O from Toro Ankole to Kivu, which reflects a progressive decrease of the melting depth towards the South (Furman & Graham, 1999; Rosenthal *et al.*, 2009). The Virunga Volcanic Province (VVP) is currently the sole volcanic province of the Western branch characterized by ongoing eruptive activity. The VVP is characterized by a massive gas release from both volcanic gas plumes (CO₂, SO₂, and halogens) and soil degassing (CO₂) across the rift valley (Tedesco *et al.*, 2010; Bobrowski *et al.*, 2017a, b; Fischer *et al.*, 2019). The VVP is also known for its high potassic magmatism, making it one of the largest within-plate potassic provinces on Earth (Minissale *et al.*, 2022). Two main magmatic series are coexisting (Poucllet *et al.*, 2016): leucite-bearing basanite and leucite–melilite-bearing nephelinite. They are best represented by the eruptive activity that occurred over 12 ka at the Nyamulagira and Nyiragongo volcanoes, respectively. Previously, it has been proposed that those magmas were related to mantle plume sources (Chakrabarti *et al.*, 2009; Castillo *et al.*, 2014). Recent studies have rather supported the melting of a heterogeneous metasomatized SCLM by thinning-driven adiabatic decompression (Poucllet *et al.*, 2016; Muravyeva *et al.*, 2021; Pitcavage *et al.*, 2021; Boudoire *et al.*, 2022; Minissale *et al.*, 2022). However, the reason for the slight enrichment in primordial helium (³He), leading to ³He/⁴He values higher than those expected from SCLM melting, in some gaseous emissions along the Western branch is still poorly understood (Tassi *et al.*, 2009; Tedesco *et al.*, 2010; Castillo *et al.*, 2014; Boudoire *et al.*, 2022).

Most studies that tried to elucidate the origin of the chemical variability of magmas in continental rifting were based either on a compilation of bulk rock analyses for the entire EARS or on the study of melt inclusions entrapped in crystals hosted by lavas from a single volcanic edifice (see Appendix A for references). In this study, we present the analyses of both melt and fluid inclusions from distinct eruptive centres in the VVP and discuss them in a global scheme including earlier analyses of mafic melt inclusions obtained for other volcanic edifices along the EARS (Fig. 1a). This innovative approach of coupling information from melt (major, volatile, and trace elements) and fluid (noble gases, barometry) inclusions from the same products offer a unique opportunity to constrain magmatic processes in the VVP and more broadly those involved in continental rifting.

SAMPLING STRATEGY AND RESULTS

Lava flow samples

Lavas flows from five eruptive sites were collected in the VVP (Fig. 1b): a melilite nephelinite from the Nyiragongo summit lava lake in 2020, a potassic basanite from the Nyamulagira summit eruption in 2020, as well as two olivine nephelinite or melilitites samples (Mudjoga, Kashaka), and an alkali basalt (Rumoka) from peripheral eruptive cones (<12 ka) in the rift valley (Poucllet *et al.*, 2016; Poucllet & Bram, 2021; Minissale *et al.*, 2022; Molendijk *et al.*, 2024). The major element composition of the bulk rocks mostly overlaps within the chemical variability previously documented in the area (Fig. 1c), in particular the two main nephelinitic and basanitic magmatic series reported in the VVP (Poucllet *et al.*, 2016).

Composition of melt and fluid inclusions

Melt and fluid inclusions in olivine, clinopyroxene, and nepheline crystals from the collected lava flows were analysed (see Appendix A for the full methodology and Appendix B for the

dataset). Melt and fluid inclusions in high (Fo_{87–90}; where Fo is the forsterite content in molar proportions), and low (Fo_{75–85}) magnesian olivine crystals were distinguished.

Three endmembers that frame the whole chemical variability in major, trace, and volatile elements were identified (Fig. 2; Appendixes B and C). (1) The 'Nyiragongo-type' nephelinitic melt exhibits the lowest Na₂O/K₂O (= 0.46), and the highest Cl (2352 ppm) and F (4866 ppm) contents. The enrichment in more incompatible elements with respect to less incompatible ones is significant (La/Yb=59.1). Noble gases measurements are not reported due to the existence of a diffusive loss and isotopic fractionation involving especially ³He and ⁴He. (2) The 'Nyamulagira-type' basanitic melt is characterized by Na₂O/K₂O=1.1, high Cl (1249 ppm) and F (2145 ppm) contents as well as intermediate ratios between more incompatible and less incompatible trace elements (La/Yb=37.1). Noble gases systematics show relatively low ³He/⁴He (Rc/Ra=7.8), ⁴He/⁴⁰Ar* (= 0.2), but a high CO₂/³He (= 7.9 × 10⁹). (3) The 'Rumoka-type' basaltic melt (in primitive Fo_{89–90} olivine crystals) is characterized by the highest Na₂O/K₂O (= 1.4) and the lowest Cl (641 ppm) and F (1038 ppm) contents. The ratio between light (LREE) and heavy (HREE) rare earth elements is the lowest (La/Yb=22.1). Noble gases show the highest ³He/⁴He (Rc/Ra=8.1), ⁴He/⁴⁰Ar* (= 3.4), and lowest CO₂/³He (= 3.5 × 10⁸). Strong variations of highly incompatible elements ratios (such as Ce/Pb, Nb/La, Th/U, Zr/Hf) are observed among these melts. The chemical composition of melt inclusions in olivine crystals from the Mudjoga and Kashaka lavas ranges between a 'Nyiragongo-type' nephelinitic melt and a 'Rumoka-type' basaltic melt (Fig. 2c, d; Appendix C). The chemical composition of melt and fluid inclusions in low magnesian olivine crystals from Rumoka lavas diverges from the composition of a 'Rumoka-type' basaltic melt towards a 'Nyamulagira-type' basanitic melt (Head *et al.*, 2011).

Barometric estimates

The dissolved CO₂ content in melt inclusions ranges from 0.11 to 0.31 wt % in 'Nyamulagira-type' basanitic melt to 0.28 to 0.35 wt % in 'Nyiragongo-type' nephelinitic melt and up to 1.69 wt % in 'Rumoka-type' basaltic melt. In melt inclusions from other peripheral products (Kashaka and Mudjoga), the dissolved CO₂ content shows intermediate values (0.13–0.77 wt %) like the ones (0.29–0.88 wt %) measured in the more evolved melt inclusions at Rumoka (Fo₈₃ olivine crystal). The CO₂ density in fluid inclusions for these peripheral products ranges from 0.57 to 0.94 g/cc (Appendix B).

Corresponding barometric estimates allow us to define four pressure ranges (Fig. 3). (1) The highest pressures (870–700 MPa) are recorded for Rumoka and Kashaka peripheral products. Intermediate pressures are centred on two modes at (2) 540–450 MPa and (3) 230–330 MPa for Nyamulagira products (mirroring a differentiation trend from Fo_{79.6} to Fo_{75.6}) and for all peripheral products. (4) The lowest pressures (70–130 MPa) are recorded in melt inclusions from Nyiragongo (in nepheline crystals) and partially from Mudjoga. These barometric estimates agree with earlier results based on petrological, geochemical, and geophysical methods (Louaradi *et al.*, 1993; Poucllet & Bram, 2021; Molendijk *et al.*, 2024) which suggest at least three major magma ponding zones beneath the VVP in similar pressure ranges.

DISCUSSION

For a single eruptive centre, the decrease in MgO, CaO/Al₂O₃, and Sc correlated to an increase in incompatible elements such




Fig. 1. (A) Location of the volcanoes of the East African Rift System (EARS) for which basic melt inclusions were documented and compiled in this study. Continuous lines show the main EARS active segments (Michon *et al.*, 2022). (B) Location of the sampling sites in the Virunga Volcanic Province (VVP) for the melt and fluid inclusions investigation of this study. (C) Composition of the host lavas (coloured dots refer to the lava sampled in the VVP) and comparison with literature data (coloured fields refer to the lava sampled along the EARS, see Appendix A for references). Service Credit Layers: Landsat 8, Google Earth.

as K and Th suggest that olivine-clinopyroxene crystallization accounts for most of the chemical evolution in major and trace elements from the primitive to the more evolved melt inclusions (Fig. 2a; Appendix C). This predominant role of olivine-clinopyroxene crystallization (followed by feldspathoids in shallower magma reservoirs) in the chemical variability was previously documented for lavas erupted in the surroundings of Nyiragongo (Pitcavage *et al.*, 2021; Molendijk *et al.*, 2024). However, the crystallization of the main mineralogical phases reported in VVP magmas cannot explain the chemical variability observed between the chemical endmembers (Minissale *et al.*, 2019), nor that observed on certain highly incompatible trace elements ratios (such as Th/U) between primitive and more evolved melt inclusions at Rumoka. Our data also confirm that

a significant role of crustal contamination may be discarded, consistent with the absence of Ce/Pb and Nb/La anomalies (Fig. 2b; Chakrabarti *et al.*, 2009; Minissale *et al.*, 2022) and thus in favour of heterogeneous mantle sources.

Melting of metasomes in the deep lithospheric mantle

The 'Nyiragongo-type' nephelinitic melt has a clear affinity with a carbonatitic component (high Zr/Hf and low Th/U, Ti/Eu), as previously suggested in the area (Chakrabarti *et al.*, 2009; Pouclet *et al.*, 2016), and for nephelinitic melts emitted at Ol Doinyo Lengai (Fig. 2c; Appendix C). Lithium-enrichment in the 'Nyamulagira-type' basanitic melt is consistent with a preferential affinity for the global average of subducted sediment (GLOSS-II component)




Fig. 2. (A) SiO_2 vs. K_2O in melt inclusions with the composition of the main mineralogic phases in the VVP indicated with crosses (Cpx. stands for clinopyroxene). (B) Nb/La vs. Ce/Pb , (C) Zr/Hf vs. Th/U , and (D) Li/Yb vs. Nb/U in melt inclusions compared with the composition of global endmembers showed with squares (see Appendix A for references). (E) K/Yb vs. Dy/Yb and (F) Ba/Rb vs. Rb/Sr in melt inclusions with the expected effect of amphibole, phlogopite, garnet and spinel in the melted mantle source (Furman & Graham, 1999). Numbers show the melting degree on the theoretical mantle sources melting paths. Coloured fields refer to the composition of basic melt inclusions from the EARS documented in the literature (see Appendix A for references).

as previously suggested also for the Bufumbira volcanic field in the VVP (Fig. 2d; Hudgins et al., 2015). While subducted sediments are thought to play a negligible role in the genesis of VVP magmas (Minissale et al., 2022), past episodes of mantle metasomatism by slab-derived fluids have led to preferential enrichment in some elements (Taniuchi et al., 2024), as attested by the elevated Li/Yb

in the Bufumbira volcanic field (Hudgins et al., 2015), and the lower $\delta^{13}\text{C}$ values of CO_2 in gaseous emissions at Nyamulagira (Boudoire et al., 2022). The chemical composition of melt inclusions also highlights a preferential signature of garnet over spinel (enrichment in LREE and MREE with respect to HREE; Fig. 2e) and of phlogopite over amphibole (low Ba/Rb , $\text{Na}_2\text{O/K}_2\text{O}$, and high Rb/Sr ;




Fig. 3. Pressure and depth estimates of magma ponding zones beneath the VVP from CO₂ densimetry in fluid inclusions (FI) and CO₂-H₂O equilibrium in melt inclusions (MI). Earlier data from literature (Lit.) for Nyamulagira and Nyiragongo about potential magma ponding zones (shaded coloured ellipsoids) and the crust structure (low velocity zones, crust–mantle transition zone) are listed in Appendix A. The melt inclusions selected as chemical endmembers in this study are highlighted: (1) for Nyiragongo-type melt, (2) for Nyamulagira-type melt, and (3) for Rumoka-type melt.

Fig. 2f) in the mantle sources of the two related magmatic series (Chakrabarti *et al.*, 2009; Condomines *et al.*, 2015; Muravyeva *et al.*, 2021; Minissale *et al.*, 2022).

In summary, our new data from melt inclusions entirely agree with the melting of two distinct types of metasomatized lithologies in the mantle beneath the Nyiragongo and Nyamulagira volcanoes ('Type V magmas'; Rooney, 2020a). The generation of a Nyiragongo primary olivine melilitic melt (leading to present nephelinitic melt by differentiation) agrees with the melting of a carbonated metasome in the garnet–phlogopite–carbonate stability field at depth > 100 km, close to the base of the lithosphere (140–160 km-depth; Chakrabarti *et al.*, 2009; Muravyeva *et al.*, 2021; Minissale *et al.*, 2022). Conversely, a distinct metasome affected by old slab fluids is expected to be at the origin of the Nyamulagira primary basanitic melt (Chakrabarti *et al.*, 2009; Minissale *et al.*, 2022).

Contemporary melting of an upper lithospheric mantle domain

The chemical composition of melt inclusions analysed in this study also clearly highlights the contemporary production of a third magmatic series (the 'Rumoka-type' basaltic melt) not yet documented in the VVP but reminiscent of 'Type IIb magmas' (Rooney, 2020a). The similarity of chemical compositions in trace elements between the related melt inclusions and those from the 2011 eruption at Nabro in the Afar region is striking (Fig. 2; Appendix C). These melts are ubiquitous along the EARS and are considered to be produced at the beginning of the transition towards mature rifts by thermobaric destabilization of a thick lithosphere (Rooney, 2020a, 2020b; Chiasera *et al.*, 2021). Along the

Western branch, these melts would be produced by melting of a less enriched lithospheric mantle domain (Chakrabarti *et al.*, 2009; Rosenthal *et al.*, 2009; Pouclet *et al.*, 2016).

The Na₂O/K₂O ratio above the unit in the primitive basaltic melt inclusions at Rumoka suggests the presence of amphibole (together with phlogopite) in the mantle source, a mineral characteristic of a shallower lithospheric melting domain (60–80 km depth; Rosenthal *et al.*, 2009; Muravyeva *et al.*, 2021). The melting of a similar shallow mantle domain was consistently proposed for the genesis of alkali basalts of similar composition in the neighbouring Kivu volcanic province (Rosenthal *et al.*, 2009) and in the genesis of Type IIb magmas (Rooney, 2020a). In the VVP, it supports the idea of contemporary melting of distinct lithospheric mantle domains involved in magma genesis. A heterogeneous SCLM with a deeper lithospheric mantle hosting metasomes (U-Th enriched lithologies) and a less enriched upper lithospheric mantle could be a reasonable explanation for the slight ³He/⁴He variability seen between Nyiragongo, Nyamulagira, and Rumoka. This hypothesis agrees with the observation that metasomatized lithologies are preferentially found near the base of the lithosphere along the EARS (Rooney, 2020a). Such vertical partition has also been suggested beneath the North Tanzanian Divergence (NTZ) of the Eastern branch, where a LVZ has been imaged at a mid-lithospheric depth (~60 km) separating the two mantle components (Plasman *et al.*, 2017). Whatever the case, the ³He/⁴He values remain within the range of values classically attributed to the DMM (8 ± 1 Ra; Tedesco *et al.*, 2010) and confirm the major contribution of the upper mantle in the genesis of the fluids under this part of the EARS (Aiuppa *et al.*, 2021; Boudoire *et al.*, 2022).

Magma mixing of distinct magmatic series

The primitive melt inclusions entrapped in the deepest roots of the magmatic systems beneath the VVP, i.e. at the crust-mantle transition zone (Muravyeva *et al.*, 2021) (Fig. 3), and those of Nyiragongo and Nyamulagira, better preserve the chemical signature of distinct magmatic series. Conversely, more evolved melt inclusions from peripheral eruptive vents have a hybrid chemical composition between these magmatic series (Fig. 2). We suggest that this fingerprint results from variable mixing of the distinct identified magmatic series in intermediate crustal magma ponding zones. In fact, we document two intermediate crustal pressure modes (540–450 MPa and 230–330 MPa) for these peripheral products (Fig. 3) that match with the top of the LVZs beneath the VVP (Mavonga *et al.*, 2010; Pouclet & Bram, 2021) with the main one extending from 18 to 28 km depth (Muravyeva *et al.*, 2021). These LVZs image magma reservoirs or several melt-rich sills where magma mixing is eased (Muravyeva *et al.*, 2021). Complex crystal zoning and textures reported in some peripheral lavas (Molendijk *et al.*, 2024) partially highlight this mixing process. A similar LVZ linked to magmatic processes has also been imaged in the lower crust (15–35 km depth) beneath the NTZ of the Eastern branch (Plasman *et al.*, 2017).

Insights on current degassing processes

Nyiragongo and Nyamulagira are among the strongest volcanic gas emitters on Earth (Fischer *et al.*, 2019). Degassing paths modelled from the initial composition of melt inclusions (see Appendix A for the methodology) were compared to the temporal evolution of CO₂/SO₂ and Cl/S from gas plumes composition reported in literature (Bobrowski *et al.*, 2017; Boudoire *et al.*, 2022). This period overlaps with the progressive replenishment of the Nyiragongo lava lake between the two flank eruptions in 2002 and 2021 (Burgi *et al.*, 2020) and the renewal of lava lake activity at Nyamulagira since 2012 (Burgi *et al.*, 2021). The increase in CO₂/SO₂ and Cl/S values suggests that magma arrival and degassing in the deeper crustal magmatic reservoirs (>250 MPa) prevail on magma degassing in the shallowest ones (<150 MPa; Fig. 4a, b) over time. At Nyiragongo, the existence of various degassing magmatic sources involved in the outgassing of the lava lake was previously proposed (Bobrowski *et al.*, 2017) and the exceptional occurrence of deep mush material in the 2021 lava from Nyiragongo (Molendijk *et al.*, 2024) supports this hypothesis.

We also note that the ³He/⁴He values in Nyiragongo fumaroles may have been slightly lower in 2020 (Rc/Ra = 7.3 ± 0.2) (Boudoire *et al.*, 2022), preceding the 2021 flank eruption than during the 2003–2007 period (Rc/Ra = 7.8 ± 0.8 and up to 8.7; Tedesco *et al.*, 2010) (Fig. 4c). This behaviour mirrors the increase of magma flux feeding the lava lake on the same time span (Burgi *et al.*, 2020, 2021). This slight decrease in ³He/⁴He values over time, which does not depend on pressure (and is, therefore, not linked to the deepening of the degassing magmatic sources over time), highlights a potential ⁴He enrichment from 2003 to 2020. It could reflect an increase of the magma production (and influx) from the Nyiragongo deep metasomatized mantle source that is enriched in U–Th (and thus ⁴He) with respect to the other endmembers described in this study. Conversely, it is worth noting the chemical affinity (higher Rc/Ra and ⁴He/⁴⁰Ar* coupled with low CO₂/³He and δ¹³C) of regional dry CO₂ gas emissions (e.g. mazuku on Fig. 4c) from the rift valley with the isotopic signature of fluid inclusions in crystals from Rumoka products. High ⁴He/⁴⁰Ar* and low CO₂/³He and δ¹³C are reminiscent of more degassed magmas (Boudoire *et al.*, 2022) and may suggest that steady-state




Fig. 4. Pressure dependence of CO₂/SO₂ and Cl/S molar ratios in gas plumes emitted at (A) Nyiragongo and (B) Nyamulagira. Degassing paths were modelled using SulfurX (Ding *et al.*, 2023) and SolEx (Witham *et al.*, 2012) considering both Fractional Equilibrium Degassing (FED) and Batch Equilibrium Degassing (BED). See Appendix A for more details about starting conditions and parameters used in the modelling. Values framed on the x-axis highlight (i) averaged measurements made in volcanic gas plumes from 2005 to 2014 (Sawyer *et al.*, 2008; Bobrowski *et al.*, 2017a, b) and (ii) minimum and maximum values measured in 2020 (Boudoire *et al.*, 2022). The theoretical pressure ranges corresponding to these measurements are highlighted on the y axis (shaded boxes) as well as the corresponding magma ponding zones (circles) from Fig. 2. (C) CO₂/³He vs. Rc/Ra in fluid inclusions with dashed black lines evidencing mixing of a mantle endmember (Rumoka-type melt) and various crustal components (Barry *et al.*, 2013). The stars show the average composition of the fumaroles at Nyiragongo and Nyamulagira (Tedesco *et al.*, 2010; Boudoire *et al.*, 2022). CO₂/³He vs. Th/U with dashed lines showing mixing between (1) a Nyiragongo-type melt and either (2) a Nyamulagira-type melt or (3) a Rumoka-type melt.

outgassing through the rift valley is mostly linked to magma storage and degassing in the LVZs that could be preferentially fed by the melting of the upper and less enriched lithospheric mantle domain (Tedesco *et al.*, 2010; Boudoire *et al.*, 2022).

CONCLUSIONS

Melt and fluid inclusions suggest that current magma production beneath the VVP is not solely linked to the melting of various mantle metasomes that preferentially feed the main volcanic edifices, as documented for immature rifts (Rooney, 2020a; Aiuppa *et al.*, 2021). The contemporary melting of a less enriched lithospheric domain, isolated at a shallower level in the mantle, is shown by the composition of primitive melt inclusions from peripheral products erupted in the rift valley. This process is rather reminiscent of magmatic production at the beginning of the transition towards mature rifts. We propose that steady-state degassing associated with continental rifting in the area is preferentially caused by magma production in the upper part of the lithosphere (at 60–80 km-depth), followed by mixing and storage of the melt in the lower crust (at 15–35 km-depth), where LVZs are imaged, rather than by the melting of isolated deep lithospheric metasomes (Muirhead *et al.*, 2020; Aiuppa *et al.*, 2021). This model would preferentially account for global CO₂ emissions along the EARS with respect to volcanic emissions (Wong *et al.*, 2019) and for the large chemical variability of erupted melts (Rooney, 2020a).

SUPPLEMENTARY MATERIAL AND DATA AVAILABILITY STATEMENT

The Supplemental Material is composed of four appendices. Appendix A describes the methodology used in this study. It provides details about the analytical conditions of measurements, data treatment, degassing models, and comparative data from literature. Appendix B is the complete dataset of measurements acquired for this study. Appendix C is composed of four complementary figures that sustain the discussion about the role of crystallization, contamination, mixing processes, and mantle sources lithologies in the chemical variability of melt inclusions. Appendix D is the completed repository template available on a recognized domain repository of the GFZ Data Services through the link <https://dataservices.gfz-potsdam.de/panmetaworks/review/0987bbb94d29daa7e406e048ef73614579375334e467578be75da9e606a69d78/> with a registered DOI (<https://doi.org/10.5880/figdeo.2024.029>) (Boudoire, 2024) for accessing the dataset and the data description in accordance with the data availability statement.

ACKNOWLEDGEMENTS

We are grateful to the Goma Volcano Observatory (OVG) for the invitation and logistic help. The United Nations program of MONUSCO is warmly thanked for the great support in the field deployment by helicopter. The ICCN is acknowledged for providing the logistic inside the National Park of Virunga. The SVG team is thanked for their priceless support in the coordination of the descent into the craters. We are in debt with M. Benbakkar, J.-L. Devidal, C. Fonquernie, F. Schiavi, E. Voyer at the LMV, E. Deloule, L. Piani at the CRPG, E. Rose-Koga at ISTO, and M. Tantillo at the INGV for their assistance during sample preparation and analysis. T. Rooney, A.Z. Tadesse, and V. Zanon are acknowledged for their constructive review of the article. This research was funded by the

French Government Laboratory of Excellence initiative n° ANR-10-LABX-0006, the Region Auvergne and the European Regional Development Fund through the project DegazRift. We thank the Istituto Nazionale di Geofisica e Vulcanologia, Italy, grant “Progetto INGV Pianeta Dinamico” - code CUP D53J19000170001 - funded by the Italian Ministry MIUR (“Fondo Finalizzato al rilancio degli investimenti delle amministrazioni centrali dello Stato e allo sviluppo del Paese”, legge 145/2018). D. Tedesco was funded by the UN-MONUSCO during this project. This is contribution no. 673 of the ClerVolc program of the International Research Center for Disaster Sciences and Sustainable Development of the University Clermont Auvergne.

REFERENCES

- Aiuppa, A., Casetta, F., Coltorti, M., Stagno, V. & Tamburello, G. (2021). Carbon concentration increases with depth of melting in Earth's upper mantle. *Nature Geoscience* **14**(9), 697–703. <https://doi.org/10.1038/s41561-021-00797-y>.
- Barry, P. H., Hilton, D. R., Fischer, T. P., de Moor, J. M., Mangasini, F., & Ramirez, C. (2013). Helium and carbon isotope systematics of cold “mazuku” CO₂ vents and hydrothermal gases and fluids from Rungwe Volcanic Province, southern Tanzania. *Chemical Geology*, **339**, 141–156. <https://doi.org/10.1016/j.chemgeo.2012.07.003>.
- Biggs, J., Ayele, A., Fischer, T. P., Fontijn, K., Hutchison, W., Kazimoto, E. & Wright, T. J. (2021). Volcanic activity and hazard in the East African rift zone. *Nature Communications* **12**(1), 6881. <https://doi.org/10.1038/s41467-021-27166-y>.
- Bobrowski, N., Giuffrida, G. B., Arellano, S., Yalire, M., Liotta, M., Brusca, L., Calabrese, S., Scaglione, S., Rüdiger, J., Castro, J. M., Galle, B., & Tedesco, D. (2017a). Plume composition and volatile flux of Nyamulagira volcano, Democratic Republic of Congo, during birth and evolution of the lava lake, 2014–2015. *Bulletin of Volcanology*, **79**(12). <https://doi.org/10.1007/s00445-017-1174-0>.
- Bobrowski, N., Giuffrida, G. B., Yalire, M., Lübcke, P., Arellano, S., Balagizi, C., Calabrese, S., Galle, B., & Tedesco, D. (2017b). Multi-component gas emission measurements of the active lava lake of Nyiragongo, DR Congo. *Journal of African Earth Sciences*, **134**, 856–865. <https://doi.org/10.1016/j.jafrearsci.2016.07.010>.
- Boudoire, G. (2024). Dataset of lava samples, melt and fluid inclusions analysis from the Virunga Volcanic Province (DR Congo). *GFZ Data Services*. <https://doi.org/10.5880/figdeo.2024.029>.
- Boudoire, G., Giuffrida, G., Liuzzo, M., Bobrowski, N., Calabrese, S., Kuhn, J. & Tedesco, D. (2022). Chemical variability in volcanic gas plumes and fumaroles along the East African Rift System: new insights from the Western branch. *Chemical Geology* **596**, 120811. <https://doi.org/10.1016/j.chemgeo.2022.120811>.
- Burgi, P. Y., Boudoire, G., Rufino, F., Karume, K. & Tedesco, D. (2020). Recent activity of Nyiragongo (Democratic Republic of Congo): new insights from field observations and numerical modeling. *Geophysical Research Letters* **47**(17), e2020GL088484. <https://doi.org/10.1029/2020GL088484>.
- Burgi, P. Y., Valade, S., Coppola, D., Boudoire, G., Mavonga, G., Rufino, F. & Tedesco, D. (2021). Unconventional filling dynamics of a pit crater. *Earth and Planetary Science Letters* **576**, 117230. <https://doi.org/10.1016/j.epsl.2021.117230>.
- Castillo, P. R., Hilton, D. R. & Halldórsson, S. A. (2014). Trace element and Sr–Nd–Pb isotope geochemistry of Rungwe Volcanic Province, Tanzania: implications for a superplume source for East Africa Rift magmatism. *Frontiers in Earth Science* **2**, 21. <https://doi.org/10.3389/feart.2014.00021>.
- Chakrabarti, R., Basu, A. R., Santo, A. P., Tedesco, D. & Vaselli, O. (2009). Isotopic and geochemical evidence for a heterogeneous

- mantle plume origin of the Virunga volcanics, Western rift, East African Rift system. *Chemical Geology* **259**(3–4), 273–289. <https://doi.org/10.1016/j.chemgeo.2008.11.010>.
- Chiasera, B., Rooney, T. O., Bastow, I. D., Yirgu, G., Grosfils, E. B., Ayalew, D. & Ramsey, M. (2021). Magmatic rifting in the Main Ethiopian Rift began in thick continental lithosphere; the case of the Galema Range. *Lithos* **406–407**, 106494. <https://doi.org/10.1016/j.lithos.2021.106494>.
- Condomines, M., Carpentier, M. & Ongendangenda, T. (2015). Extreme radium deficit in the 1957 AD Mugogo lava (Virunga volcanic field, Africa): its bearing on olivine–mellilite genesis. *Contributions to Mineralogy and Petrology* **169**, 1–19. <https://doi.org/10.1007/s00410-015-1124-9>.
- Ding, S., Plank, T., Wallace, P. J., & Rasmussen, D. J. (2023). Sulfur_X: A Model of Sulfur Degassing During Magma Ascent. *Geochemistry, Geophysics, Geosystems*, 24(4). Portico. <https://doi.org/10.1029/2022gc010552>.
- Fischer, T. P., Arellano, S., Carn, S., Aiuppa, A., Galle, B., Allard, P. & Chiodini, G. (2019). The emissions of CO₂ and other volatiles from the world's subaerial volcanoes. *Scientific Reports* **9**(1), 18716. <https://doi.org/10.1038/s41598-019-54682-1>.
- Furman, T. & Graham, D. (1999) Erosion of lithospheric mantle beneath the East African Rift system: geochemical evidence from the Kivu volcanic province. In: *Developments in Geotectonics* (Vol. 24. Elsevier, pp.237–262.
- Head, E. M., Shaw, A. M., Wallace, P. J., Sims, K. W. & Carn, S. A. (2011). Insight into volatile behavior at Nyamuragira volcano (DR Congo, Africa) through olivine-hosted melt inclusions. *Geochemistry, Geophysics, Geosystems* **12**(10). <https://doi.org/10.1029/2011GC003699>.
- Hudgins, T. R., Mukasa, S. B., Simon, A. C., Moore, G. & Barifajjo, E. (2015). Melt inclusion evidence for CO₂-rich melts beneath the western branch of the East African Rift: implications for long-term storage of volatiles in the deep lithospheric mantle. *Contributions to Mineralogy and Petrology* **169**, 1–18.
- Louaradi, D., Clochiatti, R., Pineau, F. & Javoy, M. (1993) Magma storage beneath Nyiragongo volcano (Zaire): evidence from fluid and melt inclusions. In: *Terra Abstracts* Vol. 5, p.573.
- Mavonga, T., Zana, N. & Durrheim, R. J. (2010). Studies of crustal structure, seismic precursors to volcanic eruptions and earthquake hazard in the eastern provinces of the Democratic Republic of Congo. *Journal of African Earth Sciences* **58**(4), 623–633. <https://doi.org/10.1016/j.jafrearsci.2010.08.008>.
- Michon, L., Famin, V. & Quidelleur, X. (2022). Evolution of the East African rift system from trap-scale to plate-scale rifting. *Earth-Science Reviews* **231**, 104089. <https://doi.org/10.1016/j.earscirev.2022.104089>.
- Minissale, S., Zanetti, A., Tedesco, D., Morra, V. & Melluso, L. (2019). The petrology and geochemistry of Nyiragongo lavas of 2002, 2016, 1977 and 2017 AD, and the trace element partitioning between mellilite glass and melilite, nepheline, leucite, clinopyroxene, apatite, olivine and Fe–Ti oxides: a unique scenario. *Lithos* **332–333**, 296–311. <https://doi.org/10.1016/j.lithos.2019.02.023>.
- Minissale, S., Casalini, M., Cucciniello, C., Balagizi, C., Tedesco, D., Boudoire, G. & Melluso, L. (2022). The geochemistry of recent Nyamulagira and Nyiragongo potassic lavas, Virunga Volcanic Province, and implications on the enrichment processes in the mantle lithosphere of the Tanzania–Congo craton. *Lithos* **420–421**, 106696. <https://doi.org/10.1016/j.lithos.2022.106696>.
- Molendijk, S. M., Namur, O., Kamate Kaleghetso, E., Mason, P. R., Smets, B., Vander Auwera, J. & Neave, D. A. (2024). Plumbing system architecture and differentiation processes of the Nyiragongo volcano, DR Congo. *Journal of Petrology* **65**(1), egad088. <https://doi.org/10.1093/petrology/egad088>.
- Muirhead, J. D., Fischer, T. P., Oliva, S. J., Laizer, A., van Wijk, J., Currie, C. A. & Ebinger, C. J. (2020). Displaced cratonic mantle concentrates deep carbon during continental rifting. *Nature* **582**(7810), 67–72. <https://doi.org/10.1038/s41586-020-2328-3>.
- Muravyeva, N. S., Senin, V. G., Ivanov, A. V. & Belyatsky, B. V. (2021). Leucite basanites of Virunga (east African rift): some insights into petrogenesis and source composition. *Lithos* **384–385**, 105972. <https://doi.org/10.1016/j.lithos.2021.105972>.
- Pitcavage, E., Furman, T., Nelson, W. R., Kalegga, P. K. & Barifajjo, E. (2021). Petrogenesis of primitive lavas from the Toro Ankole and Virunga Volcanic Provinces: metasomatic mineralogy beneath East Africa's Western rift. *Lithos* **396–397**, 106192. <https://doi.org/10.1016/j.lithos.2021.106192>.
- Plasman, M., Tiberi, C., Ebinger, C., Gautier, S., Albaric, J., Peyrat, S. & Gama, R. (2017). Lithospheric low-velocity zones associated with a magmatic segment of the Tanzanian Rift. *East Africa. Geophysical Journal International* **210**(1), 465–481. <https://doi.org/10.1093/gji/ggx177>.
- Poucllet, A. & Bram, K. (2021). Nyiragongo and Nyamuragira: a review of volcanic activity in the Kivu rift, western branch of the East African Rift System. *Bulletin of Volcanology* **83**(2), 10. <https://doi.org/10.1007/s00445-021-01435-6>.
- Poucllet, A., Bellon, H. & Bram, K. (2016). The Cenozoic volcanism in the Kivu rift: assessment of the tectonic setting, geochemistry, and geochronology of the volcanic activity in the south-Kivu and Virunga regions. *Journal of African Earth Sciences* **121**, 219–246. <https://doi.org/10.1016/j.jafrearsci.2016.05.026>.
- Rooney, T. O. (2020a). The Cenozoic magmatism of East Africa: part V—magma sources and processes in the East African Rift. *Lithos* **360–361**, 105296. <https://doi.org/10.1016/j.lithos.2019.105296>.
- Rooney, T. O. (2020b). The Cenozoic magmatism of East Africa: part IV—the terminal stages of rifting preserved in the northern East African Rift System. *Lithos* **360–361**, 105381. <https://doi.org/10.1016/j.lithos.2020.105381>.
- Rosenthal, A., Foley, S. F., Pearson, D. G., Nowell, G. M. & Tappe, S. (2009). Petrogenesis of strongly alkaline primitive volcanic rocks at the propagating tip of the western branch of the East African Rift. *Earth and Planetary Science Letters* **284**(1–2), 236–248. <https://doi.org/10.1016/j.epsl.2009.04.036>.
- Sawyer, G. M., Carn, S. A., Tsanev, V. I., Oppenheimer, C., & Burton, M. (2008). Investigation into magma degassing at Nyiragongo volcano, Democratic Republic of the Congo. *Geochemistry, Geophysics, Geosystems*, 9(2). Portico. <https://doi.org/10.1029/2007gc001829>.
- Taniuchi, H., Kawamoto, T., Nakatani, T., Ishizuka, O., Suzuki, T. & Tomiya, A. (2024). Compositional evolution of slab-derived fluids during ascent: implications from trace-element partition between hydrous melts and Cl-free or Cl-rich aqueous fluids. *Contributions to Mineralogy and Petrology* **179**(5), 51. <https://doi.org/10.1007/s00410-024-02122-3>.
- Tassi, F., Vaselli, O., Tedesco, D., Montegrossi, G., Darrah, T., Cuoco, E. & Delgado Huertas, A. (2009). Water and gas chemistry at Lake Kivu (DRC): geochemical evidence of vertical and horizontal heterogeneities in a multibasin structure. *Geochemistry, Geophysics, Geosystems* **10**(2). <https://doi.org/10.1029/2008GC002191>.
- Tedesco, D., Tassi, F., Vaselli, O., Poreda, R. J., Darrah, T., Cuoco, E. & Yalire, M. M. (2010). Gas isotopic signatures (He, C, and Ar) in the Lake Kivu region (western branch of the East African Rift System): geodynamic and volcanological implications. *Journal of Geophysical Research: Solid Earth* **115**(B1). <https://doi.org/10.1029/2008JB006227>.

- Witham, F., Blundy, J., Kohn, S. C., Lesne, P., Dixon, J., Churakov, S. V., & Botcharnikov, R. (2012). SolEx: A model for mixed COHSL-volatile solubilities and exsolved gas compositions in basalt. *Computers & Geosciences*, **45**, 87–97. <https://doi.org/10.1016/j.cageo.2011.09.021>.
- Wong, K., Mason, E., Brune, S., East, M., Edmonds, M. & Zahirovic, S. (2019). Deep carbon cycling over the past 200 million years: a review of fluxes in different tectonic settings. *Frontiers in Earth Science* **7**, 263. <https://doi.org/10.3389/feart.2019.00263>.



Deltech Furnaces

Sustained operating
temperatures to 1800°
Celsius

www.deltechfurnaces.com



Gas Mixing System



An ISO 9001:2015 certified company

Custom Vertical Tube



ASME NQA-1 2008 Nuclear Quality Assurance

Standard Vertical Tube



Control systems are certified by Intertek UL508A compliant

Bottom Loading Vertical Tube