

TRIDs therapy for restoring p53 expression on stop mutated cells

Michele Menditto¹, Davide Ricci¹, Elisa Ciulla¹, Ivana Pibiri¹, Andrea Pace¹, Laura Lentini¹

¹Dipartimento STEBICEF, Università degli Studi di Palermo, Palermo, Italia

Keywords: cancer, readthrough, mutations, therapy

Stop mutations are gene mutations characterized by the substitution of a single nucleotide in the coding sequence of a gene, which causes the onset of a premature stop codon (PTC) within the reading frame of the mRNA, resulting in the formation of a truncated and non-functional protein. This type of mutation accounts for approximately 11% of genetic diseases, including conditions such as Cystic Fibrosis, Duchenne Muscular Dystrophy, Choroideremia, Schwachman-Diamond syndrome, and certain types of hereditary cancers involving mutations in the TP53 gene. About 10% of TP53 mutations are stop mutations [1,2]. TP53 mutations create a favorable environment for tumor formation, and mutant p53 may exhibit loss of function, dominant-negative repression, or gain of oncogenic function, contributing to tumor stability and progression [2]. Today there is no therapy for the pathologies caused by this type of mutation, but an approach that has proven to be particularly effective is represented by molecules with readthrough activity (TRIDs; Translational Readthrough Inducing Drugs) which intervene on the ribosome allowing the overcoming of the PTC and the restoration of the synthesis and subsequent functionality of the protein [3]. In this work, we investigate the effects of TRID molecules with readthrough activity on the TP53 gene in tumor cells, which harbors the PTC R213X, the most common TP53 stop mutation, that generates a truncated and non-functional p53. We analyzed the restoration of p53 protein expression before and after induction of DNA damage by Western blot, its nuclear localization with fluorescence microscopy, the mRNA expression of p53, and its targets p21 and GADD45 to evaluate the functionality of the protein by Real-Time RT PCR. This study represents a promising path for developing targeted cancer therapies against stop mutations, a new approach to impede tumor proliferation, and a solid foundation for the formulation of novel personalized therapy modalities not only against cancer.




Figure 1: Experimental workflow for the study of the p53 rescue in H1299 p53^{R213X} cells by TRIDs molecule treatment

References

- [1] Martins-Dias P, Romão L. Nonsense suppression therapies in human genetic diseases. *Cell Mol Life Sci.* 2021 May;78(10):4677-4701.
- [2] Wang H, Guo M, Wei H, Chen Y. Targeting p53 pathways: mechanisms, structures, and advances in therapy. *Signal Transduct Target Ther.* 2023 Mar 1;8(1):92.
- [3] Fiduccia I, Corrao F, Zizzo MG, Perriera R, Genovese F, Vitale E, Ricci D, Melfi R, Tutone M, Pace A, Lentini L, Pibiri I. Promoting readthrough of nonsense mutations in CF mouse model: Biodistribution and efficacy of NV848 in rescuing CFTR protein expression. *Mol Ther.* 2024 Dec 4;32(12):4514-4523.