

Book of Abstracts



ADVANCES IN MICROBIOLOGY OF **TABLE OLIVES** International Congress

Campus di Via Lanera, Matera

September 1-2, 2025

METAOlive is funded by Ministero dell'Università e della Ricerca and the European Union 2022NN28ZZ and received funding from the European Union Next-GenerationEU, CUP C53D23005460006



Finanziato
dall'Unione europea
NextGenerationEU



Ministero
dell'Università
e della Ricerca



Italiadomani
PIANO NAZIONALE
DI RIPRESA E RESILIENZA



UNIVERSITÀ DEGLI STUDI
DELLA BASILICATA

Microbial innovation in table olive fermentation: a decade of research on *Lactiplantibacillus pentosus* OM13

Francesca, N. (1) *, Alfonzo, A. (1), Aponte, M. (2) La Croce, F. (3), Seminerio, V. (1), Craparo, V. (1), Vagnoli, P. (4), Krieger-Weber, S. (5), Petegolli, D. (3), Gaglio, R. (1), Settanni, L. (1), Moschetti, G. (1)

* corresponding author: nicola.francesca@unipa.it

(1) Department of Agricultural, Food and Forest Sciences (SAAF), University of Palermo, Viale delle Scienze, Bldg. 5, 90128, Palermo, Italy

(2) Department of Agricultural Sciences, University of Naples “Federico II”, Via Università 100, 80055, Portici (Na), Italy

(3) Geolive Belice S.r.l., S.S. 115 Km Dir, Marinella, Castelvetro, 91022, Italy

(4) Lallemand Italia, Via Rossini 14/B, Castel D'Azzano, 37060, Italy

(5) Lallemand, Office Korntal-Münchingen, In den Seiten 53, Korntal-Münchingen, 70825, Germany

Background.

Table olive fermentation is a longstanding biotechnological tradition practice integral to the Mediterranean diet and agri-food sector. Despite its cultural and economic relevance, spontaneous fermentation often suffers from unpredictable microbial dynamics, inconsistent sensory outcomes, and safety concerns due to the presence of spoilage and pathogenic microorganisms. These limitations have generated interest in controlled fermentation strategies based on microbial ecology and starter culture development and application.

At the University of Palermo, the food microbiology research group has undertaken a comprehensive programme aimed at improving the quality, safety, and reproducibility of table olive fermentation. Focusing on Sicilian cultivars, particularly Nocellara del Belice, the research group has explored both Spanish-style and Greek-style fermentation methods. This work contributes to a broader scientific effort to understand and manage microbial communities in fermented vegetables, with a special emphasis on the role of lactic acid bacteria (LAB) and yeasts in shaping product quality and safety ([10.1016/j.ijfoodmicro.2022.109670](https://doi.org/10.1016/j.ijfoodmicro.2022.109670)).

Methods.

Central to this research was the application and characterisation of the autochthonous strain

Lactiplantibacillus pentosus OM13, was consistently applied across all experimental trials. Originally isolated from Sicilian table olives, OM13 was selected for its robust fermentative capabilities, tolerance to high salinity and phenolic compounds, and ability to dominate the microbial ecosystem during fermentation. Its performance was assessed under various conditions, including different harvesting methods (manual vs. mechanical), irrigation regimes, and fermentation strategies. One of the most innovative approaches was the adaptation of the “pied de cuve” technique, commonly used in winemaking, to table olive fermentation. This involves preparing a small volume of partially fermented brine to inoculate larger batches, promoting early establishment of beneficial microbial populations while reducing the need for large quantities of starter cultures. Additional strategies included OM13 acclimatisation in brine before inoculation, supplementation with nutrients to support microbial growth, and co-inoculation with selected yeast strains (*Candida boidinii* and *Candida norvegica*) known for their enzymatic activity and contribution to flavour development. Trials were conducted at both laboratory and industrial scales in collaboration with local producers. Microbial populations were monitored using culture-based methods and molecular profiling such as randomly amplified polymorphic DNA (RAPD)-PCR and 16S rRNA gene sequencing. Physicochemical parameters (pH and salt concentration), volatile organic compounds (VOCs), and sensory attributes were systematically evaluated throughout the fermentation process. Multivariate statistical analyses were used to interpret microbial dynamics and compare fermentation treatments.

Results.

Across all experiments, *Lact. pentosus* OM13 consistently dominated the fermentation microbiota, rapidly acidified the brine, and suppressed spoilage organisms including *Enterobacteriaceae* and *Pseudomonads*. Molecular analyses confirmed that LAB isolates from inoculated trials closely matched the OM13 genetic profile based on RAPD-PCR and 16S rRNA sequencing. The strain demonstrated strong adaptability to diverse environmental conditions, including high salinity, low pH, and phenolic presence. The “pied de cuve” approach proved particularly effective in promoting native LAB growth while reducing reliance on freeze-dried cultures. This approach preserved microbial diversity and enhanced the expression of cultivar-specific sensory traits. The acclimatisation of OM13 prior to inoculation further improved fermentation performance, accelerating acidification and stabilising microbial communities. Nutrient supplementation supported OM13 persistence and improved fermentation kinetics. Sensory evaluations consistently showed that olives fermented with OM13 scored higher in aroma, flavour complexity, and overall acceptability compared to spontaneous fermentations. OM13 use was associated with a favourable VOC profile, including elevated levels of phenylethyl alcohol, ethyl acetate, and hexanal, compounds linked to freshness and green fruit notes. Co-inoculation with selected yeasts contributed to improved colour retention, pulp firmness, and reduced off-flavours.

Inoculated trials consistently demonstrated superior microbiological stability, reduced variability, and enhanced sensory quality across different processing styles and environmental conditions, underscoring the robustness and versatility of OM13 as a starter culture.

Conclusions.

The use of *Lact. pentosus* OM13 has proven to be a reliable and effective strategy for guiding table olive fermentation. Its consistent dominance, environmental resilience, and positive impact on product quality making it a key element in the advancement of table olive fermentation methods. By integrating microbial selection, process optimisation, and sensory science, producers can achieve greater consistency, safety, and authenticity in their products. The development of tailored starter cultures and fermentation aids has advanced microbial biotechnology, offering scalable solutions for both artisanal and industrial production. Moreover, the implementation of OM13 has contributed to the valorisation of local cultivars and traditional methods, enhancing the competitiveness of Sicilian table olives in domestic and international markets. Techniques such as “pied de cuve”, starter acclimatisation, and yeast co-inoculation are replicable and adaptable across the Mediterranean, providing a model for sustainable innovation in fermented vegetable production. This decade-long research effort has significantly deepened scientific understanding of microbial ecology in table olive fermentation while delivering tangible benefits to the agri-food sector. In conclusion, this work serves as a paradigm of how microbiological research can drive innovation, safeguard traditional practices, and contribute to the enhancement of the quality and identity of emblematic Mediterranean fermented foods.