



Total, red and processed meat consumption and human health: an umbrella review of observational studies

Grosso, G., La Vignera, S., Condorelli, R. A., Godos, J., Marventano, S., Tieri, M., Ghelfi, F., Titta, L., Lafranconi, A., Gambera, A., Alonzo, E., Sciacca, S., Buscemi, S., Ray, S., Del Rio, D., & Galvano, F. (2022). Total, red and processed meat consumption and human health: an umbrella review of observational studies. *International Journal of Food Sciences and Nutrition*, 73(6), 1-12. Advance online publication. <https://doi.org/10.1080/09637486.2022.2050996>

[Link to publication record in Ulster University Research Portal](#)

Published in:

International Journal of Food Sciences and Nutrition

Publication Status:

Published online: 15/03/2022

DOI:

[10.1080/09637486.2022.2050996](https://doi.org/10.1080/09637486.2022.2050996)

Document Version

Author Accepted version

For Author Accepted Manuscripts (AAM) published under Ulster University's Rights Retention Policy for Scholarly Works (RRPSW)

When citing an AAM published under Ulster University's RRPSW please use the following citation structure:

Author, A. A. (Year). Title of article. Journal Name, [Accepted Author Manuscript]. PURE Portal URL. Licensed under CC BY 4.0.

General rights

The copyright and moral rights to the output are retained by the output author(s), unless otherwise stated by the document licence.

Unless otherwise stated, users are permitted to download a copy of the output for personal study or non-commercial research and are permitted to freely distribute the URL of the output. They are not permitted to alter, reproduce, distribute or make any commercial use of the output without obtaining the permission of the author(s).

If the document is licenced under Creative Commons, the rights of users of the documents can be found at <https://creativecommons.org/share-your-work/cclicenses/>.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk

Total, red and processed meat consumption and human health: an umbrella review of observational studies

Giuseppe Grosso^{1,2}, Sandro La Vignera³, Rosita A. Condorelli³, Justyna Godos¹, Stefano Marventano⁴, Maria Tieri⁵, Francesca Ghelfi^{2,6}, Lucilla Titta⁵, Alessandra Lafranconi^{7,8}, Angelo Gambera⁹, Elena Alonzo¹⁰, Salvatore Sciacca¹¹, Silvio Buscemi¹², Sumantra Ray^{2,13,14}, Daniele Del Rio^{2,15,16}, Fabio Galvano¹

¹ *Department of Biomedical and Biotechnological Sciences, University of Catania, Catania, Italy;*

² *NNEdPro Global Centre for Nutrition and Health, St John's Innovation Centre, Cambridge, United Kingdom;*

³ *Department of Clinical and Experimental Medicine, University of Catania, Catania, Italy;*

⁴ *Rimini Women's Health, Childhood and Adolescent Department, AUSL Romagna, Rimini, Italy;*

⁵ *SmartFood Program, Department of Experimental Oncology, IEO, European Institute of Oncology IRCCS, Milan, Italy;*

⁶ *Fondazione De Marchi-Department of Pediatrics, IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy;*

⁷ *University of Milano – Bicocca, Milan, Italy;*

⁸ *Care and Public Health Research Institute, Maastricht University, Maastricht, The Netherlands;*

⁹ *Azienda Ospedaliero-Universitaria Policlinico-Vittorio Emanuele, Catania, Italy;*

¹⁰ *Food and Nutrition Security and Public Health Service, ASP Catania, Catania, Italy;*

¹¹ *Mediterranean Oncological Institute (IOM), Catania, Italy;*

¹² *Biomedical Department of Internal and Specialist Medicine (DIBIMIS), University of Palermo, Palermo, Italy;*

¹³ *School of Biomedical Sciences, Ulster University, Ulster, UK;*

¹⁴ *School of the Humanities and Social Sciences, University of Cambridge, Cambridge, UK;*

¹⁵ *Human Nutrition Unit, Department of Food and Drug, University of Parma, Parma, Italy;*

¹⁶ *School of Advanced Studies on Food and Nutrition, University of Parma, Parma, Italy.*

Corresponding author: Giuseppe Grosso, Department of Biomedical and Biotechnological Sciences, University of Catania, Via Santa Sofia 97, 95123 Catania, Italy (email: giuseppe.grosso@unict.it; Phone/Fax +39 0954781187).

Total, red and processed meat consumption and human health: an umbrella review of observational studies

Keywords: total meat; red meat; processed meat; processed foods; meta-analysis; umbrella review; evidence; non-communicable diseases; cardiovascular disease; cancer; coronary heart disease; colorectal cancer

Abstract

Meat consumption has represented an important evolutionary step for humans. However, modern patterns of consumption, including excess intake, type of meat and cooking method have been the focus of attention as a potential cause of rise in non-communicable diseases. The aim of this study was to investigate the association between total, red, and processed meat with health outcomes and to assess the level of evidence by performing a systematic search of meta-analyses of prospective cohort studies. Convincing evidence of the association between increased risk of (i) colorectal adenoma, lung cancer, CHD and stroke, (ii) colorectal adenoma, ovarian, prostate, renal, and stomach cancers, CHD and stroke, and (iii) colon and bladder cancer was found for excess intake of total, red, and processed meat, respectively. Possible negative associations with other health outcomes have been reported. Excess meat consumption may be detrimental to health, with a major impact on cardiometabolic and cancer risk.

Introduction

The inclusion of meat in the diets of human ancestors certainly contributed to human evolution and brain development (Milton 2003). Besides the obvious nutritional advantages in terms of vitamins, minerals, and protein content, primates consuming meat invested a

shorter time for eating after the discovery of fire and cooking, resulting in improvement of their social and manual skills needed for hunting compared to their solely vegetarian counterpart (Carmody & Wrangham 2009). Over the centuries, fresh meat has been typically consumed occasionally, representing a rich course in Southern European regions, while some other populations had a higher consumption of preserved meats whether living in geographical areas with more severe weather (Larsen 2003; Mann 2018). Nowadays, meat consumption is no more affected by nationality or seasonality; meat is generally always available in developed countries and probably over consumed since its introduction into fast foods (Cocking et al. 2020; Henchion et al. 2021). Health surveys and investigations aiming to test the association between meat consumption and health outcomes revealed that excess intake of meat may be detrimental for cardiometabolic diseases and certain cancers (O’Keefe & Cordain 2004). Thus, a great deal of effort has been done to determine whether and how meat could affect human health.

Current research differentiates between red meat, white meat, and processed meat: red meat refers to mammalian muscle meat (i.e., beef, veal, pork, lamb, horse, etc.) rich in myoglobin and heme iron (from which derive the red color); white meat refers to the white counterpart (i.e., chicken, turkey, and other birds); while processed meat refers to red meat or either red or white meat that has been transformed through salting, curing, smoking, or other processes aiming to improve its preservation and, lately, to enhance its flavor (Klurfeld 2015). Notably, the processing procedures include the use of additives, such as sodium, nitrites, and phosphates, which have been the focus of major interest for their potentially detrimental effects on health (Srouf & Touvier 2020). Overall, several meta-analyses have been conducted on this matter and various scientific organizations have provided rather discordant opinions: while in part agreeing that excessive consumption of processed meat would lead to

certain adverse health outcomes, the overall level of evidence, especially concerning red meat, has been questioned as in contrast with the high biological value of proteins and essential nutrients (Wyness 2016). The aim of this study is to provide a summary of existing meta-analyses exploring the relation between total, red, and processed meat consumption and various health outcomes; an attempt to measure the level of evidence for the retrieved results is also provided.

Methods

Study selection

A systematic search in Medline and Embase electronic databases of meta-analyses on meat consumption and various health outcomes was conducted until January 2017. The search strategy included: [(meat OR poultry OR chicken OR lamb OR pork) AND (meta-analysis OR meta-analyzed OR pooled analysis)]. Inclusion criteria were: (i) meta-analyses only including observational studies (prospective or case-control studies); (ii) quantitative analysis with the availability of measure of association [effect size as risk ratio (RR), odds ratio (OR), or hazard ratio (HR)]. Exclusion criteria were: (i) meta-analyses of RCTs with outcomes of intermediary biomarkers of disease (i.e., blood lipids, blood pressure, etc.) or (ii) intermediary clinical conditions (i.e., variation in body weight/BMI, etc.). Hand searching of reference lists was also undertaken. Any discrepancy on the inclusion/exclusion decision was solved through discussion.

Data extraction

From each meta-analysis included, the following information was extracted: name of the first author and year of publication, outcome, number of studies included in the meta-analysis, study design of included studies (i.e., case-control/cross-sectional and prospective), total

number of population, number of cases, type of exposure, measure of exposure [including highest versus lowest (reference) category of exposure or dose-response incremental servings per day (linear)], effect sizes.

Data evaluation and evidence synthesis

Where more than one meta-analysis was conducted on the same outcome, including the same study design, and the same population group, the concordance for the main outcome of interest, including direction and magnitude (overlapping confidence interval) of the association was evaluated. For further analyses, the most recent/exhaustive study was considered. The pooled analyses of the highest versus the lowest (reference) category of exposure and dose-response analyses were evaluated. Direction and magnitude of the association, heterogeneity (I^2) of results, and subgroup/stratified analyses for potential confounding factors were considered to have indication of the level of evidence. Criteria used for evidence categorization were modified from the Joint WHO/FAO Expert Consultation (Degrees of evidence by the Joint WHO/FAO Expert Consultation.

http://www.who.int/nutrition/topics/5_population_nutrient/en/#diet 5.1.2 Accessed

November 2015) (Table 1). Briefly, the relation between exposure and outcomes was categorized as following: suggestive/limited/contrasting evidence, when there was availability of solely meta-analyses of case-control studies, limited prospective cohort studies included in meta-analyses ($n < 3$), or evident contrasting results from meta-analyses with the same level of evidence; possible evidence, when there was availability of meta-analyses with lack of information on/significant heterogeneity ($I^2 > 50\%$) or identification of potential confounding factors (i.e., different findings in subgroups); probable association, when there was availability of meta-analyses of prospective cohort studies with no heterogeneity, no potential confounding factors identified, and eventual disagreement of results over time

reasonably explained (and evidence of dose-response relation further investigated); convincing association, when there was concordance between meta-analyses of RCTs and observational studies. Lack of fulfillment of the previous criteria was considered as insufficient evidence.

Results

Study selection

Out of 399 articles identified through the database search, 215 were excluded based on title and 97 after abstract evaluation (Figure 1). Twenty-eight studies were further excluded because meta-analyses of RCT ($n = 1$), narrative/systematic reviews without quantitative evaluation of the association between exposure and outcome ($n = 10$), pooled analysis of prospective cohort studies ($n = 3$), and investigating biomarkers ($n = 14$). Thus, a total number of 23 studies on total meat (Boyd et al. 1993; Sandhu et al. 2001; Norat et al. 2002; Faramawi et al. 2007; Aune et al. 2009; Kolahdooz et al. 2010; Micha et al. 2010; Wallin et al. 2011; Chan et al. 2011; Wang & Jiang 2012; Chen et al. 2013; Salehi et al. 2013; Zhu et al. 2013; Larsson & Orsini 2014; Zhu et al. 2014; Xu et al. 2014; Luo et al. 2014; Abete et al. 2014; Liu & Lin 2014; Alexander et al. 2015; Ge et al. 2015; Fang et al. 2015; Wu et al. 2016), 52 studies on red meat (Boyd et al. 1993; Sandhu et al. 2001; Norat et al. 2002; Larsson & Wolk 2006; Faramawi et al. 2007; Bandera et al. 2007; Alexander & Cushing 2009; Huxley et al. 2009; Aune et al. 2009; Kolahdooz et al. 2010; Micha et al. 2010; Smolińska & Paluszkiewicz 2010; Alexander, Mink, et al. 2010; Alexander, Morimoto, et al. 2010; Wallin et al. 2011; Chan et al. 2011; Alexander et al. 2011; Pan et al. 2011; Larsson & Wolk 2012; Paluszkiewicz et al. 2012; Wang & Jiang 2012; Kaluza et al. 2012; Chen et al. 2013; Huang et al. 2013; Xu et al. 2013; Choi et al. 2013; Aune et al. 2013; Salehi et al. 2013; Zhu et al. 2013; Qu et al. 2013; Fallahzadeh et al. 2014; Larsson & Orsini 2014; Zhu et

al. 2014; Luo et al. 2014; Song et al. 2014; Xue et al. 2014; Li et al. 2014; Abete et al. 2014; Guo et al. 2015; Alexander et al. 2015; Saneei et al. 2015; Yang et al. 2015; Fang et al. 2015; Bylsma & Alexander 2015; Caini et al. 2016; Wang et al. 2016; Solimini et al. 2016; Yang et al. 2016; Zhao et al. 2016; Li et al. 2016; Wu et al. 2016; Zhao et al. 2017) , and 47 studies on processed meat consumption (Sandhu et al. 2001; Norat et al. 2002; Huncharek et al. 2003; Larsson et al. 2006; Larsson & Wolk 2006; Faramawi et al. 2007; Alexander & Cushing 2009; Huxley et al. 2009; Aune et al. 2009; Kolahdooz et al. 2010; Micha et al. 2010; Alexander, Miller, et al. 2010; Alexander, Mink, et al. 2010; Alexander, Morimoto, et al. 2010; Wallin et al. 2011; Chan et al. 2011; Pan et al. 2011; Larsson & Wolk 2012; Wang & Jiang 2012; Kaluza et al. 2012; Chen et al. 2013; Huang et al. 2013; Xu et al. 2013; Choi et al. 2013; Aune et al. 2013; Salehi et al. 2013; Qu et al. 2013; Fallahzadeh et al. 2014; Larsson & Orsini 2014; Xu et al. 2014; Luo et al. 2014; Xue et al. 2014; Li et al. 2014; Abete et al. 2014; Wei et al. 2015; Guo et al. 2015; Saneei et al. 2015; Yang et al. 2015; Fang et al. 2015; Bylsma & Alexander 2015; Wang et al. 2016; Solimini et al. 2016; Yang et al. 2016; Zhao et al. 2016; Li et al. 2016; Wu et al. 2016; Zhao et al. 2017) and various health outcomes was selected for evaluation.

Characteristics of the studies on total meat consumption and health outcomes

The main characteristics of the studies exploring the risk associated to high vs. low total meat consumption for 10 unique outcomes from 8 non-overlapping meta-analyses and the risk associated to linear consumption for 9 unique outcomes from 7 non-overlapping meta-analyses are shown in Figure 2 and Supplementary Table 1, respectively. Higher intake of total meat was associated with an increased risk of colorectal and colon cancer, stroke, and all-cause mortality (Figure 2); a linear association was found for all outcomes investigated, including coronary heart disease (CHD), type-2 diabetes mellitus (T2DM), various cancers

(colorectal, colon, rectal, ovarian, and breast cancers), all-cause and cardiovascular disease (CVD) mortality (Supplementary Table 1). Two meta-analyses conducted on a limited number of prospective cohort studies (<3) and case-control studies also reported a potential association of high total meat intake and endometrial cancer (significant among case-control studies but not in the only prospective one) (Bandera et al. 2007), inflammatory bowel disease (Ge et al. 2015), and nasopharynx cancer (Xu et al. 2014). Most studies reported limited information on potential confounding factors: however, higher risk of colorectal cancer was significant only among European cohorts (Chan et al. 2011), while higher risk of all-cause mortality only among US ones (Larsson & Orsini 2014). Subgroup analyses by sex showed no significant association with risk of colorectal, colon, and rectal cancers (Chan et al. 2011).

Characteristics of the studies on red meat consumption and health outcomes

A summary of the main characteristics of the studies investigating the risk estimates for the highest vs. the lowest category of red meat consumption for 19 unique outcomes extracted from 16 non-overlapping meta-analyses in Figure 3 and the risk estimates of 14 outcomes from 12 non-overlapping meta-analyses for linear consumption of red meat are reported in Supplementary Table 2. The risk associated with high vs. low consumption of red meat included breast and lung cancers, stroke, CVD mortality, and T2DM, although with relatively high heterogeneity (>50%) between studies (Figure 3); a significant linear association was also reported for colorectal (and colon) cancer and adenomas, breast cancer, pancreatic cancer, stroke, and T2DM (Supplementary Table 2). Limited evidence was also found for higher intake of red meat and increased risk of endometrial cancer (Bandera et al. 2007), nasopharyngeal carcinoma (Li et al. 2016), and ischemic rather than hemorrhagic stroke (Yang et al. 2016). When controlling for potential confounding factors, certain variables

substantially modified the retrieved associations: for instance, increased risk of colorectal cancer was observed in European but not US or Asian cohorts (Chan et al. 2011), while other outcomes, such as all-cause mortality (Larsson & Orsini 2014), became significant when considering the same geographical area; in contrast, higher red meat intake was associated with increased risk of bladder and esophageal cancer in US cohorts but not in other regions (Zhu et al. 2014; Li et al. 2014). Moreover, increased risk of lung cancer and CVD mortality was significant only in the subgroup analysis restricted to women while all-cause mortality (Larsson & Orsini 2014) and pancreatic cancer (Zhao et al. 2017) risk was significant when restricted to men.

Characteristics of the studies on processed meat consumption and health outcomes

An overview of the main characteristics of the studies included and the risk estimates of various health outcomes for the highest versus the lowest category of processed meat consumption are reported for 19 unique outcomes extracted from 16 non-overlapping meta-analyses in Figure 4 and the risk estimates of 15 outcomes from 12 non-overlapping meta-analyses for linear consumption are reported in Supplementary Table 3. Higher intake of processed meat was associated with significantly increased risk of TD2M, all-cause mortality, various cancers (including prostate, gastric, colorectal, colon, rectal, renal, and ovarian cancers) and cardiovascular outcomes (including stroke and CVD mortality) (Figure 4). Similar findings with lower heterogeneity between studies (with exception of T2DM reporting >50% of heterogeneity) were reported when considering a linear association with increasing intake of processed meat (50 g/d or 1 serving/d increase) (Supplementary Table 3). Among meta-analyses with not sufficient evidence, an increased risk of colorectal adenomas (Aune et al. 2013), nasopharyngeal (Li et al. 2016), oral cavity, and oropharynx cancers (Xu et al. 2014), associated with higher intake of processed meat was also reported. Controlling

the results for potential confounding factors revealed that the increased risk of esophageal (Zhu et al. 2014), bladder (Li et al. 2014), pancreatic (Zhao et al. 2017) cancers and glioma (Wei et al. 2015) associated with higher processed meat consumption was significant only in US cohorts while breast cancer (Guo et al. 2015) only in European ones; some outcomes [such as CVD mortality (Abete et al. 2014) and colon cancer (Chan et al. 2011)] were significant in women only, while others (gastric (Zhu et al. 2013), lung (Xue et al. 2014), and pancreatic (Zhao et al. 2017) cancers) only in men. The analyses restricted to subgroups by sex resulted in no significant associations of processed meat intake and colorectal and rectal cancers in both men and women (Chan et al. 2011).

Summary of evidence

A detailed description of the evaluation of the evidence level is reported in Supplementary Table 4. Possible evidence of increased risk has been found for (i) adenoma of colorectum, lung cancer, CHD and stroke was found for higher intake of total meat; (ii) adenoma of colorectum, ovarian, prostate, renal, and stomach cancers, CHD and stroke for higher intake of red meat; and (iii) colon and bladder cancer for higher intake of processed meat (Table 1). Increased risk of other outcomes, including breast and colorectal cancers, T2DM, and mortality was deemed as limited mainly due to heterogeneity between results and potential otherwise inexplicable confounding factors (i.e., results were significant only in certain geographical regions or differed by sex). No probable nor convincing associations were found. Insufficient or no evidence of association has been reported for all other outcomes investigated (Table 1).

Discussion

In this umbrella review of meta-analyses investigating the relation between total, red, and processed meat consumption and various health outcomes, revealing rather limited evidence weakened by large heterogeneity across studies and geographical differences when considering grouping cohorts by regions. The reasons for such discrepancies between results are rather unclear: we may hypothesize that the differences by nationality may reflect different overall dietary patterns in which meat is consumed (for instance, hamburger with concomitant consumption of refined sugars from sugar-sweetened beverages and hydrogenated trans fats vs. lean meat with vegetables) as well as preferences for different cooking methods (for instance, a different impact of fried bacon, roasted beef, or long-cooked meat). However, a possible impact of high total, red, and processed meat on some cardiovascular and cancer outcomes have been observed. The rationale behind the detrimental effects of excess meat consumption in the diet on various health outcomes includes several mechanisms. Most studies stressed the concept that meat processing plays a central role in the detrimental effects on colorectal cancer risk: the conversion of nitrates and nitrites contained in processed meat into N-nitroso-compounds, which can act as multi-site carcinogens through the formation of covalent adducts with DNA bases (Alisson-Silva et al. 2016). Moreover, exposure to high temperature during cooking processes lead to the formation of known mutagens, such as heterocyclic amines (HCAs) and polycyclic aromatic hydrocarbons (PAHs), some of which have been associated with increased risk of colorectal adenomas (Hammerling et al. 2016). Interestingly, contrary to the common misconception that a diet rich in meat is mostly hazardous for cancer risk, in the present evaluation of the evidence we found evidence that both red and processed meat may comparably also be associated with a higher risk of CVD. Several specific mechanisms may explain this association: concerning processed meat, sodium (and specifically, NaCl) may play a detrimental role on endothelial health and raise blood pressure (Micha et al. 2013). However,

irrespectively of the processing level, the saturated fat content of meat (particularly characterized by long-chain saturated fatty acids) is under investigation as a potential trigger of low-grade sub-clinical inflammatory processes and potentially contribute to rise in post-prandial blood LDL (Astrup et al. 2020). Moreover, lipid oxidation occurring after the passage of meat through the various gastrointestinal compartments and further promoted by heme iron, high content in PUFA, and lack of antioxidants, may play a role in endothelial dysfunction and subsequent increased risk of CVD (Wu et al. 2019). Another hypothesized indirect cardiotoxic mechanism related to meat consumption may involve the transformation of L-carnitine into trimethylamine (TMA) by the gut microbiota and, subsequently, metabolized to trimethylamine-N-oxide (TMAO) in the liver: TMAO has been suggested to be involved in the atherogenesis processes, vascular inflammation, and increased platelet activity (Delgado et al. 2021).

The level of evidence reported in this umbrella review is lower than conclusions of expert scientific committees due to the stronger weight we imputed to heterogeneity of results among cohort studies (particularly evident when considering T2DM) and inexplicable differences across geographical regions (mostly concerning colorectal cancer and CVD outcomes). Scientific societies on cardiovascular health, such as the American College of Cardiology and American Heart Association, suggest limiting consumption of red and processed meat to ameliorate blood lipids and reduce the risk of CVD (Eckel et al. 2014). Experts from the American Diabetes Association and the expert panel of Diabetes UK also concluded in their recommendations to reduce meat-products for the prevention and management of T2DM (American Diabetes Association 2018; Dyson et al. 2018). Similar concerns have been reported by the World Cancer Research Fund/American Institute for Cancer Research and the Continuous Update Project, with attention to the consumption of red

and, especially processed meat and increased risk of colorectal cancer (Vieira et al. 2017). In all cases, we negatively weighted the various potential confounding factors contributing to the heterogeneity of the results and leading to an overall weakening of the level of evidence. The interpretation of such findings does not undermine the actual evidence of the retrieved associations between higher meat intake with increased risk of health conditions, but would rather suggest that other factors related to meat intake (i.e., unexplored confounding factors, effect modifiers, genetic differences among individuals, correlated variables, etc.) may play a synergistic role to define a pool or pattern of factors actually leading to the increased risk of disease. This hypothesis is supported by results from our previous study showing that red meat consumption in cohort studies globally correlates with several health risk factors, including higher BMI, higher prevalence of smokers, lower prevalence of high physical activity and university studies (Grosso et al. 2017). A recent ecological study showed that red meat had no significant impact on life expectancy in low-income or lower-middle-income countries (Ranabhat et al. 2020). A systematic review focused on studies conducted in Middle Eastern and North African countries showed an inverse association between (unprocessed) beef meat intake and colorectal cancer risk (Mint Sidi Deoula et al. 2020) and a multicenter case-control study involving the five largest university hospitals in Morocco investigated the differential association of traditional (decreased risk) vs. westernized (increased risk) processed meat products and colorectal cancer (S Deoula et al. 2020). Another individual study conducted on 21 low-, middle-, and high-income countries showed a rather null association between unprocessed red meat consumption and mortality or major CVD (Iqbal et al. 2021). Finally, concerning mechanistic studies, a case-control study conducted in non-smokers revealed the potential role of genes for S-glutathione transferase (GST), which are responsible for detoxification processes (including antioxidant activity) in the association between meat consumption and colorectal cancer risk (Klusek et al. 2019).

Albeit only preliminary, there is increasing awareness that other factors may play a role independently or together with higher meat consumption and human health. Further studies investigating intermediate outcomes and mechanistic processes in humans (i.e., gut microbiota modification, alteration of immunitary pathways, etc.) are needed to provide stronger evidence of the potential associations, ideally eliminating known confounding factors (such as, obesity, smoking habits, and scarce physical activity).

Another hypothesis that could explain the overall negative effects of meat consumption on human health, although with great heterogeneity between individuals, is the role of dietary fats on the immune system and inflammation (Ruiz-Núñez et al. 2016). There is consistent evidence from the scientific literature that intake of saturated fatty acids may trigger a number of pathways leading to local (gut level) and systemic low-grade inflammation, which in turn may affect endothelial health, cardiovascular risk, and ultimately the central nervous system through activation of the gut microbiota and pro-inflammatory processes (Ceppa et al. 2019; Salvucci 2019). Earlier meta-analyses of RCTs showed no significant differences in blood lipids levels comparing beef consumption to poultry and/or fish consumption (Maki et al. 2012) as well as consumption of ≥ 0.5 servings of total red meat/d on blood lipids, lipoproteins or blood pressure (O'Connor et al. 2017). However, a more recent meta-analysis showed that substituting red meat with high-quality plant protein sources, but not with fish or low-quality carbohydrates, may lead to favorable changes in blood lipids and lipoproteins (Guasch-Ferré et al. 2019). The findings from our recent work show substantial benefits on cardiometabolic outcomes of healthy alternative foods to meat, such as whole grains (Tieri et al. 2020), legumes and nuts (Martini et al. 2021). This may be explained by considering the different quality of fat contained in food groups, mostly omega-3 and omega-6 from vegetable sources, together with a higher fiber content

(Siri-Tarino et al. 2015; Fritsche 2015). It is under evaluation whether short-chain saturated fatty acids provided also from animal sources (i.e., dairy products and fish) may produce anti-inflammatory actions in the human body (Lordan et al. 2018). Whether confirmed by this pathway of action, the level of systemic low-grade inflammation may play a central role in nearly all non-communicable diseases, with a substantially stronger impact on the overall diet rather than individual components determining the final effect on the immune system (Stromsnes et al. 2021). Although evidence from observational studies supports the potential positive (Godos et al. 2020; Lee et al. 2020) or null effects (Marventano et al. 2020) on cardiovascular health, further research is needed to provide a stronger rationale for such potential benefits.

The present study has several limitations that should be discussed. First, the results are based on the original results of the meta-analyses of prospective studies included for evaluation, thus the same limitations applicable to observational studies should be considered (i.e., recall bias, reverse causation, under/over-reporting of meat intake, etc.). Second, the classification of the type of meat may not be univocal across individual prospective studies and across meta-analyses (i.e., some studies specified “red processed meat” while others included all types of processed meats). Similarly, total meat may include different types and proportions of meats (i.e., red, white, and processed) that may mislead the proper comparability of results.

In conclusion, excess meat consumption may be detrimental to health, potentially impacting both cardiometabolic and cancer risk. The heterogeneity of results does not allow to establish a clear ideal recommended frequency of consumption. Meat is a nutrient-dense food that significantly contributes to the nutritious dietary quality if compared to completely lacking in the average diet (McEvoy et al. 2012). Healthier dietary sources of protein with clear health

benefits are an advisable dietary choice, especially if compared to processed meats. However, the inclusion of moderate consumption of lean, unprocessed red meats in a balanced dietary pattern and a healthy lifestyle may not necessarily produce a harmful effect.

Acknowledgments

No funding.

Declaration of interests

The authors declare no conflicts of interest.

Figure legend

Figure 1. Flow chart of study selection.

Figure 2. Summary results from meta-analyses of prospective cohort studies on total meat consumption on various health outcomes included in umbrella review.

Figure 3. Summary results from meta-analyses of prospective cohort studies on red meat consumption on various health outcomes included in umbrella review.

Figure 4. Summary results from meta-analyses of prospective cohort studies on processed meat consumption on various health outcomes included in umbrella review.

Supplementary material legend

Supplementary Table 1. Summary results from meta-analyses investigating continuous linear exposure to total meat consumption and health outcomes.

Supplementary Table 2. Summary results from meta-analyses investigating continuous linear exposure to red meat consumption and health outcomes.

Supplementary Table 3. Summary results from meta-analyses investigating continuous linear exposure to processed meat consumption and health outcomes.

Supplementary Table 4. Variables investigated to address the strength of evidence from selected meta-analyses on total meat, red meat and processed meat consumption and health outcomes.

References

Abete I, Romaguera D, Vieira AR, Lopez de Munain A, Norat T. 2014. Association between total, processed, red and white meat consumption and all-cause, CVD and IHD mortality: a meta-analysis of cohort studies. *Br J Nutr.* 112(5):762–775.

Alexander DD, Cushing CA. 2009. Quantitative assessment of red meat or processed meat consumption and kidney cancer. *Cancer Detect Prev.* 32(5–6):340–351.

Alexander DD, Miller AJ, Cushing CA, Lowe KA. 2010. Processed meat and colorectal cancer: a quantitative review of prospective epidemiologic studies. *Eur J Cancer Prev.* 19(5):328–341.

Alexander DD, Mink PJ, Cushing CA, Scurman B. 2010. A review and meta-analysis of prospective studies of red and processed meat intake and prostate cancer. *Nutr J.* 9:50.

Alexander DD, Morimoto LM, Mink PJ, Cushing CA. 2010. A review and meta-analysis of red and processed meat consumption and breast cancer. *Nutr Res Rev.* 23(2):349–365.

Alexander DD, Weed DL, Cushing CA, Lowe KA. 2011. Meta-analysis of prospective studies of red meat consumption and colorectal cancer. *Eur J Cancer Prev.* 20(4):293–307.

Alexander DD, Weed DL, Miller PE, Mohamed MA. 2015. Red meat and colorectal cancer: A quantitative update on the state of the epidemiologic science. *J Am Coll Nutr.* 34(6):521–543.

Alisson-Silva F, Kawanishi K, Varki A. 2016. Human risk of diseases associated with red meat intake: Analysis of current theories and proposed role for metabolic incorporation of a non-human sialic acid. *Mol Aspects Med.* 51:16–30.

American Diabetes Association. 2018. Standards of Medical Care in Diabetes-2018 Abridged for Primary Care Providers. *Clin Diabetes.* 36(1):14–37.

Astrup A, Magkos F, Bier DM, Brenna JT, de Oliveira Otto MC, Hill JO, King JC, Mente A, Ordovas JM, Volek JS, et al. 2020. Saturated Fats and Health: A Reassessment and Proposal for Food-Based Recommendations: JACC State-of-the-Art Review. *J Am Coll Cardiol.* 76(7):844–857.

Aune D, Chan DSM, Vieira AR, Navarro Rosenblatt DA, Vieira R, Greenwood DC, Kampman E, Norat T. 2013. Red and processed meat intake and risk of colorectal adenomas:

a systematic review and meta-analysis of epidemiological studies. *Cancer Causes Control*. 24(4):611–627.

Aune D, Ursin G, Veierød MB. 2009. Meat consumption and the risk of type 2 diabetes: a systematic review and meta-analysis of cohort studies. *Diabetologia*. 52(11):2277–2287.

Bandera EV, Kushi LH, Moore DF, Gifkins DM, McCullough ML. 2007. Consumption of animal foods and endometrial cancer risk: a systematic literature review and meta-analysis. *Cancer Causes Control*. 18(9):967–988.

Boyd NF, Martin LJ, Noffel M, Lockwood GA, Trichler DL. 1993. A meta-analysis of studies of dietary fat and breast cancer risk. *Br J Cancer*. 68(3):627–636.

Bylsma LC, Alexander DD. 2015. A review and meta-analysis of prospective studies of red and processed meat, meat cooking methods, heme iron, heterocyclic amines and prostate cancer. *Nutr J*. 14:125.

Caini S, Masala G, Gnagnarella P, Ermini I, Russell-Edu W, Palli D, Gandini S. 2016. Food of animal origin and risk of non-Hodgkin lymphoma and multiple myeloma: A review of the literature and meta-analysis. *Crit Rev Oncol Hematol*. 100:16–24.

Carmody RN, Wrangham RW. 2009. The energetic significance of cooking. *J Hum Evol*. 57(4):379–391.

Ceppa F, Mancini A, Tuohy K. 2019. Current evidence linking diet to gut microbiota and brain development and function. *Int J Food Sci Nutr*. 70(1):1–19.

Chan DSM, Lau R, Aune D, Vieira R, Greenwood DC, Kampman E, Norat T. 2011. Red and processed meat and colorectal cancer incidence: meta-analysis of prospective studies. *PLoS ONE*. 6(6):e20456.

Chen GC, Lv DB, Pang Z, Liu QF. 2013. Red and processed meat consumption and risk of stroke: a meta-analysis of prospective cohort studies. *Eur J Clin Nutr*. 67(1):91–95.

Choi Y, Song S, Song Y, Lee JE. 2013. Consumption of red and processed meat and esophageal cancer risk: meta-analysis. *World J Gastroenterol*. 19(7):1020–1029.

Cocking C, Walton J, Kehoe L, Cashman KD, Flynn A. 2020. The role of meat in the European diet: current state of knowledge on dietary recommendations, intakes and contribution to energy and nutrient intakes and status. *Nutr Res Rev*. 33(2):181–189.

Delgado J, Ansorena D, Van Hecke T, Astiasarán I, De Smet S, Estévez M. 2021. Meat lipids, NaCl and carnitine: Do they unveil the conundrum of the association between red and processed meat intake and cardiovascular diseases? _Invited Review. *Meat Sci*. 171:108278.

Dyson PA, Twenefour D, Breen C, Duncan A, Elvin E, Goff L, Hill A, Kalsi P, Marsland N, McArdle P, et al. 2018. Diabetes UK evidence-based nutrition guidelines for the prevention and management of diabetes. *Diabet Med*. 35(5):541–547.

Eckel RH, Jakicic JM, Ard JD, de Jesus JM, Houston Miller N, Hubbard VS, Lee I-M, Lichtenstein AH, Loria CM, Millen BE, et al. 2014. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of

Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 63(25 Pt B):2960–2984.

Fallahzadeh H, Cheraghi M, Amoori N, Alaf M. 2014. Red meat intake and risk of non-Hodgkin lymphoma: a meta-analysis. *Asian Pac J Cancer Prev*. 15(23):10421–10425.

Fang X, Wei J, He X, An P, Wang H, Jiang L, Shao D, Liang H, Li Y, Wang F, Min J. 2015. Landscape of dietary factors associated with risk of gastric cancer: A systematic review and dose-response meta-analysis of prospective cohort studies. *Eur J Cancer*. 51(18):2820–2832.

Faramawi MF, Johnson E, Fry MW, Sall M, Zhou Y. 2007. Consumption of different types of meat and the risk of renal cancer: meta-analysis of case-control studies. *Cancer Causes Control*. 18(2):125–133.

Fritsche KL. 2015. The science of fatty acids and inflammation. *Adv Nutr*. 6(3):293S–301S.

Ge J, Han TJ, Liu J, Li JS, Zhang XH, Wang Y, Li QY, Zhu Q, Yang CM. 2015. Meat intake and risk of inflammatory bowel disease: A meta-analysis. *Turk J Gastroenterol*. 26(6):492–497.

Godos J, Tieri M, Ghelfi F, Titta L, Marventano S, Lafranconi A, Gambera A, Alonzo E, Sciacca S, Buscemi S, et al. 2020. Dairy foods and health: an umbrella review of observational studies. *Int J Food Sci Nutr*. 71(2):138–151.

Grosso G, Micek A, Godos J, Pajak A, Sciacca S, Galvano F, Boffetta P. 2017. Health risk factors associated with meat, fruit and vegetable consumption in cohort studies: A comprehensive meta-analysis. *PLoS ONE*. 12(8):e0183787.

Guasch-Ferré M, Satija A, Blondin SA, Janiszewski M, Emlen E, O'Connor LE, Campbell WW, Hu FB, Willett WC, Stampfer MJ. 2019. Meta-Analysis of Randomized Controlled Trials of Red Meat Consumption in Comparison With Various Comparison Diets on Cardiovascular Risk Factors. *Circulation*. 139(15):1828–1845.

Guo J, Wei W, Zhan L. 2015. Red and processed meat intake and risk of breast cancer: a meta-analysis of prospective studies. *Breast Cancer Res Treat*. 151(1):191–198.

Hammerling U, Bergman Laurila J, Grafström R, Ilbäck N-G. 2016. Consumption of red/processed meat and colorectal carcinoma: possible mechanisms underlying the significant association. *Crit Rev Food Sci Nutr*. 56(4):614–634.

Henchion M, Moloney AP, Hyland J, Zimmermann J, McCarthy S. 2021. Review: Trends for meat, milk and egg consumption for the next decades and the role played by livestock systems in the global production of proteins. *Animal*. 15 Suppl 1:100287.

Huang W, Han Y, Xu J, Zhu W, Li Z. 2013. Red and processed meat intake and risk of esophageal adenocarcinoma: a meta-analysis of observational studies. *Cancer Causes Control*. 24(1):193–201.

Huncharek M, Kupelnick B, Wheeler L. 2003. Dietary cured meat and the risk of adult glioma: a meta-analysis of nine observational studies. *J Environ Pathol Toxicol Oncol*. 22(2):129–137.

- Huxley RR, Ansary-Moghaddam A, Clifton P, Czernichow S, Parr CL, Woodward M. 2009. The impact of dietary and lifestyle risk factors on risk of colorectal cancer: a quantitative overview of the epidemiological evidence. *Int J Cancer*. 125(1):171–180.
- Iqbal R, Dehghan M, Mente A, Rangarajan S, Wielgosz A, Avezum A, Seron P, AlHabib KF, Lopez-Jaramillo P, Swaminathan S, et al. 2021. Associations of unprocessed and processed meat intake with mortality and cardiovascular disease in 21 countries [Prospective Urban Rural Epidemiology (PURE) Study]: a prospective cohort study. *Am J Clin Nutr*. 114(3):1049–1058.
- Kaluza J, Wolk A, Larsson SC. 2012. Red meat consumption and risk of stroke: a meta-analysis of prospective studies. *Stroke*. 43(10):2556–2560.
- Klurfeld DM. 2015. Research gaps in evaluating the relationship of meat and health. *Meat Sci*. 109:86–95.
- Klusek Justyna, Nasierowska-Guttmejer A, Kowalik A, Wawrzycka I, Chrapek M, Lewitowicz P, Radowicz-Chil A, Klusek Jolanta, Głuszek S. 2019. The Influence of Red Meat on Colorectal Cancer Occurrence Is Dependent on the Genetic Polymorphisms of S-Glutathione Transferase Genes. *Nutrients*. 11(7).
- Kolahdooz F, van der Pols JC, Bain CJ, Marks GC, Hughes MC, Whiteman DC, Webb PM, Australian Cancer Study (Ovarian Cancer) and the Australian Ovarian Cancer Study Group. 2010. Meat, fish, and ovarian cancer risk: Results from 2 Australian case-control studies, a systematic review, and meta-analysis. *Am J Clin Nutr*. 91(6):1752–1763.
- Larsen CS. 2003. Animal source foods and human health during evolution. *J Nutr*. 133(11 Suppl 2):3893S–3897S.
- Larsson SC, Orsini N, Wolk A. 2006. Processed meat consumption and stomach cancer risk: a meta-analysis. *J Natl Cancer Inst*. 98(15):1078–1087.
- Larsson SC, Orsini N. 2014. Red meat and processed meat consumption and all-cause mortality: a meta-analysis. *Am J Epidemiol*. 179(3):282–289.
- Larsson SC, Wolk A. 2006. Meat consumption and risk of colorectal cancer: a meta-analysis of prospective studies. *Int J Cancer*. 119(11):2657–2664.
- Larsson SC, Wolk A. 2012. Red and processed meat consumption and risk of pancreatic cancer: meta-analysis of prospective studies. *Br J Cancer*. 106(3):603–607.
- Lee KH, Seong HJ, Kim G, Jeong GH, Kim JY, Park H, Jung E, Kronbichler A, Eisenhut M, Stubbs B, et al. 2020. Consumption of Fish and ω -3 Fatty Acids and Cancer Risk: An Umbrella Review of Meta-Analyses of Observational Studies. *Adv Nutr*. 11(5):1134–1149.
- Liu Z-T, Lin A-H. 2014. Dietary factors and thyroid cancer risk: a meta-analysis of observational studies. *Nutr Cancer*. 66(7):1165–1178.
- Li F, An S, Hou L, Chen P, Lei C, Tan W. 2014. Red and processed meat intake and risk of bladder cancer: a meta-analysis. *Int J Clin Exp Med*. 7(8):2100–2110.
- Li F, Duan F, Zhao X, Song C, Cui S, Dai L. 2016. Red Meat and Processed Meat

Consumption and Nasopharyngeal Carcinoma Risk: A Dose-response Meta-analysis of Observational Studies. *Nutr Cancer*. 68(6):1034–1043.

Lordan R, Tsoupras A, Mitra B, Zabetakis I. 2018. Dairy fats and cardiovascular disease: do we really need to be concerned? *Foods*. 7(3).

Luo J, Yang Y, Liu J, Lu K, Tang Z, Liu P, Liu L, Zhu Y. 2014. Systematic review with meta-analysis: meat consumption and the risk of hepatocellular carcinoma. *Aliment Pharmacol Ther*. 39(9):913–922.

Maki KC, Van Elswyk ME, Alexander DD, Rains TM, Sohn EL, McNeill S. 2012. A meta-analysis of randomized controlled trials that compare the lipid effects of beef versus poultry and/or fish consumption. *J Clin Lipidol*. 6(4):352–361.

Mann NJ. 2018. A brief history of meat in the human diet and current health implications. *Meat Sci*. 144:169–179.

Martini D, Godos J, Marventano S, Tieri M, Ghelfi F, Titta L, Lafranconi A, Trigueiro H, Gambera A, Alonzo E, et al. 2021. Nut and legume consumption and human health: an umbrella review of observational studies. *Int J Food Sci Nutr*. 72(7):871–878.

Marventano S, Godos J, Tieri M, Ghelfi F, Titta L, Lafranconi A, Gambera A, Alonzo E, Sciacca S, Buscemi S, et al. 2020. Egg consumption and human health: an umbrella review of observational studies. *Int J Food Sci Nutr*. 71(3):325–331.

McEvoy CT, Temple N, Woodside JV. 2012. Vegetarian diets, low-meat diets and health: a review. *Public Health Nutr*. 15(12):2287–2294.

Micha R, Michas G, Lajous M, Mozaffarian D. 2013. Processing of meats and cardiovascular risk: time to focus on preservatives. *BMC Med*. 11:136.

Micha R, Wallace SK, Mozaffarian D. 2010. Red and processed meat consumption and risk of incident coronary heart disease, stroke, and diabetes mellitus: a systematic review and meta-analysis. *Circulation*. 121(21):2271–2283.

Milton K. 2003. The critical role played by animal source foods in human (*Homo*) evolution. *J Nutr*. 133(11 Suppl 2):3886S–3892S.

Mint Sidi Deoula M, El Kinany K, Hatime Z, Boudouaya HA, El Rhazi K. 2020. Meat and colorectal cancer in Middle Eastern and North African countries: update of literature review. *Public Health Rev*. 41:7.

Norat T, Lukanova A, Ferrari P, Riboli E. 2002. Meat consumption and colorectal cancer risk: dose-response meta-analysis of epidemiological studies. *Int J Cancer*. 98(2):241–256.

O'Connor LE, Kim JE, Campbell WW. 2017. Total red meat intake of ≥ 0.5 servings/d does not negatively influence cardiovascular disease risk factors: a systemically searched meta-analysis of randomized controlled trials. *Am J Clin Nutr*. 105(1):57–69.

O'Keefe JH, Cordain L. 2004. Cardiovascular disease resulting from a diet and lifestyle at odds with our Paleolithic genome: how to become a 21st-century hunter-gatherer. *Mayo Clin Proc*. 79(1):101–108.

- Paluszkiewicz P, Smolińska K, Dębińska I, Turski WA. 2012. Main dietary compounds and pancreatic cancer risk. The quantitative analysis of case-control and cohort studies. *Cancer Epidemiol.* 36(1):60–67.
- Pan A, Sun Q, Bernstein AM, Schulze MB, Manson JE, Willett WC, Hu FB. 2011. Red meat consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis. *Am J Clin Nutr.* 94(4):1088–1096.
- Qu X, Ben Q, Jiang Y. 2013. Consumption of red and processed meat and risk for esophageal squamous cell carcinoma based on a meta-analysis. *Ann Epidemiol.* 23(12):762-770.e1.
- Ranabhat CL, Park M-B, Kim C-B. 2020. Influence of Alcohol and Red Meat Consumption on Life Expectancy: Results of 164 Countries from 1992 to 2013. *Nutrients.* 12(2).
- Ruiz-Núñez B, Dijck-Brouwer DAJ, Muskiet FAJ. 2016. The relation of saturated fatty acids with low-grade inflammation and cardiovascular disease. *J Nutr Biochem.* 36:1–20.
- Salehi M, Moradi-Lakeh M, Salehi MH, Nojomi M, Kollahdooz F. 2013. Meat, fish, and esophageal cancer risk: a systematic review and dose-response meta-analysis. *Nutr Rev.* 71(5):257–267.
- Salvucci E. 2019. The human-microbiome superorganism and its modulation to restore health. *Int J Food Sci Nutr.* 70(7):781–795.
- Sandhu MS, White IR, McPherson K. 2001. Systematic review of the prospective cohort studies on meat consumption and colorectal cancer risk: a meta-analytical approach. *Cancer Epidemiol Biomarkers Prev.* 10(5):439–446.
- Saneei P, Willett W, Esmailzadeh A. 2015. Red and processed meat consumption and risk of glioma in adults: A systematic review and meta-analysis of observational studies. *J Res Med Sci.* 20(6):602–612.
- Siri-Tarino PW, Chiu S, Bergeron N, Krauss RM. 2015. Saturated fats versus polyunsaturated fats versus carbohydrates for cardiovascular disease prevention and treatment. *Annu Rev Nutr.* 35:517–543.
- S Deoula M, El Kinany K, Huybrechts I, Gunter MJ, Hatime Z, Boudouaya HA, Benslimane A, Nejari C, El Abkari M, Badre W, et al. 2020. Consumption of meat, traditional and modern processed meat and colorectal cancer risk among the Moroccan population: A large-scale case-control study. *Int J Cancer.* 146(5):1333–1345.
- Smolińska K, Paluszkiewicz P. 2010. Risk of colorectal cancer in relation to frequency and total amount of red meat consumption. Systematic review and meta-analysis. *Arch Med Sci.* 6(4):605–610.
- Solimini AG, Lombardi AM, Palazzo C, De Giusti M. 2016. Meat intake and non-Hodgkin lymphoma: a meta-analysis of observational studies. *Cancer Causes Control.* 27(5):595–606.
- Song P, Lu M, Yin Q, Wu L, Zhang D, Fu B, Wang B, Zhao Q. 2014. Red meat consumption and stomach cancer risk: a meta-analysis. *J Cancer Res Clin Oncol.* 140(6):979–992.
- Srour B, Touvier M. 2020. Processed and ultra-processed foods: coming to a health problem?

Int J Food Sci Nutr. 71(6):653–655.

Stromsnes K, Correas AG, Lehmann J, Gambini J, Olaso-Gonzalez G. 2021. Anti-Inflammatory Properties of Diet: Role in Healthy Aging. *Biomedicines*. 9(8).

Tieri M, Ghelfi F, Vitale M, Vetrani C, Marventano S, Lafranconi A, Godos J, Titta L, Gambera A, Alonzo E, et al. 2020. Whole grain consumption and human health: an umbrella review of observational studies. *Int J Food Sci Nutr*. 71(6):668–677.

Vieira AR, Abar L, Chan DSM, Vingeliene S, Polemiti E, Stevens C, Greenwood D, Norat T. 2017. Foods and beverages and colorectal cancer risk: a systematic review and meta-analysis of cohort studies, an update of the evidence of the WCRF-AICR Continuous Update Project. *Ann Oncol*. 28(8):1788–1802.

Wallin A, Orsini N, Wolk A. 2011. Red and processed meat consumption and risk of ovarian cancer: a dose-response meta-analysis of prospective studies. *Br J Cancer*. 104(7):1196–1201.

Wang C, Jiang H. 2012. Meat intake and risk of bladder cancer: a meta-analysis. *Med Oncol*. 29(2):848–855.

Wang X, Lin X, Ouyang YY, Liu J, Zhao G, Pan A, Hu FB. 2016. Red and processed meat consumption and mortality: dose-response meta-analysis of prospective cohort studies. *Public Health Nutr*. 19(5):893–905.

Wei Y, Zou D, Cao D, Xie P. 2015. Association between processed meat and red meat consumption and risk for glioma: a meta-analysis from 14 articles. *Nutrition*. 31(1):45–50.

Wu J, Zeng R, Huang J, Li X, Zhang J, Ho JC-M, Zheng Y. 2016. Dietary Protein Sources and Incidence of Breast Cancer: A Dose-Response Meta-Analysis of Prospective Studies. *Nutrients*. 8(11).

Wu JHY, Micha R, Mozaffarian D. 2019. Dietary fats and cardiometabolic disease: mechanisms and effects on risk factors and outcomes. *Nat Rev Cardiol*. 16(10):581–601.

Wyness L. 2016. The role of red meat in the diet: nutrition and health benefits. *Proc Nutr Soc*. 75(3):227–232.

Xue X-J, Gao Q, Qiao J-H, Zhang J, Xu C-P, Liu J. 2014. Red and processed meat consumption and the risk of lung cancer: a dose-response meta-analysis of 33 published studies. *Int J Clin Exp Med*. 7(6):1542–1553.

Xu J, Yang X, Wu Y, Li X, Bai B. 2014. Meat consumption and risk of oral cavity and oropharynx cancer: a meta-analysis of observational studies. *PLoS ONE*. 9(4):e95048.

Xu X, Yu E, Gao X, Song N, Liu L, Wei X, Zhang W, Fu C. 2013. Red and processed meat intake and risk of colorectal adenomas: a meta-analysis of observational studies. *Int J Cancer*. 132(2):437–448.

Yang C, Pan L, Sun C, Xi Y, Wang L, Li D. 2016. Red Meat Consumption and the Risk of Stroke: A Dose-Response Meta-analysis of Prospective Cohort Studies. *J Stroke Cerebrovasc Dis*. 25(5):1177–1186.

Yang L, Dong J, Jiang S, Shi W, Xu X, Huang H, You X, Liu H. 2015. Red and Processed Meat Consumption Increases Risk for Non-Hodgkin Lymphoma: A PRISMA-Compliant Meta-Analysis of Observational Studies. *Medicine (Baltimore)*. 94(45):e1729.

Zhao Z, Pu Z, Yin Z, Yu P, Hao Y, Wang Q, Guo M, Zhao Q. 2016. Dietary fruit, vegetable, fat, and red and processed meat intakes and Barrett's esophagus risk: a systematic review and meta-analysis. *Sci Rep*. 6:27334.

Zhao Z, Yin Z, Pu Z, Zhao Q. 2017. Association Between Consumption of Red and Processed Meat and Pancreatic Cancer Risk: A Systematic Review and Meta-analysis. *Clin Gastroenterol Hepatol*. 15(4):486-493.e10.

Zhu H, Yang X, Zhang C, Zhu C, Tao G, Zhao L, Tang S, Shu Z, Cai J, Dai S, et al. 2013. Red and processed meat intake is associated with higher gastric cancer risk: a meta-analysis of epidemiological observational studies. *PLoS ONE*. 8(8):e70955.

Zhu H-C, Yang X, Xu L-P, Zhao L-J, Tao G-Z, Zhang C, Qin Q, Cai J, Ma J-X, Mao W-D, et al. 2014. Meat consumption is associated with esophageal cancer risk in a meat- and cancer-histological-type dependent manner. *Dig Dis Sci*. 59(3):664–673.

Table 1. Level of evidence for the association between meat consumption and health outcomes.

Level of evidence*	Criteria§	Total meat	Red meat	Processed meat
Convincing	Meta-analyses of prospective cohort studies with evidence of dose-response relation, no heterogeneity, no potential confounding factors identified, and eventual disagreement of results over time reasonably explained [otherwise declassified as possible].	None.	None.	None.
Probable	Meta-analyses of prospective cohort studies with no heterogeneity, no potential confounding factors identified, and eventual disagreement of results over time reasonably explained [otherwise declassified as possible].	None.	None.	None.
Possible	Meta-analysis of prospective cohort studies with no heterogeneity and lack of information on potential confounding factors.	Association with increased risk of adenoma (colorectum), cancer (lung), CHD (any), stroke (total).	Association with increased risk of adenoma (colorectum), cancer (ovarian, prostate, renal, stomach), CHD (any), stroke (total).	Association with increased risk of cancer (colon, bladder).
Limited	Meta-analysis of prospective cohort studies with presence of significant heterogeneity ($I^2 > 50\%$) or identification of potential confounding factors (i.e., different findings in subgroups).	Association with increased risk of cancer (breast), mortality (CVD), T2DM.	Association with increased risk of cancer (breast, colon#, colorectum#, rectum#), mortality (all-cause, CVD), T2DM.	Association with increased risk of cancer (colorectum#), CHD (any), mortality (all-cause, cancer), stroke (total).
Insufficient	Meta-analysis of case-control studies, limited prospective cohort studies included in meta-analyses (n < 3), or evident contrasting results from meta-analyses with the same level of evidence.	Association with increased odds of cancer (nasopharynx), inflammatory bowel disease, and stroke (ischemic).	Association with increased odds of cancer (naso-opharynx).	Association with increased odds of inflammatory bowel disease.

No evidence	Meta-analyses of prospective cohort studies with evidence of dose-response relation, no heterogeneity, no potential confounding factors identified, and eventual disagreement of results over time reasonably explained [otherwise declassified as possible].	No association with risk of Barret's esophagus, cancer (bladder, colon, colorectum, esophagus, endometrial, liver, NHL, ovarian, pancreas, prostate, renal, rectum, stomach), mortality (all-cause, cancer), stroke (hemorrhagic).	No association with risk of Barret's esophagus, cancer (bladder, esophagus, glioma, liver, lung, NHL, pancreas), mortality (CHD), stroke (ischemic, hemorrhagic).	No association with risk of cancer (breast, esophagus, rectum, stomach), mortality (CHD), T2DM.
*all the associations should be biologically plausible; potential confounding factors should be taken into account. § modified from the Joint WHO/FAO Expert Consultation # presence of potential confounding factors Abbreviations: CHD, coronary heart disease; CVD, cardiovascular disease; NHL, non-Hodgkin lymphoma; T2DM, type-2 diabetes mellitus.				

Figure 1




Figure 2




Figure 3




Figure 4

