

The stunting effect of an oxylipins-containing macroalgae extract on sea urchin reproduction and neuroblastoma cells viability

Abstract

Different bioactive molecules extracted from macroalgae, including oxylipins, showed interesting potentials in different applications, from healthcare to biomaterial manufacturing and environmental remediation. Thus far, no studies reported the effects of oxylipins-containing macroalgae extracts on embryo development of marine invertebrates and on neuroblastoma cancer cells. Here, the effects of an oxylipins-containing extract from *Ericaria brachycarpa*, a canopy-forming brown algae, were investigated on the development of *Arbacia lixula* sea urchin embryos and on SH-SY5Y neuroblastoma cells viability. Embryos and cells were exposed to concentrations covering a full 0-100 % dose-response curve, with doses ranging from 0 to 40 µg/mL for embryos and from 0 to 200 µg/mL for cells. These natural marine toxins caused a dose-dependent decrease of normal embryos development and of neuroblastoma cells viability. Toxicity was higher for exposures starting from the gastrula embryonal stage if compared to the zygote and pluteus stages, with an EC₅₀ significantly lower by 33 and 68 %, respectively. Embryos exposed to low doses showed a general delay in development with a decrease in the ability to calcify, while higher doses caused 100% block of embryo growth. Exposure of SH-SY5Y neuroblastoma cells to 40 µg/mL for 72 h caused 78 % mortality, while no effect was observed on their neuronal-like cells derivatives, suggesting a selective targeting of proliferating cells. Western Blot experiments on both model systems displayed the modulation of different molecular markers (HSP60, HSP90, LC3, p62, CHOP and cleaved caspase-7), showing altered stress response and enhanced autophagy and apoptosis, confirmed by increased fragmented DNA in apoptotic nuclei. Our study gives new insights into the molecular strategies that marine invertebrates use when responding to their environmental natural toxins and suggests the *E.*

24 *brachycarpa*'s extract as a potential source for the development of innovative, environmentally
25 friendly products with larvicide and antineoplastic activity.

26 **Keywords:** marine drugs; ecotoxicology; oxylipins; developmental abnormality; SH-SY5Y;
27 apoptosis.

28 **Abbreviations:** MFSW, millipore filtered seawater; EBE, *Ericaria brachycarpa* extract; HEPes,
29 hydroxyacids; hpf, hours post fertilization; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-
30 diphenyltetrazolium bromide; HSPs, heat shock proteins; TUNEL, TdT-mediated dUTP nick-end
31 labeling; WB, western blot; EC₅₀, half maximal effective concentration; NOEC, no observed effect
32 concentration.

33

34 **1. Introduction**

35 Extracts and molecules obtained from several marine species have shown anticancer, anti-angiogenic,
36 anti-inflammatory and wound healing/skin regeneration abilities, finding applications in healthcare,
37 biomaterial manufacturing and environmental remediation (Conte et al., 2020; Pereira and Cotas,
38 2022; Librizzi et al., 2024). Among marine organisms, macroalgae are sessile organisms that need to
39 constantly adapt to either the abiotic and biotic components of the marine ecosystem and developed
40 competitive and complex adaptations to safeguard their species, including the production of a wealth
41 of bioactive molecules *via* their secondary metabolism as a chemical strategy against environmental
42 stressors and against their predators (Budzałek et al., 2021; Seth et al., 2024). Driven by the increased
43 global interest in natural products, macroalgae are currently being investigated as a primary source
44 of new classes of molecules with greater bioavailability and efficacy and less toxicity, for
45 nutraceuticals and pharmaceutical applications (Adarshan et al., 2023). Several algal-extracted
46 compounds have been characterized and their anticarcinogenic activity was demonstrated *in vivo* and
47 *in vitro* (Thaman et al., 2023). Moreover, extracts and molecules obtained from macroalgae have been

48 tested on invertebrates, showing a wide range of activities, including feeding-deterrent abilities
49 against sea urchins grazing (Budzalek et al., 2021).

50 Among secondary metabolites, oxylipins are generated from polyunsaturated fatty acids via the
51 lipoxygenase pathway, accumulating as a reaction to abiotic and biotic stressors (Barbosa et al.,
52 2016). Diatoms-derived oxylipins are produced after cell damage and during grazing by predators
53 (Ruocco et al., 2019), and their release in the marine environment can subsequently induce
54 reproductive impairment and teratogenic effects on their predators, including copepods and sea
55 urchins (Ruocco et al., 2019; Lauritano and Ianora 2020). Exposure of *Paracentrotus lividus* embryos
56 to diatoms-derived polyunsaturated aldehydes in the μM range showed a dose-dependent block of the
57 first cleavage and an increase in the percentage of malformed and delayed embryos, together with a
58 large amount of fragmented DNA (Romano et al., 2010). Other diatoms-derived oxylipins,
59 hydroxyacids (HEPEs), had low toxicity to adult copepods but highly impaired their gametes and
60 nauplii (Fontana et al., 2007) and caused a dose-dependent effect with the induction of skeletal
61 malformations in *P. lividus* embryos (Varrella et al., 2016). Moreover, diatom-derived oxylipins can
62 induce cell death in different human cancer cell lines in the μM to low mM range (Miralto et al. 1999;
63 Sansone et al. 2014; Ávila-Román et al. 2016). In macroalgae, oxylipins operate in systemic defense
64 mechanisms and are induced responding to different stresses and against grazers and pathogens
65 (Jacquemoud and Pohnert, 2015; Barbosa et al., 2016). While many studies are available on the effect
66 of diatoms-derived oxylipins on marine invertebrates' reproduction and on human cancer cell
67 viability, the biological effect of macroalgae-derived oxylipins is still poorly elucidated.

68 Among macroalgae, *Ericaria* spp. are canopy-forming brown algae belonging to the Sargassaceae
69 family that live in the Mediterranean rocky coastline. One of these species, *Ericaria brachycarpa* ((J.
70 Agardh) Molinari & Guiry, 2020), contribute to form the habitat for the sympatric sea urchins species
71 *P. lividus* and *Arbacia lixula* (Echinodermata: Echinoidea; Bonaviri et al., 2011). The grazing activity
72 of *A. lixula* exerts a primary control on the composition and abundance of canopy-forming macroalgal

assemblages, thanks to its strong Aristotle's lantern that allows the scraping of the substrate (Privitera et al., 2008; Bonaviri et al., 2011), causing the formation of barren grounds (Bonaviri et al., 2011; Lawrence, 2013). Numerous molecules have been identified from various species of the *Ericaria* genus, including terpenoids, steroids, phlorotannins and phenolic compounds, some of which showing antimicrobial, cytotoxic and antitumoral features (Amico 1995; de Sousa et al., 2017).

Here, for the first time, we identified seven oxylipins in the extract of a species of the *Ericaria* genus, (*E. brachycarpa*) and tested the effects of this extract on *A. lixula* sea urchin embryos and on the human neuroblastoma SH-SY5Y cell line and their neuronal-like cells derivatives. In order to understand the molecular mechanisms activated in exposed *A. lixula* embryos and SH-SY5Y cells after exposure, we analyzed the expression levels of different proteins involved in the stress response (HSPs) and of molecular markers of the autophagic (LC3 and p62/SQSTM1) and apoptotic (cleaved caspase-7 and CHOP-10/GADD153) pathways. Finally, we evaluated the presence of fragmented DNA through TUNEL staining. To our knowledge, this is the first time that an oxylipin-containing extract from a macroalgae shows growth-restraining activity against embryos of a marine Invertebrate and against neuroblastoma cancer cells.

2. Materials and methods

2.1 *E. brachycarpa* sampling, extraction and HPLC/MS analysis

The sporophytes of *E. brachycarpa* and adult *A. lixula* sea urchins were sampled at the littoral of Palermo, in the Northern Sicilian coast, where they naturally coexist. The sporophytes were washed and rinsed to remove debris and epiphytes, then dried at 40°C in an oven for 48 h. The dried thalli were subsequently finely ground to obtain a powder using a stainless-steel grinder. Metabolites extraction was performed by treating 1 g of the obtained powder with 20 mL of CHCl₃/MeOH (2:1 v/v). HPLC/MS analysis of the *Ericaria brachycarpa* extract (EBE) was performed as previously described (Faddetta et al., 2023). Mass spectrum data were analyzed for metabolites annotation using MassHunter Qualitative Analysis B.06.00 and the Metabolomic Workbench database

98 [https://www.metabolomicsworkbench.org/search/ms.php]. Before use, a mother solution of EBE
99 diluted 100 mg/mL in DMSO was prepared and then stored at – 20°C.

100 **2.2 *A. lixula* sampling, fertilization and exposure**

101 Adult *A. lixula* sea urchins were collected monthly in shallow rocky reefs (2– 8 m), from March to
102 June 2023, during their main reproductive peak (Wangensteen et al., 2013). All animal experiments
103 were carried out following the ARRIVE guidelines in accordance with the U.K. Animals (Scientific
104 Procedures) Act, 1986 and the EU Directive 2010/63/EU for animal experiments. For each
105 experiment, 10 adult-sized sea urchins (weight 41.1 ± 6.4 g) were brought to the tanks in laboratory
106 and fertilized within 24 h of collection. Gametes were collected from at least 3 males and 3 females
107 and examined under the microscope to select quality gametes later used for fertilization, as previously
108 described (Martino et al., 2021). Seawater was collected in the same area and filtered (0.22 µm
109 Millipore filtered seawater, hereafter MFSW). Ten minutes after fertilization, the fertilization success
110 (> 95 %) was inspected by microscopic examination. After three rinsing in MFSW, 20 mL of fertilized
111 eggs (4000 eggs/mL) were placed in 20 mm Petri dishes and exposed to different concentrations of
112 EBE (1, 4, 10, 20 and 40 µg/mL, hereafter EBE 1, EBE 4, EBE 10, EBE 20 and EBE 40), obtained
113 by dilution of the 100 µg/mL mother solution in MFSW, with 3 replicate dishes per treatment. A
114 schematic drawing of the experimental design is depicted in Figure 1 A. Embryos were reared at
115 18°C, with three developmental endpoints for each treatment: 24 h (G, gastrulae), 48 h (ePl, early
116 plutei) and 72 h (lPl, late plutei, see Fig. 1 A). Black rods in lines 1, 3 and 6 represent control
117 treatments at 24, 48 and 72 h (Fig. 1 A). Grey rods in lines 2, 4 and 7 represent the treatments for 24,
118 48 and 72 h with different concentrations of EBE added at 0 hours post fertilization (hpf, hereafter 0
119 hpf EBE 1/4/10/20/40, Fig. 1 A). Lines 5 and 8 represent treatments with EBE added at 24 and 48
120 hpf, respectively (hereafter 24 hpf EBE 1/4/10/20/40 and 48 hpf EBE 1/4/10/20/40) (see Fig. 1 A).
121 Line 9 represents the recovery experiment, with embryos reared for 24 h in the EBE 40 treatment and
122 then cultivated in MFSW till the 72 h endpoint (Fig. 1 A). To determine whether DMSO influenced

development, control embryos were exposed to the sole vector and the effect on development evaluated.

2.3 Sea urchin embryos morphological and morphometrical analysis

Every 24 h, 100 embryos were sampled, examined microscopically (Leica DCF420 C), photographed using a digital camera and scored for normal/abnormal development. A total of 100 embryos per each replicate dish per each fertilization experiment were scored. Since controls and embryos exposed to the lowest doses of EBE (EBE 1 and 4) reached the gastrula stage at 24 h and the four-armed pre-feeding echinopluteus stage at 48-72 h, using ImageJ (ver 1.46 r) we measured the changes of three morphological parameters to compare the ability to correctly grow and calcify: the full diameter at 24 h, the length of the post-oral arm (PO), and larval body width (BW) at 48 and 72 h (Fig. 1 B, $n = 11$). Post-oral arm length represents a well-known metric of echinopluteal growth and calcification and BW was used to analyze changes in larval shape in response to EBE exposure (Byrne et al., 2013; Martino et al., 2021; Wangenstein et al., 2013). It was not possible to obtain these morphometric measures at the higher doses (EBE 10, 20 and 40) because embryos presented major developmental aberrations.

2.4 SH-SY5Y cell culture, differentiation and exposure

Undifferentiated proliferating and differentiated SH-SY5Y cells were cultured as previously described by Nuzzo et al., 2023. To establish the EBE effect on cells viability, undifferentiated and differentiated cells were planted respectively at the density of 20000 and 14000 cells/well on 96-well plates in a final volume of 100 μ L/well and subsequently exposed to different EBE doses (1, 10, 40, 50, 80, 100, 200 μ g/mL) for 24 h, diluting the mother solution in growth medium. Then, a time course experiment was performed in the EBE 40 treatment at three different time points (24, 48 and 72 h). In all the experiments, control groups received an equal volume of the solvent only.

2.5 Cell treatment and MTT assay

Cell viability was evaluated after 3 h incubation at 37°C with 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT, 0.5 mg/mL, Sigma-Aldrich), as previously described (Nuzzo et al., 2021). The obtained formazan was dissolved with DMSO and the relative absorbance was read at 570 nm in a Thermo Scientific™ Multiskan™ GO microplate spectrophotometer.

2.6 Dose-Response Curves

To test whether the sensitivity of *A. lixula* embryos to EBE exposure differed with the developmental stage at which exposures were initiated, we performed three dose–response curves for 24 h treatments starting at 0, 24 and 48 hpf. Also, to test the toxic effects of EBE on neuroblastoma proliferating cells, SH-SY5Y cells were exposed for 24 h at different concentrations. The range of concentrations tested was chosen to cover the full 0-100 % curves, with doses ranging from 0 to 40 µg/mL for embryos and from 0 to 200 µg/ml for SH-SY5Y cells. The dose-response curves were calculated plotting the percentages of abnormal embryos/dead cells across increasing EBE concentrations fitting a four-parameter logistic function. Two ecotoxicological parameters were determined: EC₅₀, the concentration that gives the half-maximal response, and NOEC, the no observed effect concentration. Based on the EC₅₀ and NOEC values, EBE 40 was the dose chosen for subsequent investigations on neuroblastoma cells since it caused a significant reduction of undifferentiated SH-SY5Y cell viability and no effect on differentiated cells. For WB and TUNEL assays on embryos, two doses were chosen, i.e., EBE 4, a sub-lethal concentration below the EC₅₀ values, and EBE 40, a cytotoxic concentration causing 100 % block of first cleavage. Data were analyzed using GraphPad Prism 9.0.2. software (GraphPad Software, Inc., La Jolla, CA, USA).

2.7 Protein extraction, electrophoretic analysis and protein gel blotting

Protein extraction, electrophoresis, immunoblotting analysis and image acquisition were performed according to Chiarelli et al. (2022) for embryos and to Scordino et al. (2024) for SH-SY5Y cells. The following antibodies were used: HSP 60 (H3524 Sigma-Aldrich) diluted 1:500, HSP 90 (sc-13119 Santa Cruz Biotechnology) diluted 1:1000, Cleaved Caspase-7 (9491 Cell Signaling) diluted 1:1000, LC3 (L8918 Sigma-Aldrich) diluted 1:1000, p62/SQSTM1 (Sigma-Aldrich, P0068) diluted 1:500;

173 CHOP-10/GADD153 (sc-7351 Santa Cruz Biotechnology) diluted 1:300, β -actin (sc-47778 Santa
174 Cruz Biotechnology) diluted 1:10000. As secondary antibodies, we used the ECLTM anti-rabbit IgG
175 horseradish peroxidase linked whole antibody from donkey (GE Healthcare, NA934) or anti-mouse
176 from sheep (GE Healthcare, NA931) diluted 1:2500 with TBS-T for embryos, and goat anti-rabbit
177 IgG (sc-2004, Santa Cruz Biotechnology) horseradish peroxidase-conjugated diluted 1:8000, anti-
178 mouse IgG (sc-2005 Santa Cruz Biotechnology) diluted 1:6500 or Cy3 anti-rabbit IgG (115-165-003
179 Jackson ImmunoResearch) diluted 1:5000 for SH-SY5Y cells. In embryos, ponceau S staining was
180 used for total protein normalization, since the levels of housekeeping proteins, including actin,
181 changed during developmental phases and under different experimental conditions as many of these
182 proteins, including actin. In SH-SY5Y, β -actin was used for total protein normalization. Data are
183 presented as mean of optical density values $\pm$ standard deviation (SD) of triplicate experiments ($n =$
184 3).

185 **2.8 DNA fragmentation assay**

186 At all the experimental endpoints (see Fig. 1 A), 2 mL of the embryo cultures taken from each of the
187 three dishes per treatment were mixed and subsequently fixed with 4% paraformaldehyde for 1 h at
188 room temperature, washed in MFSW and stored until use in pure methanol at - 20 °C. SH-SY5Y cells
189 were grown in 12 mm diameter round glass coverslips at the density of $8 \cdot 10^4$. Cells exposed to EBE
190 40 were collected at three different time points, 24, 48 and 72 h, fixed with 4% paraformaldehyde for
191 1 h, washed twice with PBS 1X and processed for DNA fragmentation evaluation. DNA
192 fragmentation was analyzed, for both embryo cultures and SH-SY5Y cells, by TUNEL (TdT-
193 mediated dUTP nick-end labelling) assay (Promega) as described by Chiarelli et al. (2022). Positive
194 controls were pre-treated with DNAase before the TdT assay. Negative controls were incubated with
195 nucleotide mix (fluorescein 12-dUTP), without the TdT enzyme. The fragmented labelled chromatin
196 was observed under a 20X objective, using a fluorescence microscopy (Olympus BX50). Quantitative

197 analysis was carried out using ImageJ 1.46r software after acquisition through a digital camera (Nikon
198 Sight DS-U1).

199 **2.9 Statistical analysis**

200 For sea urchin embryos, statistical analysis was performed as described by Martino et al. (2021), with
201 some modifications. Briefly, since each dish had a different set of embryo genotypes and the
202 experiment was repeated three times resulting in 9 replicate dishes per each treatment, $n = 9$ (Martino
203 et al., 2021). Percentage data for each of the morphological categories, data on PO and BW lengths,
204 WB results and TUNEL assay results were analyzed by two-way analysis of variance (ANOVA), with
205 EBE concentration and the exposure starting time (hpf) as fixed factors. For SH-SY5Y cells, data
206 were collected from three independent experiments, with at least 3 replicates per treatment. For SH-
207 SY5Y cells, controls were considered equal to 1 in WB and to 100% in cell viability assays. Outliers
208 were identified using theROUT method ($Q=5\%$) and results were analyzed by one-way ANOVA.
209 Tukey's HSD test was used as Post-hoc test for mean comparison and homogeneity of variance was
210 checked and confirmed using Levene's test for embryos and Shapiro-Wilk's test for cells. For data
211 not normally distributed, statistical evaluations were performed by Kruskal-Wallis test, followed by
212 Dunn's multiple comparison test. The analyses were performed using the Statistica 13.2 software
213 (StatSoft, Tulsa, OK, USA) for embryos and GraphPad Prism 9.0.2. software (GraphPad Software,
214 Inc., La Jolla, CA, USA) for cells. $P < 0.05$ was set as the level of significance.

215 **3. Results**

216 **3.1 Characterization of the oxylipins in EBE**

217 Considering the potential interest toward oxylipins as agents of reproductive failure and antitumoral
218 compounds (Miralto et al., 1999; Luo et al., 2006; Christensen 2020), a qualitative analysis of the
219 lipids from EBE was performed by HPLC-MS in negative mode, allowing their characterization.
220 Seven oxygenated lipids were identified (Table 1), among these cyclic derivative oxo-phytodienoic

acid 1 and linear hydroxylated fatty acids 2-7 were found, similarly to what previously found in brown algae (Ritter et al., 2008). The carbon lengths of these oxylipins (C16 to C20) are in line with the vast majority of these compounds, derived from lipoxygenases metabolism of polyunsaturated fatty acids precursors found in marine organisms, that display carbon lengths ranging from C16 to C22 (Gerwick and Singh, 2002). Except for compound 1 (Oxo-phytodienoic acid), all the other compounds belong to the hydroxyacids (HEPEs) class of oxylipins. Although oxylipins are known from other macroalgal species (Barbosa et al., 2016), to our knowledge this is the first time they have been extracted from species of the *Ericaria* genus.

3.2. Dose-response curves

3.2.1 Sea urchin embryos

First, we established the general response to different concentrations of EBE in *A. lixula* embryos and checked whether the developmental exposure windows influenced EBE toxicity. The levels of normally developed embryos observed in controls were > 95% and did not significantly differ with those observed in DMSO treatments. Treatments were performed adding EBE just after fertilization (0 hpf) or at 24 and 48 hpf, thus respectively at the gastrula or pluteus stage, respectively, to test whether the sensitivity of embryos to EBE exposure differed with the developmental stage at which exposures were initiated (see Fig. 1 A). EBE exposure had a significant dose-dependent effect on the development of *A. lixula* embryos ($F_{17,36} = 581.2, p < 0.0001$), as well as the developmental stage at which exposures were initiated ($F_{17,36} = 128.4, p < 0.0001$) and their interaction (EBE x hpf: $F_{17,36} = 43.8; p < 0.0001$). Exposure from fertilization significantly reduced levels of normally developed larvae in parallel with the increase of the EBE concentration by 19.6, 63, and 100%, respectively for the exposure to EBE 4, 10 and 20. Strikingly, the 24 hpf EBE 1 treatment resulted in 21 % abnormal embryos already, while this dose had no effect in the 0 and 48 hpf treatments. On the contrary, while for the 0 and 24 hpf treatment there was 100 % abnormality at EBE 20, in the 48 hpf EBE 20 treatment there was a reduction in levels of normally developed larvae by 58.8 %. The dose-response curves,

obtained by plotting the percentages of abnormal development against the concentration used, are shown in Fig. 2 A and resulted in an EC₅₀ value of 8.0 µg/mL for the 0 hpf treatment (R squared = 0.9948), 5.3 µg/mL for the 24 hpf treatment (R squared = 0.9868) and 16.4 µg/mL for the 48 hpf treatment (R squared = 0.9903), while the NOEC values were respectively 0.5, 0.08 and 0.7 µg/mL.

3.2.2 SH-SY5Y cells

In order to establish the EBE effects on SH-SY5Y cell viability, we performed an MTT cell sensitivity assay, finding that 24 h EBE exposure produced a dose-dependent effect on undifferentiated SH-SY5Y cell viability ($F_{7, 99} = 147.4$, $p < 0.0001$, Fig. 2 B). No toxic effects on SH-SY5Y cells was observed at concentrations up to EBE 10, while a decrease of cell viability was observed above this dose. The dose-response curve showed an EC₅₀ value of 45.7 µg/mL (R squared = 0.9998) and a NOEC value of 14.4 µg/mL. No toxic effects were detected for doses lower than EBE 10, while maximum percentage of cell death was observed starting from EBE 80. EBE 40 was chosen for the next investigations. At this concentration, EBE exposure produced a significant time-dependent reduction in undifferentiated SH-SY5Y cell viability (Fig. 2 C), with 78 % of dead cells at 72 h, while no significant toxicity was observed in differentiated SH-SY5Y (Fig. 2 D), non-proliferating cells with functional and morphological characteristics similar to neurons. Taken together, these data demonstrated that at concentrations up to EBE 40 a specific toxic effect on undifferentiated SH-SY5Y cells but not on their neuronal-like cells derivatives, was exerted, suggesting a preferential selectivity for proliferating cells.

3.3. Morphological analysis of embryos exposed to EBE

3.3.1. Larval morphometry at low doses

EBE exposure from fertilization at the two concentrations below the EC₅₀ values (EBE 1 and 4) had no effect until gastrulation (24 h), as both controls and exposed embryos reached all the gastrula milestones with the correct timing, including biomineral deposition (Fig. 3 A). The diameter of

270 gastrulae was not significantly different between controls and EBE exposed embryos (not shown). At
271 48 and 72 h, controls were plutei showing a complex skeletal structure (Fig. 3 B, C). Since most
272 embryos exposed to low doses showed delayed development with a decrease in size (Fig. 3 D), it was
273 possible to measure their larval size, measured as post-oral rod lengths (PO) and body width (BW),
274 to compare with controls the ability of exposed larvae *vs* controls to correctly grow and calcify (see
275 Fig. 1 B).

276 In control larvae, while the post-oral (PO) rod showed a physiological 25 % increase in length from
277 $178.6 \pm 12.8 \mu\text{m}$ at 48 h to $221.9 \pm 20 \mu\text{m}$ at 72 h, no difference was observed in BW, showing that
278 the growth in the plutei size during development is mainly due to increased length and not in width,
279 eventually resulting in a slenderer larval shape (Fig. 3 B, C and Fig. 4).

280 At 48 h, exposure to EBE significantly affected both PO arm length ($F_{5,60} = 193.1, p < 0.0005$) and
281 larval body width (BW, $F_{5,60} = 28.2; p < 0.0005$), as well as the developmental stage at which
282 exposures were initiated (PO: $F_{5,60} = 165, p < 0.0005$; BW: $F_{5,60} = 12, p < 0.005$) and their interaction
283 (PO: $F_{5,60} = 42.5, p < 0.0005$; BW: $F_{5,60} = 14.75, p < 0.0005$). Larvae exposed from fertilization to
284 EBE 1 and 4 had respectively 15 % ($153.9 \pm 30.6 \mu\text{m}$) and 36 % ($116.7 \pm 30.4 \mu\text{m}$) shorter PO rods
285 at 48 h, if compared to controls (Fig. 4 A). Strikingly, the maximum reduction in PO length was
286 observed in the 24 hpf treatments at 48 h, with a 68 % reduction in PO arm length in the EBE 1
287 treatment ($58.1 \pm 5.5 \mu\text{m}$) and an 87 % reduction in the EBE 4 treatment ($23.9 \pm 7.08 \mu\text{m}$) if compared
288 to 48 h controls (Tukey's HSD PO 48 h: 24 hpf EBE 4 < 24 hpf EBE 1 < EBE 4 < EBE 1 < C, see
289 Fig. 3 D and Fig. 4 A). BW had a significant 17 % reduction in the EBE 4 treatment at 48 h ($141.6 \pm$
290 $8.4 \mu\text{m}$), but not at EBE 1, if compared to 48 h control plutei ($170.4 \pm 11.42 \mu\text{m}$). Strikingly, in the
291 24 hpf treatments there was a significant 23 % reduction in BW at EBE 1 ($138.8 \pm 12.9 \mu\text{m}$) and a 32
292 % reduction at EBE 4 ($122.5 \pm 15.1 \mu\text{m}$, Tukey's HSD BW 48 h: 24 hpf EBE 4 = 24 hpf EBE 1 =
293 EBE 4 < EBE 1 = C, see Fig. 4 A).

At 72 h, EBE exposure had a significant effect both on PO ($F_{5,60} = 45.4, p < 0.0001$) and BW ($F_{5,60} = 14.8; p < 0.0005$), as well as the developmental stage at which exposures were initiated (PO: $F_{5,60} = 77.2, p < 0.0005$; BW: $F_{5,60} = 4.31, p < 0.05$) and their interaction (PO: $F_{5,60} = 49.5, p < 0.0005$; BW: $F_{5,60} = 14.7, p < 0.0005$). Larvae in the 0 hpf EBE 1 and 4 treatments had respectively 13 % ($193.9 \pm 15.7 \mu\text{m}$) and 46 % ($121.2 \pm 13.7 \mu\text{m}$) shorter PO rods at 72 h, if compared to controls ($156.9 \pm 8.9 \mu\text{m}$) (Fig. 4 B, Tukey's HSD PO 72 h: EBE 4 < EBE 1 < C = 48 hpf EBE 1 = 48 hpf EBE 4). For BW, a significant 21 % reduction was observed at EBE 4 ($135.1 \pm 9.4 \mu\text{m}$), while no difference was present at EBE 1 (Fig. 4 B). Strikingly, no difference was present in the 48 hpf EBE 1 and 4 treatments either for PO or BW (Fig. 4 B, Tukey's HSD BW 72 h: EBE 4 < EBE 1 = C = 48 hpf EBE 1 = 48 hpf EBE 4).

304

3.3.2. Impairment of development at high doses

Retarded development and major developmental aberrations were observed at the three concentrations tested above the 0 hpf EC_{50} value (EBE 10, 20 and 40), showing a dose-dependent block of the first cleavage. Only 37 % of embryos exposed to EBE 10 normally developed to the pluteus stage at 48 h, while 33 % were delayed or abnormal embryos (Fig. 3 D, E) and 30 % showed cleavage inhibition (Fig. 3 F). Treatments with the highest dose (EBE 40) showed 100 % block of first cleavage and after 48 h apoptotic shrinkage and disassembly into apoptotic bodies were easily recognized (Fig. 3 G).

In the recovery experiment, embryos exposed to EBE 40 for 24 h and then transferred in MSFW for further 48 h showed 100 % cleavage inhibition with the first signals of apoptotic cell death, such as convoluted and extension-forming outlines (the phenotype shown in Fig. 3 F), demonstrating that once the apoptotic process is initiated after exposure to EBE 40, the cell will be inevitably led to death.

318 In the 24 and 48 hpf EBE 40 treatments, the progression through developmental stages resulted totally
319 blocked, as gastrulae showed acute toxicity (Fig. 3 H) and plutei displayed evident apoptotic features
320 with cells forming extensions and detaching from the skeletal structure (Fig. 3 I).

321 **3.4. EBE induces a modulation of Heat shock proteins**

322 **3.4.1 Sea urchin embryos**

323 Both EBE exposure ($F_{12,26} = 387.8$; $p < 0.0001$) and the developmental stage at which exposures were
324 initiated ($F_{12,26} = 248.2$; $p < 0.0001$) had a significant effect on HSP60 expression, as well as their
325 interaction (EBE x hpf: $F_{12,26} = 108.5$; $p < 0.0001$). While no increase was observed in the EBE 4
326 treatment, EBE 40 treatments induced the up-regulation of HSP 60 protein levels at 24 h but,
327 strikingly, was not present at 48 and 72 h (Fig. 5 A), suggesting the activation of other processes. On
328 the contrary, larvae in the 24 and 48 hpf treatments at both EBE concentrations had significantly ~2-
329 fold higher HSP 60 levels if exposed from the gastrula stage and ~5-fold higher levels if exposed
330 from the pluteus stage, compared to 48 and 72 h controls (Fig. 5 A).

331 **3.4.2 SH-SY5Y cells**

332 As shown in Fig. 5 B and C, exposure of undifferentiated SH-SY5Y cells for 24 h to EBE 40 treatment
333 induced a significant down regulation of HSP 60 and HSP 90. A significant 24 % and 34% reduction
334 in HSP 60 protein levels was respectively observed at 24 h and 48 h, and a further decrease up to 60
335 % was observed at 72 h ($F_{3,36} = 16.09$; $p < 0.0001$). HSP 90 protein levels showed a significant
336 progressive reduction from 43 % at 24 h up to 84 % at 72 h ($F_{3,36} = 22.38$; $p < 0.0001$).

337 **3.5 EBE induces a dose-dependent apoptotic-mediated cell death**

338 **3.5.1 Cleaved caspase-7 and CHOP-10/GADD153 proteins**

339 **3.5.1.1 Sea urchin embryos**

Both EBE exposure ($F_{12,26} = 946.9$ $p < 0.0001$) and the hpf treatments ($F_{12,26} = 680.9$; $p < 0.0001$) had a significant effect on caspase-7 expression, as well as their interaction (EBE x hpf: $F_{12,26} = 593.9$; $p < 0.0001$). While only low levels of the protein were present in most treatments, embryos exposed to EBE 40 had a 13-fold increase at 48 h and a 108-fold increase at 72 h, if compared to controls (Fig. 6 A). Strikingly, also embryos exposed to EBE 40 from the gastrula and pluteus stages had respectively a 57- and a 17-fold increase (Fig. 6 A), respectively, demonstrating the massive induction of the execution-phase of cell apoptosis in embryos exposed to the highest dose of EBE.

For CHOP-10/GADD153 expression, WB analysis showed that EBE exposure ($F_{12,26} = 105.5$; $p < 0.0001$) and the developmental stage at which exposures were initiated ($F_{12,26} = 87.5$; $p < 0.0001$) had a significant effect on its protein levels, as well as their interaction (EBE x hpf: $F_{12,26} = 27.2$; $p < 0.0001$). Strikingly, CHOP-10/GADD153 was upregulated in the EBE 40 treatment at 24 h but not present at 48 and 72 h (Fig. 6 A), when on the contrary there was a massive expression of caspase-7, likely because the action of CHOP-10/GADD153 in cell cycle arrest is temporally precedent to that of caspase-7. Exposure from the gastrula and pluteus stages also induced higher levels of the CHOP-10/GADD153 protein at both concentrations (Fig. 6 A). We also observed a 2-fold increase in the EBE 4 treatment at 72 h if compared to 72 h controls (Fig. 6 A).

3.5.1.2 SH-SY5Y cells

Similarly, EBE had a significant effect on CHOP-10/GADD153 expression ($F_{3,34} = 6.734$; $p = 0.0011$) and on caspase-7 expression ($H_{30} = 17.1$; $p = 0.0007$) in undifferentiated SH-SY5Y cells. WB analysis showed very low protein levels of both CHOP-10/GADD153 and cleaved isoform of Caspase-7 in control group (Fig. 6 D). Otherwise, a 10-fold increase of CHOP-10/GADD153 was observed in the EBE 40 treatment at 24 h and a significant increase of cleaved isoform of caspase-7 was detected in the EBE 40 treatment at 48 h and 72 h (Fig. 6 D).

3.5.2 Chromatin fragmentation

364 3.5.2.1 Sea urchin embryos

365 Results of total fragmented apoptotic DNA displayed by TUNEL assay corroborated the WB data
366 about apoptosis. Fig. 6 B-C shows equatorial optical sections of representative larvae at 48/72 h and
367 the histogram reporting the amount of apoptotic fragmented DNA. Both EBE exposure ($F_{9,20} = 578$;
368 $p < 0.0001$) and the developmental stage at which exposures were initiated ($F_{9,20} = 93.4$; $p < 0.0001$)
369 had a significant effect on the amount of apoptotic fragmented DNA, as well as their interaction (EBE
370 x hpf: $F_{9,20} = 50.3$; $p < 0.0001$). As previously reported in *P. lividus* larvae, low levels of apoptotic
371 fragmented DNA were present in controls both at 48 and 72 h (Chiarelli et al., 2021; Martino et al.,
372 2021, see Fig. 6 B), indicating only basal physiological apoptosis. While no difference was present
373 between controls and larvae in the EBE 4 treatment at 48 h, a significant 10-fold increase was
374 highlighted at 72 h, hallmark of a time-dependent activation of apoptosis (Fig. 6 B). No caspase-7
375 signal was evidenced by WB analysis in the EBE 4 treatment at 72 h since cells were in an advanced
376 stage of apoptosis, where caspase-7 role is already ended but fragmented DNA is still present and
377 visible. Strikingly, the EBE 40 treatment at 48 h determined a 55-fold increase compared to controls,
378 indicating the presence of advanced apoptosis (Fig. 6 B). It was not possible to analyze EBE 40
379 treatment at 72 h by TUNEL because embryos were disassembled into small apoptotic bodies.

380 Embryos exposed from the gastrula and pluteus stages had a dose-dependent increase of chromatin
381 fragmentation, with a significant 10- and 54- fold increase in the 24 hpf EBE 4 and 40 treatments,
382 and a significant 9.5- and 20.5- fold increase in the 48 hpf EBE 4 and 40 treatments, respectively
383 (Fig. 6 B).

384 Regardless the exposure starting time, embryos exposed to EBE 4 displayed apoptotic nuclei with
385 fragmented DNA only in a restricted group of cells (Fig. 6 C), while embryos exposed to EBE 40
386 showed massive apoptosis, widespread throughout the embryo (Fig. 6 C).

387 3.5.2.2 Chromatin fragmentation in SH-SY5Y cells

EBE exposure ($F_{3,8} = 61$; $p < 0.0001$) led to a significant increase in the amount of apoptotic fragmented DNA in undifferentiated SH-SY5Y cells. As shown in Fig. 6 E-F, no signal of fragmented DNA was detected in control cells. Cells exposed to EBE 40 showed a time-dependent response, as the percentage of fragmented nuclei was 3.5 %, 21 % and 34 % in cells treated for 24, 48 and 72 h, respectively (Fig. 6 E-F).

3.6.EBE modulates the autophagic proteins LC3 and p62/SQSTM1

3.6.1 Sea urchin embryos

The amount of these proteins showed that EBE exposure significantly affected their expression (LC3: $F_{12,26} = 609.3$, $p < 0.0001$; p62: $F_{12,26} = 969.3$, $p < 0.0001$), as well as the exposure starting time (LC3: $F_{12,26} = 246.2$, $p < 0.0001$; p62: $F_{12,26} = 1004.1$, $p < 0.0001$) and their interaction (LC3: $F_{12,26} = 177.6$, $p < 0.0001$; p62: $F_{12,26} = 530$, $p < 0.0001$). The two proteins had similar expression profiles, being down-regulated in the EBE 40 treatments from fertilization (Fig. 7 A). In treatments starting at the gastrula and pluteus stages these proteins showed opposite trends, as LC3 was upregulated in the EBE 4 treatments with a ~4-fold increase while p62/SQSTM1 was upregulated in the EBE 40 treatments with a ~12-fold increase at 48 h and a 2.4-fold increase at 72 h (Fig. 7 A).

3.6.2 SH-SY5Y cells

The WB analysis showed that EBE 40 treatment induced autophagy in undifferentiated SH-SY5Y cells ($F_{3,36} = 8.578$, $p = 0.0002$), as seen by a significant 2.4-fold increase in LC3 levels following 48 h exposure (Fig. 7 B).

4. Discussion

Our results provide morphological and molecular evidence of the toxic effects of the oxylipin-containing *E. brachycarpa* extract on *A. lixula* embryos and larvae and on neuroblastoma cells.

411 Oxylipins' diversity in chemical structure and the associated biological effects make them particularly
412 promising for therapeutic strategies, as widely seen for oxylipins derived from plants and microalgae
413 (Blasio and Balzano, 2021; Savchenko et al., 2022), while only few studies exist about the beneficial
414 properties of oxylipins derived from macroalgae (Barbosa et al., 2016). Here, for the first time, we
415 identified seven oxylipins in the extract of a species of the *Ericaria* genus, providing a new insight
416 into the secondary metabolites produced by these brown algae and their potential as therapeutic agents
417 and larvicidal compounds.

418 Since macroalgae can secrete their secondary metabolites either by direct contact for compounds
419 contained in epidermal glands and secretory trichomes or through masses of water (Gomes et al.,
420 2017), it is likely that sea urchin embryos and larvae may come into direct contact with these
421 molecules. An important finding of this study was that EBE was able to induce a marked delay in
422 development at concentrations below the EC_{50} values, namely EBE 1 and 4. We quantified this delay
423 measuring the decrease of PO and BW lengths and used these values as a proxy of larval calcification
424 (Byrne et al., 2013). Our results showed a major reduction in PO length for larvae exposed from
425 fertilization with respect to controls, posing a threat to larval fitness due to impaired feeding and
426 swimming (Byrne et al., 2013; Martino et al., 2021). Strikingly, the maximum reduction in PO length
427 was observed in the 24 hpf treatments at 48 h, while no difference was present in the 48 hpf EBE 1
428 and 4 treatments either for PO or BW. These results are in agreement with the developmental stage-
429 specific sensitivity highlighted by the three dose-response curves, confirming gastrulae as the most
430 sensitive stage to the extract with the lowest EC_{50} (5.366 $\mu\text{g/mL}$) and plutei as the most resistant stage
431 with an EC_{50} equal to 16.46 $\mu\text{g/mL}$, probably because of their incomplete development and
432 suboptimal protective mechanisms, such as the induction of HSPs and autophagy (see below).
433 Exposure to the highest concentration (EBE 40) caused a 100 % inhibition of cleavage and inevitably
434 lead to an apoptotic-driven cell death regardless the exposure starting time, with no possibility of
435 recovery.

436 Several biomarkers were used to detect the activation of the cellular stress response to EBE in *A.*
437 *lixula* embryos, showing that it strongly depended on the developmental exposure window. The first
438 cellular stress response mechanisms activated were the induction of HSP 60 and autophagy, similarly
439 to our previous results on the hierarchical cellular stress response of *P. lividus* embryos exposed to
440 environmental stressors (Matranga et al., 2013; Chiarelli et al., 2016). Larvae exposed from the
441 pluteus stage had a HSP 60 induction greater than larvae exposed from the gastrula stage, probably
442 accounting for the better resistance of plutei. As seen from the LC3 and p62/SQSTM1 markers,
443 autophagy was massively induced in both 24 and 48 hpf treatments, indicating the effort of gastrulae
444 and plutei to remove toxic metabolites, misfolded/damaged proteins and organelles. On the contrary,
445 in the 0 hpf treatments we found a time-dependent decrease of autophagy, in parallel with the
446 induction of apoptosis as seen with two markers: cleaved caspase-7, known to play a key role in the
447 execution-phase of apoptosis, and CHOP-10/GADD153, whose overexpression is linked to cell cycle
448 arrest leading to apoptosis. The *in situ* analysis of fragmented DNA showed that apoptosis was acting
449 either as a selective process at low doses (EBE 4) affecting only some cells, likely the most damaged,
450 or as a process involving the whole embryo at high doses (EBE 40), leading to death. This selective
451 apoptosis is reasonably acting to preserve the developmental program since other cells may benefit
452 from the materials liberated by apoptotic cells, e.g. amino acids and substrates for ATP (Somero,
453 2020), as was observed in *P. lividus* embryos exposed to vanadium and to the combined exposure of
454 gadolinium and increased temperatures (Chiarelli et al., 2021; Martino et al., 2021).

455 To interpretate why this seaweed should produce natural toxins with a toxic effect on sea urchin
456 embryos, we have to consider the ecological interactions between these two species, with the seaweed
457 producing chemical substances against its grazers and the sea urchin embryos adopting molecular
458 strategies to respond to these environmental natural toxins. Oxylipins are the probable agents of this
459 reproductive failure in *A. lixula*, similarly to the stunting action generated by diatoms-derived
460 oxylipins on *P. lividus* and copepods reproduction (Miralto et al., 1999; Varrella et al., 2014). As *A.*
461 *lixula* normally grazes on *E. brachycarpa* present in its environment, the harmful effect on embryos

462 but not on adults suggests that there may be greater vulnerability of the embryos, similarly to
463 copepods that can be successfully cultivated from embryo to adult stage on diatom food (Koski et al.,
464 1998). Extracts from different brown algae, including *Ericaria crinita*, showed larvicidal effects
465 against the larvae of different insects (Fouda et al., 2022; Aly et al., 2023), suggesting their potential
466 as natural resources for insecticides production.

467 As previously mentioned, oxylipins have been shown to have toxic effects on several tumoral cell
468 line (Cohen and Flescher, 2009; Christensen, 2020), suggesting potential therapeutic uses as anti-
469 cancer drug. Our study showed that the treatment of neuroblastoma tumoral cells with the oxylipin-
470 containing *E. brachycarpa* extract induced a dose-dependent reduction in cells viability, with 40
471 µg/mL killing 78 % of the cells at 72 h, while no toxic effect was detected at this dose on the neuronal-
472 like cells derivatives, demonstrating the EBE selectivity versus the proliferating phenotype. This
473 selective effect of EBE on proliferating cells may be associated with a decrease of the HSPs,
474 molecular chaperones involved in maintaining the structure of proteins under different stress
475 conditions (Hu et al., 2022). Both HSP 60 and HSP 90 play a key role in tumorigenesis by regulating
476 protein homeostasis in response to stress, and inhibitors of these chaperons are widely considered as
477 cancer therapeutics (Park et al., 2020; Zhou et al., 2020). The strong and progressive reduction of
478 both HSP 60 and HSP 90 in response to EBE treatment reveals a new mechanism underlying the
479 potential anti-neoplastic activity of this extract of macroalgae containing oxylipins. Inhibition of HSP
480 60 and HSP 90, associated with the lack of HSP 70 expression in SH-SY5Y cells (Scordino et al.,
481 2023) leads to the accumulation of unfolded or misfolded proteins in the endoplasmic reticulum,
482 triggering the activation of a defense mechanism known as unfolded protein response, finally leading
483 to apoptosis of SH-SY5Y cells (Read et al., 2021). According to our results, EBE treatment produces
484 a significant increase in CHOP/GADD153, a protein involved in ER stress-mediated apoptosis
485 (Oyadomari and Mori 2004), and the consequent increase of the cleaved form of Caspase-7. The
486 activation of apoptosis was also confirmed by the increase in the percentage of fragmented DNA.

487 Interestingly, we observed an increase in LC3 expression in response to 48 h EBE treatment,
488 suggesting the induction of the autophagy, a process that can be stimulated by several types of cellular
489 stress, including ER stress. However, the role of autophagy induced by ER stress is controversial and
490 likely dependent on the extent of ER stress (Rashid et al., 2015). Although most studies indicate that
491 autophagy is a conserved cellular process that maintains cellular homeostasis, exerting pro-survival
492 functions following ER stress, it has been also demonstrated that autophagy and apoptosis often occur
493 in the same cell, mostly in a sequence in which autophagy precedes and facilitates apoptosis (Gump
494 and Thorburn, 2011; Song et al, 2017) Here, the increase in the autophagic process in response to
495 EBE treatment may indicate an attempt of proliferating cells to avoid being killed, or the effect of an
496 intense ER stress which leads to autophagy-dependent cell apoptosis. Therefore, more research is
497 needed to clarify the complex interplay between autophagy and apoptosis in EBE-exposed SH-SY5Y
498 cells.

499 **5. Conclusions**

500 Our study is the first to test an oxylipin-containing macroalgae extract on invertebrate reproduction,
501 giving new perspectives to elucidate the molecular strategies that marine invertebrates use when
502 responding to their environmental natural toxins. Moreover, this is the first time that an oxylipins-
503 containing extract from a macroalgae shows antineoplastic activity in a neuroblastoma cell line,
504 showing the potential to selectively trigger apoptosis of cancer cells but not in the corresponding
505 normal cells. Further studies are currently being carried out to fully characterize the chemical
506 composition of EBE and to isolate each compound. Once the effect of single molecules on embryo
507 development and cancer cells will be well known, it will be possible to synthesize the molecules with
508 the best anti-growth activities and further explore their potential as a prospective source for the
509 development of innovative, environmentally friendly larvicides and antineoplastic compounds.

510

511 **Figure captions**

512 Figure 1. A) Schematic drawing of the experimental design for *A. lixula* embryos exposure. Black
513 rods, culture in MFSW; grey rods, culture in EBE-containing MFSW. F, fertilization; G, gastrula; ePl,
514 early pluteus; lPl, late pluteus. Black arrows indicate the start of EBE exposure in the different
515 treatments; white arrow indicates the start of recovery in MFSW. B) Morphological traits measured
516 on *A. lixula* plutei: BW, body width; PO, post-oral skeletal rod. Bar: 50 μ m.

517 Figure 2. Dose-response curves across increasing EBE concentrations for percentages of: A)
518 abnormal embryos exposed to EBE from 0, 24 and 48 hpf; B) dead undifferentiated SH-SY5Y cells.
519 Time-course analysis of cell viability following EBE 40 exposure for 24, 48 and 72 h in: C)
520 undifferentiated SH-SY5Y cells and D) differentiated SH-SY5Y cells. Treatments with the same letter
521 do not differ (Tukey HSD).

522 Figure 3. Morphotypes in control (A-C) and EBE-exposed embryos (D-I). A) control embryo at 24
523 h; B) control embryo at 48 h; C) control embryo at 72 h; D) delayed embryo at 48 h; E) malformed
524 embryo at 48 h; F) zygote with segmentation inhibition at 24 h; G) apoptotic shrinkage and
525 disassembly into apoptotic bodies at 72 h; H) acute toxicity in the 24 hpf EBE 40 treatment; I)
526 apoptotic cell death in 48 hpf EBE 40 plutei. Bar = 60 μ M in A-B, D-H; 75 μ M in C, I.

527 Figure 4. Larval post-oral arm length and body width measured at 48 (A) and 72 h (B) in response to
528 EBE exposure. Treatments with the same letter do not differ (Tukey HSD).

529 Figure 5. Western blot analysis of protein levels for: A) HSP 60 in sea urchin embryos; B) HSP 60
530 and C) HSP 90 after EBE 40 exposure in SH-SY5Y cells. Treatments with the same letter do not
531 differ (Tukey HSD).

532 Figure 6. Induction of apoptosis in sea urchin embryos and in SH-SY5Y cells. Western blot analysis
533 of protein levels for CHOP10/GADD153 and Cleaved Caspase-7 in A) sea urchin embryos and D)
534 EBE 40-exposed SH-SY5Y cells. The histograms report the quantification of fluorescence related to

535 fragmented DNA in B) sea urchin embryos and E) SH-SY5Y cells. Treatments with the same letter
 536 do not differ (Tukey HSD). Merge of signals from total nuclei stained with propidium iodide and
 537 from nuclei with DNA fragmentation (see bright spots indicated by arrows) showing: C) Images of
 538 representative embryos at 48 (a-e) and 72 h (f-i) of development. Control embryo at 48 h (a); exposed
 539 embryos at 48 h: EBE 4 (b); EBE 40 (c); 24 hpf EBE 4 (d); 24 hpf EBE 40 (e); control embryo at 72
 540 h (f); exposed embryos at 72 h: EBE 4 (g); 48 hpf EBE 4 (h); 48 hpf EBE 40 (i). Positive control
 541 embryo (j). Negative control embryo (k). F) Images of SH-SY5Y cells. Control (a) and EBE 40-
 542 exposed cells at 24 (b), 48 (c) and 72 h (d). Positive control cells (e). Negative control cells (f). Bar
 543 = 60 μ m in C) and 45 μ m in F).

544 Figure 7. Western blot analysis of protein levels for: A) LC3 and B) p62/SQSTM1 in sea urchin
 545 embryos; C) LC3 after EBE 40 exposure in SH-SY5Y cells. Treatments with the same letter do not
 546 differ (Tukey HSD).

547

548 **Acknowledgments.** The authors would like to thank the National Biodiversity Future Center –
 549 NBFC, Project funded under the National Recovery and Resilience Plan (NRRP), Mission 4
 550 Component 2 Investment 1.4 - Call for tender No. 3138 of 16 December 2021, rectified by Decree
 551 n.3175 of 18 December 2021 of Italian Ministry of University and Research funded by the European
 552 Union – NextGenerationEU; Award Number: Project code CN_00000033, Concession Decree No.
 553 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUP
 554 B73C22000790001, Project title “National Biodiversity Future Center - NBFC” .

555 **Declaration of Competing Interest.** The authors declare no conflict of interest.

556

557 References

558 Adarshan, S., Sree, V.S.S., Muthuramalingam, P., Nambiar, K.S., Sevanan, M., Satish, L., Venkidasamy,
 559 B., Jeelani, P.G., Shin, H., 2023. Understanding Macroalgae: A Comprehensive Exploration of

560 Nutraceutical, Pharmaceutical, and Omics Dimensions. *Plants* (Basel) 13.
561 <https://doi.org/10.3390/plants13010113>
562
563 Aly, S.H., Elissawy, A.M., Salah, D., Alfuhaid, N.A., Zyaan, O.H., Mohamed, H.I., Singab, A.N.B., Farag,
564 S.M., 2023. Phytochemical Investigation of Three. *Mar. Drugs* 21. <https://doi.org/10.3390/md21020117>
565
566 Amico, V., 1995. Marine brown algae of family *Cystoseiraceae*: Chemistry and chemotaxonomy.
567 *Phytochem.* 39, 1257-1279.
568
569 Ávila-Román, J., Talero, E., de Los Reyes, C., Zubía, E., Motilva, V., García-Mauriño, S., 2016. Cytotoxic
570 Activity of Microalgal-derived Oxylipins against Human Cancer Cell lines and their Impact on ATP
571 Levels. *Nat. Prod. Commun.* 11, 1871-1875. <https://doi.org/10.1177/1934578X1601101225>
572
573 Barbosa, M., Valentão, P., Andrade, P.B., 2016. Biologically Active Oxylipins from Enzymatic and
574 Nonenzymatic Routes in Macroalgae. *Mar. Drugs* 14, 23. <https://doi.org/10.3390/md14010023>
575
576 Blasio, M., Balzano, S., 2021. Fatty Acids Derivatives From Eukaryotic Microalgae, Pathways and
577 Potential Applications. *Front. Microbiol.* 12, 718933. <https://doi.org/10.3389/fmicb.2021.718933>
578
579 Bonaviri, C., Vega Fernández, T., Fanelli, G., Badalamenti, F., Gianguzza, P., 2011. Leading role of the sea
580 urchin *Arbacia lixula* in maintaining the barren state in southwestern Mediterranean. *Mar. Biol.* 158, 2505-
581 2513. <https://doi.org/10.1007/s00227-011-1751-2>
582
583 Budzałek, G., Śliwińska-Wilczewska, S., Wiśniewska, K., Wochna, A., Bubak, I., Latała, A., Wiktor, J.M.,
584 2021. Macroalgal Defense against Competitors and Herbivores. *Int. J. Mol. Sci.* 22.
585 <https://doi.org/10.3390/ijms22157865>
586
587 Byrne, M., Lamare, M., Winter, D., Dworjanyn, S.A., Uthicke, S., 2013. The stunting effect of a high CO₂
588 ocean on calcification and development in sea urchin larvae, a synthesis from the tropics to the poles.
589 *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 368, 20120439 <https://doi.org/10.1098/rstb.2012.0439>.
590
591 Chiarelli, R., Martino, C., Agnello, M., Bosco, L., Roccheri, M.C., 2016. Autophagy as a defense strategy
592 against stress: focus on *Paracentrotus lividus* sea urchin embryos exposed to cadmium. *Cell Stress*
593 *Chaperones* 21, 19-27. <https://doi.org/10.1007/s12192-015-0639-3>
594
595 Chiarelli, R., Martino, C., Roccheri, M.C., Cancemi, P., 2021. Toxic effects induced by vanadium on sea
596 urchin embryos. *Chemosphere* 274, 129843.
597
598 Christensen, L.P., 2020. Bioactive C. *Molecules* 25. <https://doi.org/10.1016/j.chemosphere.2021.129843>
599
600 Cohen, S., Flescher, E., 2009. Methyl jasmonate: a plant stress hormone as an anti-cancer drug. *Phytochem.*
601 70, 1600-1609. <https://doi.org/10.1016/j.phytochem.2009.06.007>
602
603 Conte, M., Fontana, E., Nebbioso, A., Altucci, L., 2020. Marine-Derived Secondary Metabolites as
604 Promising Epigenetic Bio-Compounds for Anticancer Therapy. *Mar. Drugs* 19.
605 <https://doi.org/10.3390/md19010015>
606
607 Faddetta, T., Polito, G., Abbate, L., Alibrandi, P., Zerbo, M., Caldiero, C., Reina, C., Puccio, G., Vaccaro,
608 E., Abenavoli, M.R., Cavalieri, V., Mercati, F., Palumbo Piccionello, A., Gallo, G., 2023. Bioactive
609 Metabolite Survey of Actinobacteria Showing Plant Growth Promoting Traits to Develop Novel
610 Biofertilizers. *Metabolites* 13. <https://doi.org/10.3390/metabo13030374>

- Fontana, A., d'Ippolito, G., Cutignano, A., Miralto, A., Ianora, A., Romano, G., Cimino, G., 2007. Chemistry of oxylipin pathways in marine diatoms. *Pure Appl. Chem.* 79(4), 481-490. <https://doi.org/10.1351/pac200779040481>
- Fouda, A., Eid, A.M., Abdel-Rahman, M.A., El-Belely, E.F., Awad, M.A., Hassan, S.E., Al-Faifi, Z.E., Hamza, M.F., 2022. Enhanced Antimicrobial, Cytotoxicity, Larvicidal, and Repellence Activities of Brown Algae. *Front. Bioeng. Biotechnol.* 10, 849921. <https://doi.org/10.3389/fbioe.2022.849921>
- Gerwick, W.H., 1996. Epoxy allylic carbocations as conceptual intermediates in the biogenesis of diverse marine oxylipins. *Lipids* 31, 1215-1231. <https://doi.org/10.1007/BF02587906>
- Gomes, M.P., Garcia, Q.S., Barreto, L.C., Pimenta, L.P.S., Matheus, M.T., Figueredo, C.C., 2017. Allelopathy: An overview from micro-to macroscopic organisms, from cells to environments, and the perspectives in a climate-changing world. *Biologia* 72, 113–129. <https://doi.org/10.1515/biolog-2017-0019>
- Gump, J.M., Thorburn, A., 2011. Autophagy and apoptosis: what is the connection? *Trends Cell Biol.* 21, 387-392. <https://doi.org/10.1016/j.tcb.2011.03.007>
- Hu, C., Yang, J., Qi, Z., Wu, H., Wang, B., Zou, F., Mei, H., Liu, J., Wang, W., Liu, Q., 2022. Heat shock proteins: Biological functions, pathological roles, and therapeutic opportunities. *Med. Comm.* 3, e161. <https://doi.org/10.1002/mco2.161>
- Jacquemoud, D., Pohnert, G., 2015. Extraction and Analysis of Oxylipins from Macroalgae Illustrated on the Example *Gracilaria vermiculophylla*. *Methods Mol. Biol.* 1308, 159-172.
- Koski, M., Klein, B.W., Schogt, N., 1998. Effect of food quality on rate of growth and development of the pelagic copepod *Pseudocalanus elongatus* (Copepoda, Calanoida). *Mar. Ecol. Prog. Ser.* 170:169–187.
- Lauritano, C., Ianora, A., 2020. Chemical Defense in Marine Organisms. *Mar. Drugs.* 18;18(10):518. doi: 10.3390/md18100518.
- Lawrence, J. M., Agatsuma, Y., 2013. Tripneustes, in: *Developments in aquaculture and fisheries science.* Elsevier 38, 491-507.
- Librizzi, M., Martino, C., Mauro, M., Abruscato, G., Arizza, V., Vazzana, M., Luparello, C., 2023. Natural Anticancer Peptides from Marine Animal Species: Evidence from In Vitro Cell Model Systems. *Cancers (Basel)* 16. <https://doi.org/10.3390/md18100518>
- Luo, X., Li, F., Hong, J., Lee, C.O., Sim, C.J., Im, K.S., Jung, J.H., 2006. Cytotoxic oxylipins from a marine sponge *Topsentia* sp. *J. Nat. Prod.* 69, 567-571. <https://doi.org/10.1021/np0503552>
- Martino, C., Byrne, M., Roccheri, M.C., Chiarelli, R., 2021. Interactive effects of increased temperature and gadolinium pollution in *Paracentrotus lividus* sea urchin embryos: a climate change perspective. *Aquat. Toxicol.* 232, 105750. <https://doi.org/10.1016/j.aquatox.2021.105750>
- Matranga, V., Pinsino, A., Bonaventura, R., Costa, C., Karakostis, K., Martino, C., Russo, R., Zito, F., 2013. Cellular and molecular bases of biomineralization in sea urchin embryos. *Cah. Biol. Mar.* 54, 467–478.

661 Miralto, A., Barone, G., Romano, G., Poulet, S. A., Ianora, A., Russo, G. L., Buttino, I., Mazzarella, G.,
662 Laabir, M., Giacobbe, M.G., 1999. The insidious effect of diatoms on copepod reproduction. *Nature*,
663 402(6758), 173-176.
664

665 Nuzzo, D., Frinchi, M., Giardina, C., Scordino, M., Zuccarini, M., De Simone, C., Di Carlo, M., Belluardo,
666 N., Mudò, G., Di Liberto, V., 2023. Neuroprotective and Antioxidant Role of Oxotremorine-M, a Non-
667 selective Muscarinic Acetylcholine Receptors Agonist, in a Cellular Model of Alzheimer Disease. *Cell*.
668 *Mol. Neurobiol.* 43, 1941-1956.
669

670 Nuzzo, D., Picone, P., Giardina, C., Scordino, M., Mudò, G., Pagliaro, M., Scurria, A., Meneguzzo, F.,
671 Ilharco, L.M., Fidalgo, A., Alduina, R., Presentato, A., Ciriminna, R., Di Liberto, V., 2021. New
672 Neuroprotective Effect of Lemon IntegroPectin on Neuronal Cellular Model. *Antioxidants* (Basel) 10.
673 <https://doi.org/10.3390/antiox10050669>
674

675 Oyadomari, S., Mori, M., 2004. Roles of CHOP/GADD153 in endoplasmic reticulum stress. *Cell Death*
676 *Differ.* 11, 381-389.
677

678 Park, H.K., Yoon, N.G., Lee, J.E., Hu, S., Yoon, S., Kim, S.Y., Hong, J.H., Nam, D., Chae, Y.C., Park,
679 J.B., Kang, B.H., 2020. Unleashing the full potential of Hsp90 inhibitors as cancer therapeutics through
680 simultaneous inactivation of Hsp90, Grp94, and TRAP1. *Exp. Mol. Med.* 52, 79-91.
681

682 Pereira, L., Cotas, J., 2023. Therapeutic Potential of Polyphenols and Other Micronutrients of Marine
683 Origin. *Mar. Drugs* 21. <https://doi.org/10.3390/md21060323>
684

685 Privitera, D., Chiantore, M., Mangialajo, L., Glavic, N., Kozul, W., Cattaneo-Vietti, R., 2008. Inter-and
686 intra-specific competition between *Paracentrotus lividus* and *Arbacia lixula* in resource-limited barren
687 areas. *J. Sea Res.* 60(3), 184-192. <https://doi.org/10.1016/j.seares.2008.07.001>
688

689 Rashid, H.O., Yadav, R.K., Kim, H.R., Chae, H.J., 2015. ER stress: Autophagy induction, inhibition and
690 selection. *Autophagy* 11, 1956-1977. <https://doi.org/10.1080/15548627.2015.1091141>
691

692 Read, A., Schröder, M., 2021. The Unfolded Protein Response: An Overview. *Biology* (Basel) 10.
693 <https://doi.org/10.3390/biology10050384>
694

695 Ritter, A., Goulitquer, S., Salaün, J. P., Tonon, T., Correa, J. A., Potin, P., 2008. Copper stress induces
696 biosynthesis of octadecanoid and eicosanoid oxygenated derivatives in the brown algal kelp *Laminaria*
697 *digitata*. *New Phytol.* 180(4), 809–821. <https://doi.org/10.1111/j.1469-8137.2008.02626.x>
698

699 Romano, G., Miralto, A., Ianora, A., 2010. Teratogenic effects of diatom metabolites on sea urchin
700 *Paracentrotus lividus* embryos. *Mar. Drugs* 8, 950-967. <https://doi.org/10.3390/md8040950>
701

702 Ruocco, N., Annunziata, C., Ianora, A., Libralato, G., Manfra, L., Costantini, S., Costantini, M., 2019.
703 Toxicity of diatom-derived polyunsaturated aldehyde mixtures on sea urchin *Paracentrotus lividus*
704 development. *Sci. Rep.* 9, 517. <https://doi.org/10.1038/s41598-018-37546-y>
705

706 Sansone, C., Braca, A., Ercolesi, E., Romano, G., Palumbo, A., Casotti, R., Francone, M., Ianora, A., 2014.
707 Diatom-derived polyunsaturated aldehydes activate cell death in human cancer cell lines but not normal
708 cells. *PloS one*, 9(7), e101220. <https://doi.org/10.1371/journal.pone.0101220>
709

710 Savchenko, T., Degtyaryov, E., Radzyukevich, Y., Buryak, V., 2022. Therapeutic Potential of Plant
711 Oxylinins. *Int. J. Mol. Sci.* 23. <https://doi.org/10.3390/ijms232314627>

- Scordino, M., Frinchi, M., Urone, G., Nuzzo, D., Mudò, G., Di Liberto, V., 2023. Manipulation of HSP70-SOD1 Expression Modulates SH-SY5Y Differentiation and Susceptibility to Oxidative Stress-Dependent Cell Damage: Involvement in Oxotremorine-M-Mediated Neuroprotective Effects. *Antioxidants* (Basel) 12. <https://doi.org/10.3390/antiox12030687>
- Scordino, M., Urone, G., Frinchi, M., Valenza, C., Bonura, A., Cipollina, C., Ciriminna, R., Meneguzzo, F., Pagliaro, M., Mudò, G., Di Liberto, V., 2024. Anti-Apoptotic and Anti-Inflammatory Properties of Grapefruit IntegroPectin on Human Microglial HMC3 Cell Line. *Cells* 13. <https://doi.org/10.3390/cells13040355>
- Seth, T., Asija, S., Umar, S., Gupta, R., 2024. The intricate role of lipids in orchestrating plant defense responses. *Plant Sci.* 338, 111904. <https://doi.org/10.1016/j.plantsci.2023.111904>
- Somero, G.N., 2020. The cellular stress response and temperature: Function, regulation, and evolution. *J. Exp. Zool. A Ecol. Integr. Physiol.* 333, 379-397. <https://doi.org/10.1002/jez.2344>
- Song, S., Tan, J., Miao, Y., Li, M., Zhang, Q., 2017. Crosstalk of autophagy and apoptosis: Involvement of the dual role of autophagy under ER stress. *J. Cell Physiol.* 232, 2977-2984. <https://doi.org/10.1002/jcp.25785>
- Thaman, J., Pal, R. S., Chaitanya, M. V. N. L., Yanadaiah, P., Thangavelu, P., Sharma, S., Amoateng, P., Arora, S., Sivasankaran, P., Pandey, P., & Mazumder, A., 2023. Reconciling the Gap between Medications and their Potential Leads: The Role of Marine Metabolites in the Discovery of New Anticancer Drugs: A Comprehensive Review. *Curr. Pharm. Des.* 29(39), 3137–3153. <https://doi.org/10.2174/0113816128272025231106071447>
- Varrella, S., Romano, G., Ianora, A., Bentley, M. G., Ruocco, N., Costantini, M., 2014. Molecular response to toxic diatom-derived aldehydes in the sea urchin *Paracentrotus lividus*. *Mar. drugs* 12(4), 2089-2113. <https://doi.org/10.3390/md12042089>
- Varrella, S., Romano, G., Ruocco, N., Ianora, A., Bentley, M. G., Costantini, M., 2016. First morphological and molecular evidence of the negative impact of diatom-derived hydroxyacids on the sea urchin *Paracentrotus lividus*. *Tox. Sci.* 151(2), 419-433. <https://doi.org/10.1093/toxsci/kfw053>
- Wangenstein, O.S., Turon, X., Casso, M., Palacín, C., 2013. The reproductive cycle of the sea urchin *Arbacia lixula* in northwest Mediterranean: potential influence of temperature and photoperiod. *Mar. Biol.* 160, 3157–3168. <https://doi.org/10.1007/s00227-013-2303-8>
- Zhou, C., Sun, H., Zheng, C., Gao, J., Fu, Q., Hu, N., Shao, X., Zhou, Y., Xiong, J., Nie, K., Zhou, H., Shen, L., Fang, H., Lyu, J., 2018. Oncogenic HSP 60 regulates mitochondrial oxidative phosphorylation to support Erk1/2 activation during pancreatic cancer cell growth. *Cell Death Dis.* 9(2), 161. <https://doi.org/10.1038/s41419-017-0196-z>