

# NONLINEAR VECTOR DUFFING INCLUSIONS WITH NO GROWTH RESTRICTION ON THE ORIENTOR FIELD

NIKOLAOS S. PAPAGEORGIU, CALOGERO VETRO, FRANCESCA VETRO

ABSTRACT. We consider nonlinear multivalued Dirichlet Duffing systems. We do not impose any growth condition on the multivalued perturbation. Using tools from the theory of nonlinear operators of monotone type, we prove existence theorems for the convex and the nonconvex problems. Also we show the existence of extremal trajectories and show that such solutions are  $C_0^1(T, \mathbb{R}^N)$ -dense in the solution set of the convex problem (strong relaxation theorem).

## 1. INTRODUCTION

In this paper, we continue our work on multivalued nonlinear Duffing systems initiated in Papageorgiou-Vetro-Vetro [14]. So, the system under consideration is the following:

$$(1) \quad \begin{cases} -a(u'(t))' - r(t)|u'(t)|^{p-2}u'(t) \in F(t, u(t)) & \text{for a.a. } t \in T = [0, b], \\ u(0) = u(b) = 0, & 1 < p < +\infty. \end{cases}$$

Here  $a : \mathbb{R}^N \rightarrow \mathbb{R}^N$  is a monotone homeomorphism and incorporates as special cases many differential operators of interest. In [14] we proved existence theorems for both the convex and nonconvex problems (that is,  $F$  is convex valued and respectively nonconvex valued). Also, we proved a relaxation theorem showing that the solutions of the nonconvex problem are  $C_0^1(T, \mathbb{R}^N)$ -dense in the solution set of the convex problem. In [14] the hypotheses on the multivalued perturbation  $F(t, x)$  (the orientor field), dictated a sublinear growth for  $F(t, \cdot)$ . In contrast here we do not impose any global growth condition on  $F(t, \cdot)$ . Instead, we employ a Hartman-type condition on  $F(t, \cdot)$ . As in [14], we prove existence theorems for the convex and nonconvex problems. We also show the existence of “extremal solutions”, that is, solutions of (1) when  $F(t, x)$  is replaced by  $\text{ext}F(t, x)$  (= the extreme points of  $F(t, x)$ ). In the context of control systems such solutions correspond to states generated by “bang-bang controls”. Finally we prove a “strong relaxation theorem” showing that the extremal solutions are  $C_0^1(T, \mathbb{R}^N)$ -dense in the solution set of the convex problem. The last two results were mentioned as open problems in [14].

The presence of the drift term  $r(\cdot)|u'(\cdot)|^{p-2}u'(\cdot)$ , characterizes problem (1) as non-variational. So, our method of proof is topological but it is different from the one in [14]. There, the main tool was fixed point theory. Here our arguments are based on the theory of nonlinear operators of monotone type.

We mention that the starting point for the work in [14], was the recent paper of Kalita-Kowalski [10], where the authors studied scalar semilinear Duffing inclusions

---

*Key words and phrases.* Duffing system, extremal solutions, nonlinear operator of monotone type, strong relaxations.

*2010 Mathematics Subject Classification:* 34A60, 34B15.

(the convex problem only). Earlier results for single-valued such equations, can be found in Galewski [2], Kowalski [11], Tomiczek [15].

## 2. MATHEMATICAL BACKGROUND - HYPOTHESES

Let  $X$  be a reflexive Banach space and  $X^*$  its topological dual. By  $\langle \cdot, \cdot \rangle$  we denote the duality brackets for the pair  $(X^*, X)$ . A multivalued map  $A : D \subseteq X \rightarrow 2^{X^*}$  is said to be “monotone”, if

$$\langle u^* - x^*, u - x \rangle \geq 0 \quad \text{for all } (u, u^*), (x, x^*) \in \text{Gr}A.$$

The map  $A(\cdot)$  is “strictly monotone”, if it is monotone and

$$\langle u^* - x^*, u - x \rangle = 0 \quad \implies \quad u = x.$$

We say that  $A(\cdot)$  is “maximal monotone”, if it is monotone and

$$\langle u^* - x^*, u - x \rangle \geq 0 \quad \text{for all } (u, u^*) \in \text{Gr}A \quad \implies \quad (x, x^*) \in \text{Gr}A.$$

This condition is equivalent to saying that  $\text{Gr}A$  is maximal with respect to inclusion among the graphs of all monotone maps.

A nonlinear operator  $K : X \rightarrow X^*$  is said to be of type  $(S)_+$ , if the following property holds

$$u_n \xrightarrow{w} u \text{ in } X \text{ and } \limsup_{n \rightarrow +\infty} \langle K(u_n), u_n - u \rangle \leq 0 \quad \implies \quad u_n \rightarrow u \text{ in } X.$$

A multivalued map  $V : X \rightarrow 2^{X^*}$  is said to be “pseudomonotone”, if

- (a) for every  $x \in X$ ,  $V(x) \subseteq X^*$  is nonempty, convex and  $w$ -compact;
- (b) for any sequences  $\{u_n\}_{n \geq 1} \subseteq X$ ,  $\{u_n^*\}_{n \geq 1} \subseteq X^*$  such that

$$u_n \xrightarrow{w} u \text{ in } X, u_n^* \xrightarrow{w} u^* \text{ in } X^*, u_n^* \in V(u_n) \text{ for all } n \in \mathbb{N},$$

$$\limsup_{n \rightarrow +\infty} \langle u_n^*, u_n - u \rangle \leq 0,$$

we have  $u^* \in V(u)$  and  $\langle u_n^*, u_n \rangle \rightarrow \langle u^*, u \rangle$ .

Pseudomonotone maps exhibit remarkable surjectivity properties. Recall that a multivalued map  $V : X \rightarrow 2^{X^*} \setminus \{\emptyset\}$  is said to be “coercive”, if

$$\frac{\inf[\langle u^*, u \rangle : u^* \in V(u)]}{\|u\|} \rightarrow +\infty \text{ as } \|u\| \rightarrow +\infty.$$

We have the following surjectivity result for pseudomonotone maps (see Gasiński-Papageorgiou [3], Theorem 3.2.52, p. 336), see also Francù[1].

**Theorem 1.** *If  $V : X \rightarrow 2^{X^*}$  is pseudomonotone and coercive, then  $V(\cdot)$  is surjective.*

Let  $Y$  be a separable Banach space. We introduce the following families of subsets of  $Y$ :

$$P_{f(c)}(Y) = \{A \subseteq Y : A \text{ is nonempty, closed (and convex)}\},$$

$$P_{(w)k(c)}(Y) = \{A \subseteq Y : A \text{ is nonempty, (w-) compact (and convex)}\}.$$

Let  $(\Omega, \Sigma)$  be a measurable space and  $F : \Omega \rightarrow 2^Y \setminus \{\emptyset\}$ . We say that  $F$  is “graph measurable” if  $\text{Gr}F = \{(\omega, y) \in \Omega \times Y : y \in F(\omega)\} \in \Sigma \otimes B(Y)$  with  $B(Y)$  being the Borel  $\sigma$ -field of  $Y$ . Suppose that  $\mu(\cdot)$  is a finite measure defined on  $\Sigma$ . By the Yankov-von Neumann-Aumann selection theorem (see Hu-Papageorgiou [8], Theorem 2.14, p. 158), a graph measurable multifunction  $F : \Omega \rightarrow 2^Y \setminus \{\emptyset\}$  admits a measurable

selection, that is, there exists a function  $f : \Omega \rightarrow Y$  which is  $(\Sigma, B(Y))$ -measurable and  $f(w) \in F(w)$   $\mu$ -a.e. Given a graph measurable multifunction  $F : \Omega \rightarrow 2^Y \setminus \{\emptyset\}$  and  $1 \leq p \leq +\infty$  we set

$$S_F^p = \{f \in L^p(\Omega, Y) : f(\omega) \in F(\omega) \text{ } \mu\text{-a.e. on } \Omega\}.$$

It is easy to check that

$$S_F^p \neq \emptyset \text{ if and only if } \inf[\|y\| : y \in F(\omega)] \leq \psi(\omega) \text{ } \mu\text{-a.e., with } \psi \in L^p(\Omega).$$

The set  $S_F^p$  is “decomposable”, that is,

$$(A, f_1, f_2) \in \Sigma \times S_F^p \times S_F^p \implies \chi_A f_1 + \chi_{\Omega \setminus A} f_2 \in S_F^p.$$

Recall that for  $C \subseteq \Omega$ ,  $\chi_C$  is the characteristic function of  $C$  defined by

$$\chi_C(\omega) = \begin{cases} 1 & \text{if } \omega \in C \\ 0 & \text{if } \omega \notin C \end{cases}.$$

On  $P_f(Y)$  we can define an extended metric, known as the “Hausdorff metric”, by setting

$$h(C, K) = \max\{\max_{c \in C} d(c, K), \max_{k \in K} d(k, C)\} \text{ for all } C, K \in P_f(Y).$$

We know that  $(P_f(Y), h)$  is a complete metric space. We say that a multifunction  $F : Y \rightarrow P_f(Y)$  is “ $h$ -continuous”, if it is continuous from  $Y$  into  $(P_f(Y), h)$ .

Suppose that  $E, Z$  are Hausdorff topological spaces and  $G : E \rightarrow 2^Z \setminus \{\emptyset\}$  a multifunction. We say that  $G(\cdot)$  is “upper semicontinuous (usc)” (resp. “lower semicontinuous (lsc)”), if for all  $C \subseteq Z$  closed,  $G^-(C) = \{e \in E : G(e) \cap C \neq \emptyset\}$  (resp.  $G^+(C) = \{e \in E : G(e) \subseteq C\}$ ) is closed in  $E$ .

For any  $A \subseteq \mathbb{R}^N$ , we set  $|A| = \sup\{|a| : a \in A\}$  (hereafter  $|\cdot|$  denotes the Euclidean norm on  $\mathbb{R}^N$ ). Let  $T = [0, b]$ . By  $L_w^1(T, \mathbb{R}^N)$  we denote the Lebesgue space  $L^1(T, \mathbb{R}^N)$  equipped with the weak norm  $\|\cdot\|_w$  defined by

$$\|u\|_w = \sup \left[ \left| \int_s^t u(\tau) d\tau \right| : 0 \leq s \leq t \leq b \right], \quad u \in L^1(T, \mathbb{R}^N).$$

Equivalently, we can define the weak norm of  $u$  by

$$\|u\|_w = \sup \left[ \left| \int_0^t u(\tau) d\tau \right| : 0 \leq t \leq b \right].$$

On the other hand by  $L^\theta(T, \mathbb{R}^N)_w$  ( $1 \leq \theta < +\infty$ ), we denote the space  $L^\theta(T, \mathbb{R}^N)$  furnished with the weak topology. The following simple fact can be found in Papageorgiou [9], Lemma 2.8, p. 24.

**Proposition 1.** *If  $\{u_n, u\}_{n \geq 1} \subseteq L^p(T, \mathbb{R}^N)$ ,  $1 < p < +\infty$ ,  $u_n \xrightarrow{\|\cdot\|_w} u$  and  $\sup_{n \geq 1} \|u_n\|_p < +\infty$ , then  $u_n \xrightarrow{w} u$  in  $L^p(T, \mathbb{R}^N)$ .*

Consider the following nonlinear vector eigenvalue problem:

$$-(|u'(t)|^{p-2} u'(t))' = \widehat{\lambda} |u(t)|^{p-2} u(t) \text{ a.e. on } T, \quad u(0) = u(b) = 0.$$

This problem has a smallest eigenvalue  $\widehat{\lambda}_1 > 0$ , which admits the following variational characterization

$$(2) \quad \widehat{\lambda}_1 = \inf \left[ \frac{\|u'\|_p^p}{\|u\|_p^p} : u \in W_0^{1,p}((0, b), \mathbb{R}^N), u \neq 0 \right]$$

(see Gasiński-Papageorgiou [3], p. 768).

Now we introduce the hypotheses on the map  $a : \mathbb{R}^N \rightarrow \mathbb{R}^N$  and on the drift coefficient  $r(\cdot)$ .

$H(a)$ :  $a : \mathbb{R}^N \rightarrow \mathbb{R}^N$  is continuous, monotone,  $a(0) = 0$  and

$$a(y) = \widehat{c}(|y|)y \quad \text{for all } y \in \mathbb{R}^N,$$

with  $\widehat{c} : (0, +\infty) \rightarrow (0, +\infty)$  continuous,  $c_0 t^p \leq \widehat{c}(t)t^2$  for all  $t > 0$ , some  $c_0 > 0$  and

$$|a(y)| \leq c_1[1 + |y|^{p-1}] \quad \text{for some } c_1 > 0, \text{ all } y \in \mathbb{R}^N.$$

*Remark 1.* Evidently  $a : \mathbb{R}^N \rightarrow \mathbb{R}^N$  is maximal monotone and

$$c_0|y|^p \leq (a(y), y)_{\mathbb{R}^N} \quad \text{for all } y \in \mathbb{R}^N.$$

Hypotheses  $H(a)$  are more restrictive than those in [14], where no growth restriction was imposed on  $a(\cdot)$ . Nevertheless, as the examples which follow illustrate, conditions  $H(a)$  are still very general and incorporate many differential operators of interest.

**Example 1.** The following maps  $a : \mathbb{R}^N \rightarrow \mathbb{R}^N$  satisfy hypotheses  $H(a)$ :

$$a(y) = |y|^{p-2}y \quad \text{for all } y \in \mathbb{R}^N, 1 < p < +\infty$$

(this map corresponds to the vector  $p$ -Laplacian),

$$a(y) = |y|^{p-2}y + |y|^{q-2}y \quad \text{for all } y \in \mathbb{R}^N, 1 < q < p < +\infty$$

(this map corresponds to the vector  $(p, q)$ -Laplacian),

$$a(y) = (1 + |y|^2)^{\frac{p-2}{2}}y \quad \text{for all } y \in \mathbb{R}^N, 1 < p < +\infty.$$

$H(r)$ :  $r \in L^\infty(T)$  ( $T = [0, b]$ ),  $r(t) \geq 0$  for a.a.  $t \in T$  and  $\|r\|_\infty < c_0 \widehat{\lambda}^{1/p}$ .

Finally in what follows by  $\|\cdot\|$  we denote the norm of the Sobolev space  $W_0^{1,p}((0, b), \mathbb{R}^N)$ . On account of the Poincaré inequality, we have

$$\|u\| = \|u'\|_p \quad \text{for all } u \in W_0^{1,p}((0, b), \mathbb{R}^N).$$

In what follows to simplify our notation, we write

$$W_0^{1,p} = W_0^{1,p}((0, b), \mathbb{R}^N).$$

Also we write

$$C_0(T, \mathbb{R}^N) = \{u \in C(T, \mathbb{R}^N) : u(0) = u(b) = 0\}$$

and

$$C_0^1(T, \mathbb{R}^N) = C^1(T, \mathbb{R}^N) \cap C_0(T, \mathbb{R}^N).$$

Finally if  $1 \leq p < +\infty$ , then  $1 < p' \leq +\infty$  is defined by  $\frac{1}{p} + \frac{1}{p'} = 1$ .

### 3. THE CONVEX PROBLEM

In this section we prove an existence theorem for the convex problem (that is,  $F$  is convex valued). The precise hypotheses on  $F(t, x)$  are the following:

$H(F)_1$ :  $F : T \times \mathbb{R}^N \rightarrow P_{kc}(\mathbb{R}^N)$  is a multifunction such that

- (i) for all  $x \in \mathbb{R}^N$ , the multifunction  $t \rightarrow F(t, x)$  admits a measurable selection;
- (ii) for a.a.  $t \in T$ ,  $\text{Gr}F(t, \cdot) \subseteq \mathbb{R}^N \times \mathbb{R}^N$  is closed;

(iii) for every  $\eta > 0$ , there exists  $a_\eta \in L^{p'}(T)_+$  such that

$$|F(t, x)| \leq a_\eta(t) \quad \text{for a.a. } t \in T, \text{ all } |x| \leq \eta$$

and there exists  $M > 0$  such that

$$(h, x)_{\mathbb{R}^N} < 0 \quad \text{for a.a. } t \in T, \text{ all } |x| \geq M, \text{ all } h \in F(t, x).$$

*Remark 2.* Hypothesis  $H(F)_1(i)$  is satisfied if for all  $x \in \mathbb{R}^N$  the multifunction  $t \rightarrow F(t, x)$  is graph measurable. Hypotheses  $H(F)_1(ii)$ , (iii) imply that for a.a.  $t \in T$ ,  $F(t, \cdot)$  is usc (see Hu-Papageorgiou [8], Proposition 2.23, p. 43). The second part of hypothesis  $H(F)_1(iii)$  is a Hartman-type condition on  $F(t, \cdot)$  (see Hartman [6, 7]). In the context of scalar, single-valued Duffing equations, this hypothesis is in fact a sign condition saying that  $F(t, x) < 0$  for  $x \geq M$  and  $F(t, x) > 0$  for  $x \leq -M$ . We stress that no global growth condition is imposed on  $F(t, \cdot)$ . This distinguishes our work here from that in [14].

Consider the nonlinear operator  $A : W_0^{1,p} \rightarrow W^{-1,p'} = (W_0^{1,p})^*$  defined by

$$\langle A(u), y \rangle = \int_0^b (a(u'), y')_{\mathbb{R}^N} dt \quad \text{for all } u, y \in W_0^{1,p}.$$

From Gasiński-Papageorgiou [5] (Problem 2.192, p. 279), we have:

**Proposition 2.** *If hypotheses  $H(a)$  hold, then  $A(\cdot)$  is continuous, monotone (hence maximal monotone too) and of type  $(S)_+$ .*

Then let  $V : W_0^{1,p} \rightarrow W^{-1,p'}$  be defined by

$$\langle V(u), y \rangle = \langle A(u), y \rangle - \int_0^b r(t) |u'|^{p-2} (u', y)_{\mathbb{R}^N} dt \quad \text{for all } u, y \in W_0^{1,p},$$

that is,

$$(3) \quad V(u) = A(u) - r(\cdot) |u'(\cdot)|^{p-2} u'(\cdot) \quad \text{for all } u \in W_0^{1,p}.$$

**Proposition 3.** *If hypotheses  $H(a)$ ,  $H(r)$  hold, then  $V : W_0^{1,p} \rightarrow W^{-1,p'}$  is pseudomonotone and coercive.*

*Proof.* Suppose that

$$(4) \quad u_n \xrightarrow{w} u \text{ in } W_0^{1,p}, V(u_n) \xrightarrow{w} u^* \text{ in } W^{-1,p'}, \limsup_{n \rightarrow +\infty} \langle V(u_n), u_n - u \rangle \leq 0.$$

From (4) and since  $W_0^{1,p}$  is embedded compactly in  $C_0(T, \mathbb{R}^N)$ , we have

$$\begin{aligned} & u_n \rightarrow u \text{ in } C_0(T, \mathbb{R}^N), \\ & \Rightarrow \int_0^b r(t) |u'_n|^{p-2} (u'_n, u_n - u)_{\mathbb{R}^N} dt \rightarrow 0, \\ & \Rightarrow \limsup_{n \rightarrow +\infty} \langle A(u_n), u_n - u \rangle \leq 0 \quad (\text{see (3), (4)}), \\ (5) \quad & \Rightarrow u_n \rightarrow u \text{ in } W_0^{1,p} \quad (\text{see Proposition 2}). \end{aligned}$$

From (5) it follows that

$$u'_n \rightarrow u' \text{ in } L^p(T, \mathbb{R}^N).$$

So, by passing to a subsequence if necessary, we may assume that

$$(6) \quad u'_n(t) \rightarrow u'(t) \text{ for a.a. } t \in T, |u'_n(t)|, |u'(t)| \leq \xi(t) \text{ for a.a. } t \in T, \text{ all } n \in \mathbb{N},$$

with  $\xi \in L^p(T)_+$ . Then from (6) and the dominated convergence theorem, we have

$$(7) \quad \begin{aligned} & \int_0^b \left| |u'_n|^{p-2}u'_n - |u'|^{p-2}u' \right|^{p'} dt \rightarrow 0 \\ \Rightarrow & r|u'_n|^{p-2}u'_n \rightarrow r|u'|^{p-2}u' \text{ in } L^{p'}(T, \mathbb{R}^N) \text{ (see } H(r)). \end{aligned}$$

On account of (5) and Proposition 2, we have

$$(8) \quad A(u_n) \rightarrow A(u) \text{ in } W^{-1,p'}.$$

From (7) and (8) it follows that for the original sequence we have

$$\begin{aligned} & \langle V(u_n), u_n \rangle \rightarrow \langle V(u), u \rangle, \quad V(u) = u^*, \\ \Rightarrow & V(\cdot) \text{ is pseudomonotone.} \end{aligned}$$

Also for all  $u \in W_0^{1,p}$  we have

$$(9) \quad \begin{aligned} \langle V(u), u \rangle &= \int_0^b (a(u'), u')_{\mathbb{R}^N} dt - \int_0^b r(t) |u'|^{p-2} (u', u)_{\mathbb{R}^N} dt \\ &\geq c_0 \|u'\|_p^p - \|r\|_\infty \|u'\|_p^{p-1} \|u\|_p \\ &\text{(see hypothesis } H(a) \text{ and use Hölder's inequality)} \\ &\geq \left[ c_0 - \frac{\|r\|_\infty}{\widehat{\lambda}_1^{1/p}} \right] \|u'\|_p^p \quad \text{(see (2)),} \\ \Rightarrow & V(\cdot) \text{ is coercive (see } H(r)). \end{aligned}$$

□

Let  $N : W_0^{1,p} \rightarrow 2^{L^{p'}(T, \mathbb{R}^N)}$  be the multifunction defined by

$$N(u) = S_{F(\cdot, u(\cdot))}^{p'} \quad \text{for all } u \in W_0^{1,p}.$$

From [14, Proposition 3], we have:

**Proposition 4.** *If hypotheses  $H(r)$ ,  $H(F)_1$  hold, then the multifunction  $N(\cdot)$  has values in  $P_{wkc}(L^{p'}(T, \mathbb{R}^N))$  and it is usc from  $W_0^{1,p}$  into  $L^{p'}(T, \mathbb{R}^N)_w$ .*

Recall that  $W_0^{1,p} \hookrightarrow L^p(T, \mathbb{R}^N)$  compactly. Hence

$$(10) \quad \begin{aligned} & L^{p'}(T, \mathbb{R}^N) \hookrightarrow W^{-1,p'} \text{ compactly} \\ & \text{(see Gasiński-Papageorgiou [3], Lemma 2.2.27, p. 141),} \\ \Rightarrow & N : W_0^{1,p} \rightarrow P_{kc}(W^{-1,p'}) \text{ is compact} \\ & \text{(that is, } N(\cdot) \text{ maps bounded sets to relatively compact ones),} \\ \Rightarrow & u \rightarrow V(u) - N(u) \text{ is pseudomonotone} \\ & \text{(see Gasiński-Papageorgiou [3], Proposition 3.2.51, p. 334).} \end{aligned}$$

Also for all  $u \in W_0^{1,p}$  and all  $f \in S_{F(\cdot, u(\cdot))}^{p'}$  we have

$$\begin{aligned} & \langle V(u), u \rangle - \int_0^b (f, u)_{\mathbb{R}^N} dt \\ & \geq \left[ c_0 - \frac{\|r\|_\infty}{\widehat{\lambda}_1^{1/p}} \right] \|u'\|_p^p - \int_{\{|u| \geq M\}} (f, u)_{\mathbb{R}^N} dt - \int_{\{|u| < M\}} (f, u)_{\mathbb{R}^N} dt \quad (\text{see (9)}), \\ & \geq \left[ c_0 - \frac{\|r\|_\infty}{\widehat{\lambda}_1^{1/p}} \right] \|u'\|_p^p - c_2 \quad \text{for some } c_2 > 0 \quad (\text{see hypothesis } H(F)_1(\text{iii})), \end{aligned}$$

(11)  $\Rightarrow u \rightarrow V(u) - N(u)$  is coercive.

**Theorem 2.** *If hypotheses  $H(a)$ ,  $H(r)$ ,  $H(F)_1$  hold, then problem (1) has a solution  $u_0 \in C_0^1(T, \mathbb{R}^N)$ .*

*Proof.* On account of (10), (11) and Theorem 1, we can find  $u_0 \in W_0^{1,p}$  such that

$$\begin{aligned} & 0 \in V(u_0) - N(u_0), \\ & \Rightarrow V(u_0) = f \quad \text{for some } f \in N(u_0) = S_{F(\cdot, u_0(\cdot))}^{p'}, \\ & \Rightarrow \langle A(u_0), y \rangle - \int_0^b r(t)|u_0'|^{p-2}(u_0', y)_{\mathbb{R}^N} dt = \int_0^b (f, y)_{\mathbb{R}^N} dt \quad \text{for all } y \in W_0^{1,p}, \\ (12) \Rightarrow & -a(u_0'(t))' - r(t)|u_0'(t)|^{p-2}u_0'(t) = f(t) \quad \text{for a.a. } t \in T, \quad u_0(0) = u_0(b) = 0. \end{aligned}$$

From (12) as in the proof of [14, Proposition 2], we conclude that  $u_0 \in C_0^1(T, \mathbb{R}^N)$ .  $\square$

*Remark 3.* An interesting consequence of the above proof, is that if by  $\widehat{S}_c \subseteq C_0^1(T, \mathbb{R}^N)$  we denote the solution set of the convex problem, then  $\widehat{S}_c \in P_k(C_0^1(T, \mathbb{R}^N))$  (see also [14]).

#### 4. THE NONCONVEX PROBLEM

In this section we prove an existence theorem for the nonconvex problem (that is,  $F$  has nonconvex values). The precise hypotheses on the orientor field  $F(t, x)$  are the following:

$H(F)_2$ :  $F : T \times \mathbb{R}^N \rightarrow P_f(\mathbb{R}^N)$  is a multifunction such that

- (i)  $(t, x) \rightarrow F(t, x)$  is graph measurable;
- (ii) for a.a.  $t \in T$ ,  $x \rightarrow F(t, x)$  is lsc;
- (iii) the same as hypothesis  $H(F)_1(\text{iii})$ .

**Theorem 3.** *If hypotheses  $H(a)$ ,  $H(r)$ ,  $H(F)_2$  hold, then problem (1) has a solution  $u_0 \in C_0^1(T, \mathbb{R}^N)$ .*

*Proof.* Consider the multifunction  $N : W_0^{1,p} \rightarrow P_f(L^{p'}(T, \mathbb{R}^N))$  defined by

$$N(u) = S_{F(\cdot, u(\cdot))}^{p'} \quad \text{for all } u \in W_0^{1,p}.$$

According to Theorem 7.27, p. 237, of Hu-Papageorgiou [8],  $N(\cdot)$  is lsc and of course it has decomposable values. So, we can apply Theorem 8.7, p. 245, of Hu-Papageorgiou [8] and produce a continuous map  $g : W_0^{1,p} \rightarrow L^{p'}(T, \mathbb{R}^N)$  such that

$$g(u) \in N(u) \quad \text{for all } u \in W_0^{1,p}.$$

We consider the following nonlinear Duffing system

$$\begin{cases} -a(u'(t))' - r(t)|u'(t)|^{p-2}u'(t) = g(u)(t) & \text{for a.a. } t \in T, \\ u(0) = u(b) = 0. \end{cases}$$

Then this problem has a solution  $u_0 \in C_0^1(T, \mathbb{R}^N)$  (see Theorem 2). Evidently  $u_0 \in C_0^1(T, \mathbb{R}^N)$  is a solution of (1).  $\square$

## 5. EXTREMAL SOLUTIONS

Let  $\text{ext}F(t, x)$  denote the set of extreme points of  $F(t, x)$ . In this section we deal with the following nonlinear multivalued Duffing system:

$$(13) \quad \begin{cases} -a(u'(t))' - r(t)|u'(t)|^{p-2}u'(t) \in \text{ext}F(t, u(t)) & \text{for a.a. } t \in T, \\ u(0) = u(b) = 0. \end{cases}$$

The solutions of (13) are of course solutions of (1) and are known as ‘‘extremal solutions’’.

To produce extremal solutions, first we prove an a priori pointwise bound for the solutions of (1), using hypothesis  $H(F)_1(\text{iii})$ .

**Proposition 5.** *If hypotheses  $H(a)$ ,  $H(r)$ ,  $H(F)_1(\text{iii})$  hold and  $u \in C_0^1(T, \mathbb{R}^N)$  is a solution of problem (1), then  $\|u\|_\infty \leq M$ .*

*Proof.* Let  $\theta(t) = \frac{1}{2}|u(t)|^2$  for all  $t \in T$ . Suppose that the proposition is not true. Then

$$\theta(t_0) = \max[\theta(t) : t \in T] > \frac{1}{2}M^2.$$

Since  $\theta(0) = \theta(b) = 0$ , we see that we may assume that  $t_0 \in (0, b)$ . Then we can find  $\delta > 0$  small such that

$$(14) \quad |u(t)| \geq M \text{ and } \frac{d}{dt}|u(t)|^2 \leq 0 \text{ for all } t \in [t_0, t_0 + \delta] \subseteq T.$$

We have

$$-a(u'(t))' - r(t)|u'(t)|^{p-2}u'(t) = f(t) \text{ for a.a. } t \in T, u(0) = u(b) = 0,$$

with  $f \in S_{F(\cdot, u(\cdot))}^{p'}$ .

We take inner product with  $u(t)$ , integrate over  $[t_0, t]$  with  $t \leq t_0 + \delta$  and perform integration by parts. We obtain

$$(15) \quad \begin{aligned} & (a(u'(t)), u(t))_{\mathbb{R}^N} - (a(u'(t_0)), u(t_0))_{\mathbb{R}^N} - \int_{t_0}^t (a(u'), u')_{\mathbb{R}^N} ds \\ & + \int_{t_0}^t r(s)|u'|^{p-2}(u', u)_{\mathbb{R}^N} ds = \int_{t_0}^t (-f, u)_{\mathbb{R}^N} ds. \end{aligned}$$

Note that

$$\begin{aligned} & \frac{d}{dt}\theta(t) \Big|_{t=t_0} = 0, \\ \Rightarrow & \frac{1}{2} \frac{d}{dt}|u(t)|^2 \Big|_{t=t_0} = 0, \\ \Rightarrow & (u'(t_0), u(t_0))_{\mathbb{R}^N} = 0. \end{aligned}$$



Therefore we have

$$(16) \quad (a(u'(t_0)), u(t_0))_{\mathbb{R}^N} = \widehat{c}(|u'(t_0)|)(u'(t_0), u(t_0))_{\mathbb{R}^N} = 0 \quad (\text{see hypothesis } H(a)).$$

Also we have

$$(17) \quad \int_{t_0}^t r(s)|u'|^{p-2}(u', u)_{\mathbb{R}^N} ds = \frac{1}{2} \int_{t_0}^t r(s)|u'|^{p-2} \frac{d}{ds} |u|^2 ds \leq 0$$

(see hypothesis  $H(r)$  and (14)).

Finally on account of (14) and hypothesis  $H(F)_1(\text{iii})$ , we have

$$(18) \quad \int_{t_0}^t (-f, u)_{\mathbb{R}^N} ds > 0.$$

Returning to (15) and using (16), (17), (18), we obtain

$$\begin{aligned} & (a(u'(t)), u(t))_{\mathbb{R}^N} - \int_{t_0}^t (a(u'), u')_{\mathbb{R}^N} dt > 0, \\ \Rightarrow & (a(u'(t)), u(t))_{\mathbb{R}^N} > 0 \quad \text{for all } t \in (t_0, t_0 + \delta] \quad (\text{see hypothesis } H(a)), \\ \Rightarrow & \widehat{c}(|u'(t)|)(u'(t), u(t))_{\mathbb{R}^N} > 0 \quad \text{for all } t \in (t_0, t_0 + \delta], \\ \Rightarrow & (u'(t), u(t))_{\mathbb{R}^N} = \frac{1}{2} \frac{d}{dt} |u(t)|^2 > 0 \quad \text{for all } t \in (t_0, t_0 + \delta], \end{aligned}$$

a contradiction (see (14)). Therefore

$$\begin{aligned} \theta(t_0) &= \frac{1}{2} |u(t_0)|^2 \leq \frac{1}{2} M^2, \\ \Rightarrow |u(t)| &\leq M \quad \text{for all } t \in T. \end{aligned}$$

□

Let  $p_M : \mathbb{R}^N \rightarrow \mathbb{R}^N$  be the  $M$ -radial retraction defined by

$$p_M(x) = \begin{cases} x & \text{if } |x| \leq M \\ \frac{Mx}{|x|} & \text{if } M < |x| \end{cases} \quad \text{for all } x \in \mathbb{R}^N.$$

Evidently  $p_M(\cdot)$  is nonexpansive, that is,

$$|p_M(x) - p_M(v)| \leq |x - v| \quad \text{for all } x, v \in \mathbb{R}^N.$$

On account of Proposition 5, in (13) we may replace  $F$  by  $F_0$  defined by

$$F_0(t, x) = F(t, p_M(x)) \quad \text{for all } t \in T, \text{ all } x \in \mathbb{R}^N.$$

Note that  $F_0(t, x)$  satisfies the same conditions as  $F(t, x)$  and in addition

$$|F_0(t, x)| \leq a_M(t) \quad \text{for a.a. } t \in T, \text{ all } x \in \mathbb{R}^N, \text{ with } a_M \in L^{p'}(T)_+.$$

Therefore without any loss of generality we may assume that

$$|F(t, x)| \leq a_M(t) \quad \text{for a.a. } t \in T, \text{ all } x \in \mathbb{R}^N.$$

Let  $\mathcal{D} = \{h \in L^{p'}(T, \mathbb{R}^N) : |h(t)| \leq a_M(t) \text{ for a.a. } t \in T\}$ . We consider the following Duffing system

$$(19) \quad -a(u'(t))' - r(t)|u'(t)|^{p-2}u'(t) = h(t) \text{ for a.a. } t \in T, \quad u(0) = u(b) = 0.$$

Let  $C_0 = \{u \in C_0^1(T, \mathbb{R}^N) : u \text{ is a solution of (19) with } h \in \mathcal{D}\}$ .

**Proposition 6.** *If hypotheses  $H(a)$ ,  $H(r)$  hold, then  $C_0 \subseteq C_0^1(T, \mathbb{R}^N)$  is compact.*

*Proof.* Let  $\{u_n\}_{n \geq 1} \subseteq C_0$ . We have

$$(20) \quad -a(u'_n)' - r(t)|u'_n|^{p-2}u'_n = h_n, \quad u_n(0) = u_n(b) = 0 \quad \text{for all } n \in \mathbb{N}.$$

We act with  $u_n$  in (20) and after integration by parts, we obtain

$$\begin{aligned} c_0 \|u'_n\|_p^p - \|r\|_\infty \|u'_n\|_p^{p-1} \|u_n\|_p &\leq \int_0^b (h_n, u_n)_{\mathbb{R}^N} dt \leq c_3 \|u'_n\|_p \quad \text{for some } c_3 > 0, \\ \Rightarrow \left[ c_0 - \frac{\|r\|_\infty}{\lambda_1^{1/p}} \right] \|u'_n\|_p^{p-1} &\leq c_3 \quad \text{for all } n \in \mathbb{N}, \end{aligned}$$

$$\Rightarrow \{u_n\}_{n \geq 1} \subseteq W_0^{1,p} \text{ is bounded (see hypothesis } H(r)),$$

$$(21) \quad \Rightarrow \{u_n\}_{n \geq 1} \subseteq C_0(T, \mathbb{R}^N) \text{ is relatively compact.}$$

From (20) we have

$$(22) \quad a(u'_n(t)) = a(u'_n(0)) - \int_0^t r(s)|u'_n|^{p-2}u'_n ds - \int_0^t h_n ds \quad \text{for all } t \in T, \text{ all } n \in \mathbb{N},$$

$$(23) \quad \Rightarrow u'_n(t) = a^{-1} \left[ a(u'_n(0)) - \int_0^t [r(s)|u'_n|^{p-2}u'_n + h_n] ds \right] \quad \text{for all } t \in T, \text{ all } n \in \mathbb{N}.$$

Note that  $\int_0^b u'_n(t) dt = 0$ . So, from (23) and Proposition 3.1(ii) of Manásevich-Mawhin [12], we have that

$$(24) \quad \{a(u'_n(0))\}_{n \geq 1} \subseteq \mathbb{R}^N \text{ is bounded.}$$

Then from (22), (24) and Arzelà-Ascoli theorem, it follows that

$$(25) \quad \{a(u'_n(\cdot))\}_{n \geq 1} \subseteq C(T, \mathbb{R}^N) \text{ is relatively compact.}$$

Consider the map  $\widehat{a}^{-1} : C(T, \mathbb{R}^N) \rightarrow C(T, \mathbb{R}^N)$  defined by

$$\widehat{a}^{-1}(u)(\cdot) = a^{-1}(u(\cdot)) \quad \text{for all } u \in C(T, \mathbb{R}^N).$$

Evidently this map is continuous and bounded (that is, maps bounded sets to bounded sets). So, from (25) it follows that

$$(26) \quad \{u'_n\}_{n \geq 1} \subseteq C(T, \mathbb{R}^N) \text{ is relatively compact.}$$

From (21) and (26), we have that

$$\{u_n\}_{n \geq 1} \subseteq C_0^1(T, \mathbb{R}^N) \text{ is relatively compact.}$$

So, we may assume that

$$u_n \rightarrow u \text{ in } C_0^1(T, \mathbb{R}^N).$$

Evidently  $\mathcal{D} \subseteq L^{p'}(T, \mathbb{R}^N)$  is  $w$ -compact and so we may assume that

$$h_n \xrightarrow{w} h \text{ in } L^{p'}(T, \mathbb{R}^N), \quad h \in \mathcal{D}.$$

We have

$$\begin{aligned} \int_0^b (a(u'_n), y')_{\mathbb{R}^N} dt - \int_0^b r(t)|u'_n|^{p-2}(u'_n, y)_{\mathbb{R}^N} dt &= \int_0^b (h_n, y)_{\mathbb{R}^N} dt \\ &\text{for all } y \in W_0^{1,p}, \text{ all } n \in \mathbb{N}. \end{aligned}$$

Passing to the limit as  $n \rightarrow +\infty$  and using (24) and (25), we obtain

$$\int_0^b (a(u'), y')_{\mathbb{R}^N} dt - \int_0^b r(t) |u'|^{p-2} (u', y)_{\mathbb{R}^N} dt = \int_0^b (h, y)_{\mathbb{R}^N} dt$$

for all  $y \in W_0^{1,p}$ ,

$$\Rightarrow u \in C_0.$$

Therefore  $C_0 \subseteq C_0^1(T, \mathbb{R}^N)$  is compact.  $\square$

Now let  $C = \overline{\text{conv}} C_0 \in P_{kc}(C_0^1(T, \mathbb{R}^N))$  (see Gasiński-Papageorgiou [4], Theorem 5.86, p. 852). If by  $\widehat{S}_c$  we denote the solution set of the convex problem, then  $\widehat{S}_c \subseteq C$ .

To produce extremal solutions (that is, solutions of (13)), we introduce the following conditions on the orientor field  $F(t, x)$ .

$H(F)_3$ :  $F : T \times \mathbb{R}^N \rightarrow P_{kc}(\mathbb{R}^N)$  is a multifunction such that

- (i) for all  $x \in \mathbb{R}^N$ ,  $t \rightarrow F(t, x)$  is graph measurable;
- (ii) for a.a.  $t \in T$ ,  $x \rightarrow F(t, x)$  is  $h$ -continuous;
- (iii) the same as hypothesis  $H(F)_1$ (iii).

*Remark 4.* Hypotheses  $H(F)_3$ (i), (ii) imply that  $(t, x) \rightarrow F(t, x)$  is graph measurable (see Proposition 7.9, p. 229 of Hu-Papageorgiou [8]). As we already mentioned, by replacing  $F$  with  $F_0$  if necessary (see Proposition 5), without any loss of generality we may assume that

$$(27) \quad |F(t, x)| \leq a_M(t) \quad \text{for a.a. } t \in T, \text{ all } x \in \mathbb{R}^N, \text{ with } a_M \in L^{p'}(T)_+.$$

**Theorem 4.** *If hypotheses  $H(a)$ ,  $H(r)$ ,  $H(F)_3$  hold, then problem (13) has a solution  $u_0 \in C_0^1(T, \mathbb{R}^N)$ .*

*Proof.* Recall that  $C = \overline{\text{conv}} C_0 \in P_{kc}(C_0^1(T, \mathbb{R}^N))$  (see Proposition 6). We consider the multifunction  $G : C \rightarrow P_{wkc}(L^{p'}(T, \mathbb{R}^N))$  defined by

$$G(u) = S_{F(\cdot, u(\cdot))}^{p'} \quad \text{for all } u \in C.$$

Using Theorem 8.31, p. 260, of Hu-Papageorgiou [8], we can find a continuous map  $g : C \rightarrow L_w^1(T, \mathbb{R}^N)$  such that

$$(28) \quad g(u) \in \text{ext} G(u) = \text{ext} S_{F(\cdot, u(\cdot))}^{p'} = S_{\text{ext} F(\cdot, u(\cdot))}^{p'} \quad \text{for all } u \in C$$

(see Hu-Papageorgiou [8], Theorem 4.6, p. 192). On account of (27) and Proposition 1, we have that  $g : C \rightarrow L_w^{p'}(T, \mathbb{R}^N)$  is continuous. We consider the following Duffing system:

$$\begin{cases} -a(u'(t))' - r(t) |u'(t)|^{p-2} u'(t) = g(u)(t) & \text{for a.a. } t \in T, \\ u(0) = u(b) = 0. \end{cases}$$

By Theorem 2, this problem has a solution  $u_0 \in C_0^1(T, \mathbb{R}^N)$ . Evidently (28) implies that  $u_0$  is an extremal solution (that is, solves problem (13)).  $\square$

## 6. STRONG RELAXATION THEOREM

In this section, under stronger conditions on  $a(\cdot)$  and  $F(\cdot, \cdot)$  we prove a strong relaxation theorem.

Recall that if  $E \subseteq \mathbb{R}^N$  is compact, then  $\text{conv}E \in P_{kc}(\mathbb{R}^N)$  (see Gasiński-Papageorgiou [4], Problem 5.93, p. 889). We consider the following multivalued Duffing systems:

$$(29) \quad -a(u'(t))' - r(t)u(t) \in \text{ext conv}F(t, u(t)) \quad \text{for a.a. } t \in T, u(0) = u(b) = 0,$$

$$(30) \quad -a(u'(t))' - r(t)u(t) \in \text{conv}F(t, u(t)) \quad \text{for a.a. } t \in T, u(0) = u(b) = 0.$$

Let  $\widehat{S}_e \subseteq C_0^1(T, \mathbb{R}^N)$  be the solution set of (29) and  $\widehat{S}_c \subseteq C_0^1(T, \mathbb{R}^N)$  the solution set of (30). We know that  $\widehat{S}_c \in P_k(C_0^1(T, \mathbb{R}^N))$ .

Our aim is to show that

$$\overline{\widehat{S}_e}^{C_0^1(T, \mathbb{R}^N)} = \widehat{S}_c.$$

Such a density result is known as “strong relaxation theorem”. It is important in control theory in connection with the “bang-bang principle”.

To have a strong relaxation theorem, we need stronger conditions on  $a(\cdot)$  and  $F(\cdot, \cdot)$ .  $H(a)'$ :  $a : \mathbb{R}^N \rightarrow \mathbb{R}^N$  is continuous, monotone,  $a(0) = 0$ ,

$$a(y) = \widehat{c}(|y|)y \quad \text{for all } y \in \mathbb{R}^N,$$

with  $\widehat{c} : (0, +\infty) \rightarrow (0, +\infty)$  continuous,  $c_0 t^2 \leq \widehat{c}(t)t^2$  for all  $t > 0$ , some  $c_0 > 0$ , for every  $\eta > 0$ , there exists  $\widetilde{c}_\eta > 0$  such that

$$\widetilde{c}_\eta |y - v|^2 \leq (a(y) - a(v), y - v)_{\mathbb{R}^N} \quad \text{for all } |y|, |v| \leq \eta$$

and

$$|a(y)| \leq c_1(1 + |y|) \quad \text{for some } c_1 > 0, \text{ all } y \in \mathbb{R}^N.$$

*Remark 5.* Evidently  $a(\cdot)$  is strictly monotone and maximal monotone. Also  $a(\cdot)$  is a homeomorphism onto  $\mathbb{R}^N$  and  $|a^{-1}(y)| \rightarrow +\infty$  as  $|y| \rightarrow +\infty$ .

**Example 2.** The following maps satisfy hypotheses  $H(a)'$ :

$$a(y) = y \quad \text{for all } y \in \mathbb{R}^N,$$

$$a(y) = |y|^{q-2}y + y \quad \text{for all } y \in \mathbb{R}^N, \text{ with } 1 < q < 2,$$

$$a(y) = \begin{cases} 2|y|^{q-2}y & \text{if } |y| \leq 1 \\ 2y & \text{if } 1 < |y| \end{cases} \quad \text{for all } y \in \mathbb{R}^N, \text{ with } 1 < q < 2,$$

$$a(y) = (1 + |y|^q)^{\frac{2-q}{q}} |y|^{q-2}y \quad \text{for all } y \in \mathbb{R}^N, \text{ with } 1 < q \leq 2.$$

$H(F)_4$ :  $F : T \times \mathbb{R}^N \rightarrow P_k(\mathbb{R}^N)$  is a multifunction such that

- (i) for all  $x \in \mathbb{R}^N$ ,  $t \rightarrow F(t, x)$  is graph measurable;
- (ii) for every  $\eta > 0$ , there exists  $k_\eta \in L^1(T)_+$  such that

$$\widetilde{c}_\eta - \frac{\|r\|_\infty}{\widehat{\lambda}_1^{1/p}} - \|k_\eta\|_\infty b > 0,$$

and

$$h(\text{conv}F(t, x), \text{conv}F(t, v)) \leq k_\eta(t)|x - v|$$

for a.a.  $t \in T$ , all  $|x|, |v| \leq \eta$ ;

- (iii) the same as hypothesis  $H(F)_1$ (iii).

*Remark 6.* As before on account of Proposition 5 and by replacing  $F$  with  $F_0$ , without any loss of generality, we may assume that

$$(31) \quad |F(t, x)| \leq a_M(t) \quad \text{for a.a. } t \in T, \text{ all } x \in \mathbb{R}^N, \text{ with } a_M \in L^{p'}(T)_+.$$

Hypotheses  $H(F)_4$ (i), (ii) imply that  $(t, x) \rightarrow F(t, x)$  is graph measurable and then so is  $(t, x) \rightarrow \text{conv}F(t, x)$ . Finally note that

$$\text{ext conv}F(t, x) \subseteq F(t, x) \quad \text{for all } (t, x) \in T \times \mathbb{R}^N.$$

**Theorem 5.** *If hypotheses  $H(a)'$ ,  $H(r)$ ,  $H(F)_4$  hold, then  $\widehat{S}_e^{C_0^1(T, \mathbb{R}^N)} = \widehat{S}_c$ .*

*Proof.* As before  $C = \overline{\text{conv}}C_0 \in P_{kc}(C_0^1(T, \mathbb{R}^N))$  (see Proposition 6). Let  $u \in \widehat{S}_c$ . We have

$$-a(u')' - r(t)|u'|^{p-2}u' = f \quad \text{with } f \in S_{\text{conv}F(\cdot, u(\cdot))}^{p'}.$$

For  $v \in C$  and  $\varepsilon > 0$ , let  $R_\varepsilon^v : T \rightarrow 2^{\mathbb{R}^N} \setminus \{\emptyset\}$  be defined by

$$R_\varepsilon^v(t) = \{h \in \mathbb{R}^N : |f(t) - h| < \varepsilon + d(f(t), \text{conv}F(t, v(t))), h \in \text{conv}F(t, v(t))\}.$$

Clearly  $R_\varepsilon^v(\cdot)$  is graph measurable. So, we can use the Yankov-von Neumann-Aumann selection theorem (see Hu-Papageorgiou [8], Theorem 2.14, p. 158) and infer that  $R_\varepsilon^v(\cdot)$  admits a measurable selection. Such a selection belongs in  $L^{p'}(T, \mathbb{R}^N)$  (see (31)).

Next let  $L_\varepsilon : C \rightarrow 2^{L^{p'}(T, \mathbb{R}^N)}$  be defined by

$$L_\varepsilon(v) = S_{R_\varepsilon^v}^{p'} \quad \text{for all } v \in C.$$

From the previous argument, we see that  $L_\varepsilon(v) \neq \emptyset$  for all  $v \in C$ . Also  $v \rightarrow L_\varepsilon(v)$  is lsc (see Hu-Papageorgiou [8], Lemma 8.3, p. 239). It follows that  $v \rightarrow \overline{L_\varepsilon(v)}$  is lsc and has decomposable values. So, we can find  $l_\varepsilon : C \rightarrow L^{p'}(T, \mathbb{R}^N)$  continuous such that

$$l_\varepsilon(v) \in \overline{L_\varepsilon(v)} \quad \text{for all } v \in C$$

(see Hu-Papageorgiou [8], Theorem 8.7, p. 245). Then on account of Theorem 8.31, p. 260, of [8], we can find a continuous map  $\theta_\varepsilon : C \rightarrow L_w^{p'}(T, \mathbb{R}^N)$  such that

$$(32) \quad \theta_\varepsilon(v) \in S_{\text{ext conv}F(\cdot, v(\cdot))}^{p'}, \|\theta_\varepsilon(v) - l_\varepsilon(v)\|_w < \varepsilon \quad \text{for all } v \in C.$$

Now let  $\varepsilon_n \rightarrow 0^+$  and set  $l_n = l_{\varepsilon_n}$ ,  $\theta_n = \theta_{\varepsilon_n}$  for all  $n \in \mathbb{N}$ . We consider the following Duffing system

$$-a(u'(t))' - r(t)u'(t) = \theta_n(u)(t) \quad \text{for a.a. } t \in T, u(0) = u(b) = 0, n \in \mathbb{N}.$$

This problem has a solution  $u_n \in C_0^1(T, \mathbb{R}^N)$  (see Theorem 2). Note that  $u_n \in C$  for all  $n \in \mathbb{N}$  (see (31)) and  $C \in P_{kc}(C_0^1(T, \mathbb{R}^N))$ . So, we may assume that

$$(33) \quad u_n \rightarrow \widehat{u} \text{ in } C_0^1(T, \mathbb{R}^N).$$

We have

$$-a(u'_n)' + a(u')' - r(t)[u'_n - u'] = \theta_n(u_n) - f \quad \text{for all } n \in \mathbb{N}.$$

We act with  $u_n - u$  and after integration by parts, we have

$$(34) \quad \begin{aligned} & \int_0^b (a(u'_n) - a(u'), u'_n - u')_{\mathbb{R}^N} dt - \int_0^b r(t)(u'_n - u', u_n - u)_{\mathbb{R}^N} dt \\ &= \int_0^b (\theta_n(u_n) - f, u_n - u)_{\mathbb{R}^N} dt. \end{aligned}$$

Let  $\eta = \max\{\sup_{n \geq 1} \|u_n\|_{C_0^1(T, \mathbb{R}^N)}, \|u\|_{C_0^1(T, \mathbb{R}^N)}\} < +\infty$  (see (33)). Hypothesis  $H(a)'$  implies that

$$(35) \quad \tilde{c}_\eta \|u'_n - u'\|_2^2 \leq \int_0^b (a(u'_n) - a(u'), u'_n - u')_{\mathbb{R}^N} dt.$$

Also, we have

$$(36) \quad \begin{aligned} \int_0^b r(t)(u'_n - u', u_n - u)_{\mathbb{R}^N} dt &\leq \|r\|_\infty \|u'_n - u'\|_2 \|u_n - u\|_2 \\ &\leq \frac{\|r\|_\infty}{\widehat{\lambda}_1^{1/2}} \|u'_n - u'\|_2^2. \end{aligned}$$

Finally note that

$$(37) \quad \begin{aligned} & \int_0^b (\theta_n(u_n) - f, u_n - u)_{\mathbb{R}^N} dt \\ &= \int_0^b (\theta_n(u_n) - l_n(u_n), u_n - u)_{\mathbb{R}^N} dt + \int_0^b (l_n(u_n) - f, u_n - u)_{\mathbb{R}^N} dt \\ &\leq \int_0^b (\theta_n(u_n) - l_n(u_n), u_n - u)_{\mathbb{R}^N} dt + \varepsilon_n b + \int_0^b k_\eta(t) |u_n - u|^2 dt \quad \text{for all } n \in \mathbb{N}. \end{aligned}$$

From (32) and Proposition 1, we have

$$(38) \quad \int_0^b (\theta_n(u_n) - l_n(u_n), u_n - u)_{\mathbb{R}^N} dt \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

We return to (34), use (35), (36), (37), pass to the limit as  $n \rightarrow +\infty$  and finally use (33), (38) and Jensen's inequality. Then

$$\begin{aligned} & \left[ \tilde{c}_\eta - \frac{\|r\|_\infty}{\widehat{\lambda}_1^{1/2}} - \|k_\eta\|_\infty b \right] \|\widehat{u}' - u'\|_2^2 \leq 0, \\ & \Rightarrow \widehat{u} = u \quad (\text{see hypothesis } H(F)_4(\text{ii})). \end{aligned}$$

Since  $u_n \in \widehat{S}_e$  and  $u_n \rightarrow u$  in  $C_0^1(T, \mathbb{R}^N)$  (see (33)), we conclude that  $\widehat{S}_e \stackrel{\text{---}}{=} \widehat{S}_c \stackrel{\text{---}}{=} C_0^1(T, \mathbb{R}^N) = \widehat{S}_e$ . □

*Remark 7.* A careful inspection of the proofs, reveals that the positivity of the drift coefficient  $r(\cdot)$  (see hypothesis  $H(r)$ ) was first used in the proof of Proposition 5. So, Theorems 2 and 3 are valid without the assumption that  $r(t) \geq 0$  for a.a.  $t \in T$ . It will be interesting to know if we can remove this restriction also in Theorems 4 and 5. Theorems 4 and 5 provide answers to questions raised at the end of [14]. Extremal trajectories and a strong relaxation theorem, were proved for a different class of multivalued second order systems, in Papageorgiou-Vetro-Vetro [13].

We conclude with an example of a control system.

**Example 3.** Consider the following control system:

$$(39) \quad \begin{cases} -a(u'(t))' - r(t)|u'(t)|^{p-2}u'(t) = f(t, u(t))v(t) & \text{a.e. on } T, \\ u(0) = u(b) = 0, v(t) \in V(t) & \text{for a.a. } t \in T. \end{cases}$$

In this system,  $f : T \times \mathbb{R}^N \rightarrow \mathcal{L}(\mathbb{R}^m, \mathbb{R}^N)$  is a Carathéodory function and  $V : T \rightarrow P_k(\mathbb{R}^m)$  is graph measurable. We assume that

$$\begin{aligned} |f(t, x)| &\leq a_\eta(t) \quad \text{for a.a. } t \in T, \text{ all } |x| \leq \eta \text{ with } a_\eta \in L^p(T), \\ |U(t)| &\leq \theta \quad \text{for a.a. } t \in T, \text{ some } \theta > 0, \end{aligned}$$

and there exists  $M$  such that

$$(f(t, x)u, x)_{\mathbb{R}^N} < 0 \quad \text{for a.a. } t \in T, \text{ all } |x| \geq M, \text{ all } u \in U(t).$$

Then the control system (39) has admissible state-control pairs and if  $f(t, \cdot)$  is locally Lipschitz with local Lipschitz constant  $\widehat{k}_\eta(\cdot) \in L^1(T)$ , then the states of the nonconvex problem are  $C_0^1(T, \mathbb{R}^N)$ -dense in those of the convexified system (control constraint set  $\text{conv } U(t)$  for all  $t \in T$ , see Theorem 5).

#### ACKNOWLEDGEMENT

The authors wish to thank the two referees for their useful remarks and suggestions.

#### REFERENCES

- [1] J. FRANÇÙ, *Monotone operators. A survey directed to applications to differential equations*, Appl. Math. **35** (1990), 257–301.
- [2] M. GALEWSKI, *On the Dirichlet problem for a Duffing type equation*, Electron. J. Qual. Theory Differ. Equ. **2011**:15 (2011), 1–12.
- [3] L. GASIŃSKI, N. S. PAPAGEORGIOU, *Nonlinear Analysis*, Ser. Math. Anal. Appl., 9, Chapman and Hall/CRC Press, Boca Raton, Florida, 2006.
- [4] ———, *Exercises in Analysis*, Part 1, Problem Books in Mathematics, Springer, Cham, 2014.
- [5] ———, *Exercises in Analysis*, Part 2, Problem Books in Mathematics, Springer, Cham, 2016.
- [6] P. HARTMAN, *On boundary value problems for systems of ordinary nonlinear second order differential equations*, Trans. Amer. Math. Soc. **96** (1960), 493–509.
- [7] ———, *Ordinary Differential Equations*, Wiley, New York, 1964.
- [8] S. HU, N. S. PAPAGEORGIOU, *Handbook of Multivalued Analysis*. Vol. I: Theory, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1997.
- [9] ———, *Handbook of Multivalued Analysis*. Vol. II: Applications, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2000.
- [10] P. KALITA, P. M. KOWALSKI, *On multivalued Duffing equation*, J. Math. Anal. Appl. **462** (2018), 1130–1147.
- [11] P. M. KOWALSKI, *Well-posed Dirichlet problems pertaining to the Duffing equation*, Electron. J. Qual. Theory Differ. Equ. **2014**:59 (2014), 1–15.
- [12] R. MANÁSEVICH, J. MAWHIN, *Boundary value problems for nonlinear perturbations of vector  $p$ -Laplacian-like operators*, J. Korean Math. Soc. **37** (2000), 665–685.
- [13] N. S. PAPAGEORGIOU, C. VETRO, F. VETRO, *Extremal solutions and strong relaxation for nonlinear multivalued systems with maximal monotone terms*, J. Math. Anal. Appl. **461** (2018), 401–421.
- [14] ———, *Nonlinear multivalued Duffing systems*, J. Math. Anal. Appl. **468** (2018), 376–390.
- [15] P. TOMICZEK, *Remark on Duffing equation with Dirichlet boundary condition*, Electron. J. Differential Equations **2007**:81 (2007), 1–3.

(N.S. Papageorgiou) DEPARTMENT OF MATHEMATICS, NATIONAL TECHNICAL UNIVERSITY, ZOGRAFOU CAMPUS, 15780, ATHENS, GREECE  
 Email address: npapg@math.ntua.gr

(C. Vetro) DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF PALERMO,  
VIA ARCHIRAFI 34, 90123, PALERMO, ITALY

*Email address:* `calogero.vetro@unipa.it`

(F. Vetro) NONLINEAR ANALYSIS RESEARCH GROUP, TON DUC THANG UNIVERSITY, HO CHI  
MINH CITY, VIETNAM; FACULTY OF MATHEMATICS AND STATISTICS, TON DUC THANG UNIVER-  
SITY, HO CHI MINH CITY, VIETNAM

*Email address:* `francescavetro@tdtu.edu.vn`