

Relaxed selection diminishes social memory and expression of host defenses against cuckoos

Deryk Tolman ^{1,2}, Katja Rönkä ^{1,2}, Edward Kluen ^{1,2}, Juho Jolkkonen ³, Pietro Di Bari ¹, Renzo Ientile ⁴, Daniela Campobello ⁵, Rose Thorogood ^{1,2}

¹HiLIFE Helsinki Institute of Life Science, University of Helsinki, Helsinki, Finland

²Research Programme in Organismal and Evolutionary Biology, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland

³Department of Biological and Environmental Science, University of Jyväskylä, Jyväskylä, Finland

⁴Department of Biological, Geological and Environmental Sciences, Section of Animal Biology “M. La Greca,” University of Catania, Catania, Italy

⁵Dipartimento STEBICEF, Università degli Studi di Palermo, Palermo, Italy

Corresponding author: Research Programme in Organismal and Evolutionary Biology, Faculty of Biological and Environmental Sciences, University of Helsinki, Biokeskus 3, Viikinkaari 1, 00790 Helsinki, Finland. Email: deryk.tolman@helsinki.fi

Abstract

Behavioral defenses are a key adaptation across animals, but confirming decay under relaxed selection is challenging. If expression involves decision-making, then defenses may only be expressed when the right environmental cues are available (i.e., cryptic plasticity). Even if defenses wane, they could be quickly restored once selection resumes as animals can learn by observing others (i.e., social memory). Here we tested cryptic plasticity and social memory under relaxed selection in a cuckoo host, the common reed warbler, by manipulating perceived parasitism risk with social information in geographically distinct areas differing in allopatry (100–1,000 years). Despite predicting that social information of a cuckoo vs. a control would elicit otherwise cryptic behavioral defenses, only birds that mobbed model cuckoos prior to manipulation upregulated this defense, and only in the recently allopatric population. Social information also had no effect on low-to-absent rates of foreign egg rejection, a second line of defense common to cuckoo host species. These results contrast with parasitized locations where similar methods provoke mobbing and increase egg rejection 10-fold. Our study therefore suggests that, even when manipulating information to account for reaction norms, behavioral defenses may degrade rapidly under relaxed selection, demonstrating a process facilitating a geographic mosaic of coevolution.

Keywords: relaxed selection, geographic mosaic of coevolution, host–parasite coevolution, behavioral plasticity, social information

Introduction

As species’ ranges expand or contract, their exposure to enemies and selection pressures can also change (Gilman et al., 2010). Where selection relaxes (i.e., with allopatry), traits that were previously adaptive are expected to reduce (e.g., due to drift), particularly if maintaining the trait is costly (Lahti et al., 2009). However, while behavioral defenses are common across the animal kingdom (Candolin & Wong, 2015), whether these respond to relaxed selection is rarely tested (Foster, 2013; Lahti et al., 2009). Behaviors are assumed unlikely to be lost due to complex genetic architectures (pleiotropy) and underlying trait variation that can be co-opted in response to other selective pressures (Lahti et al., 2009). However, another key issue is that loss of behavior can be challenging to measure. Behavioral responses depend on prior experience and availability of information about the relative costs or benefits of taking action (i.e., information ecology; Dall et al., 2005). Therefore, the absence of a response does not necessarily indicate a lack of enemy recognition but rather cryptic plasticity (Foster, 2013). Taking an informational approach (Gil et al., 2018; O’Connor et al., 2019) and manipulating cues in allopatry, however, makes it

possible to approximate the retention of behavioral reaction norms in the wild.

Social “memory” could play a key role in determining whether behaviors are lost or retained. For example, prey and hosts observe the behavior of others to learn about novel threats (Feeney & Langmore, 2013; Langmore et al., 2012; Rowell et al., 2020) and assess the risk of predation or parasitism (Gil et al., 2018), while predators use social information (SI) to find suitable prey (Hämäläinen et al., 2022). Therefore, how individuals adjust their behavior in response to SI (i.e., social responsiveness; de Groot et al., 2023) is likely to affect both their own fitness and the strength of selection on interacting parties (Cantor et al., 2021; Thorogood & Davies, 2012). Social responsiveness may also determine whether populations are resilient to environmental change (Gil et al., 2019; Thorogood et al., 2018), especially as the utility of SI depends on the availability of more-informed “demonstrators.” For example, gene flow among populations experiencing variable selection pressures (i.e., trait re-mixing) is thought to determine how quickly a host population responds to shifts in the strength of selection by parasites (Geographic Mosaic The-

ory of Coevolution; Thompson, 2005). If experienced individuals from “hotspots” of current selection also introduce SI to naïve “cold spots” of relaxed selection, social learning could speed up the recovery of defenses in host populations that are later reinvaded by a previous enemy. However, while recent experimental evolution studies show that social behaviors can diminish very rapidly after selection is removed (Bladon et al., 2023), the role of social cues in maintaining behaviors in “cold spots” is still largely unknown (Lahti et al., 2009; Langmore et al., 2012).

An ideal system to test if behavioral defenses and social responsiveness decline in allopatry, altering the potential for hosts to respond to future (re)invasion, is avian brood parasitism. Virulent brood parasites lay their eggs in the nests of host species and exert strong selection for defensive behaviors such as mobbing or rejecting suspicious eggs. As many brood parasites are aggressive or Batesian mimics (Feeney et al., 2015; Welbergen & Davies, 2011), hosts are also expected to adjust defenses according to local risk to minimize costly mistakes (Lindholm & Thomas, 2000; Thorogood & Davies, 2013). Furthermore, parasitism can also vary in time and space, with some host populations or species escaping parasitism for many generations (Grim & Stokke, 2016). Over time in allopatry, plasticity is expected to be lost with hosts either becoming resistant to reinvasion (i.e., fixation of defenses) or susceptible to extinction if defenses wane (Soler, 2014), but it is unclear which trajectory will occur as there are examples of both. Some species long separated from brood parasites (i.e., hundreds of thousands to millions of years) show persistent egg rejection when experimentally induced (Bolen et al., 2000; Briskie et al., 1992; Chaumont et al., 2021; Hale & Briskie, 2007; Moskat et al., 2002; Rothstein, 2001); however, it appears to reduce in others (Briskie et al., 1992; Davies & Brooke, 1989; Soler & Møller, 1990; Yang et al., 2015, 2021). Frontline defenses, such as mobbing of adult brood parasites, also appear to either degrade (Briskie et al., 1992; Hale & Briskie, 2007; Lawson et al., 2020, 2023) or be preserved (Briskie et al., 1992; Chaumont et al., 2021; Kuehn et al., 2016) under allopatry. These defenses are not necessarily coupled; for example, egg rejection may be expressed while mobbing is not (Hale & Briskie, 2007; Thorogood & Davies, 2016). In addition, apparently naïve species can obtain host defenses very rapidly if invaded (Nakamura, 1991), but whether these changes represent the cryptic persistence of defenses in the population, gene flow from parasitized areas, or novel learning is unknown.

Here, we use allopatric populations of Common reed warblers (*Acrocephalus scirpaceus*) at the northern- and southernmost extents of their European breeding range to test for the retention of behavioral defenses and social responsiveness to brood-parasitic common cuckoos (*Cuculus canorus*). Reed warblers colonized Finland ca. 100 years ago (Järvinen & Ulfstrand, 1980), most likely expanding from the southwest (Bergman et al., 2025) and away from regularly parasitized areas in the range core (Avilés et al., 2006). Cuckoos in Finland use different host species, away from reedbeds. In Sicily, reed warblers are likely to have escaped parasitism by cuckoos for much longer—cuckoos only occur on mountainsides, while reed warblers are largely restricted to coastal areas, and mentions of both species in historical records are not in association with each other (Arnott, 2007). At parasitized sites in their range-core, reed warblers upregulate mobbing

(Campobello & Sealy, 2011; Davies & Welbergen, 2009) and egg rejection (Thorogood & Davies, 2016) defenses after observing their neighbors attack cuckoos. Among populations not currently exposed to parasitism, however, egg rejection rates vary widely (e.g., from 0% to 40%; Thorogood & Davies, 2013), and mobbing responses are reduced (e.g., 25% of pairs; Welbergen & Davies, 2012). However, as in other studies of brood-parasite host defenses in allopatry (see also Rothstein, 2001), perceived parasitism risk has not yet been manipulated (e.g., via SI), so the extent to which host defenses reduce under relaxed selection remains unclear.

Therefore, we experimentally manipulated SI and simulated parasitism in field experiments to test if defenses and/or social responsiveness decay in allopatry. We predicted that if defenses are retained, then providing SI should induce plasticity (i.e., social responsiveness) and lead to an increase in mobbing (propensity and intensity) and egg rejection, as previously shown in range-core populations. However, if defenses are not upregulated in response to our manipulated cues, then this would provide evidence that these behaviors are being lost under relaxed selection. Additionally, we assess changes in latency to approach threats at the nest as a more subtle indication of how these hosts perceive cuckoos, since we expect reed warblers to be more attentive when they perceive parasitism risk to be greater (Davies & Welbergen, 2009), and more cautious to approach if they mistake cuckoos for predatory hawks (Thorogood & Davies, 2012; Welbergen & Davies, 2011).

Materials and methods

Study areas and host populations

We studied unparasitized populations of common reed warblers breeding in May–July around Helsinki, southern Finland (30 discrete reed-lined bays along 100 km of coastline, 60.19848N 24.07305E to 60.34039N 25.71162E, in 2019 and 2020) and in the Fiume Ciane nature reserve near Siracusa, Sicily, Italy (37.04285N 15.24273E, in 2022), avoiding re-use of areas across years in Finland to avoid contamination of experience or pseudoreplication. Reed warblers nest at relatively low densities in Helsinki (Tolman et al., 2021) but are more numerous in Fiume Ciane (median distance to closest nest = 36 m, interquartile range = 24–53 m). We found nests by cold-searching or monitoring singing males and recorded location and distance to the nearest potential cuckoo vantage point (tree > 3 m in height, a possible indirect cue of parasitism risk; Oien et al., 1996; Welbergen & Davies, 2009) using handheld GPS (accuracy ± 1 m).

Experimental protocols

To compare results to previously studied parasitized populations in the range-core, we adopted published protocols (Davies & Welbergen, 2009; Thorogood & Davies, 2016) as follows. Experiments started on the day the fourth egg was laid (clutch sizes = 3–5 eggs, incubation begins with the penultimate egg; Cramp & Brooks, 1992). At each nest, we measured the change in mobbing response to a cuckoo after presentation of SI of either a cuckoo or a control. We used the same 3D-printed model cuckoos (painted by an artist, see Tolman et al., 2021 for details) in Finland and Sicily but controls differed. In Finland, we used a commercially available plastic common magpie (*Pica pica*; Live Decoys) to assess




Figure 1. Reed warbler mobbing defenses (calls per 5 min) in allopatry (Finland: A–C, Sicily: D–F) before (gray dots) and after (colored dots) experimental manipulation of social information (“SI”) about cuckoos (B, E) vs. a control (C—magpie, F—teal). Details of experimental design are shown in (A) and (D). In Finland (A), the first model presented at nests in Finland (P1) was always a cuckoo, followed by either a cuckoo or magpie (P2) depending on social information treatment (SI), and finished with a cuckoo presentation (P3) for the “after” comparison. In Sicily (D), cuckoo and control model presentation order was alternated before (P1–P2) and after (P3–P4) the social information presentations.

dence of damage was treated as a rejection attempt. Finally, in Sicily, we also assessed non-mimetic egg rejection without any information of cuckoo presence ($n = 11$) and with personal information of parasitism risk (i.e., by presenting a cuckoo model at the nest; $n = 12$) by coloring one egg in the nest entirely with a blue felt-tip pen (Copic Ciao B). Immaculate blue eggs (resembling redstart-gens cuckoo eggs) are highly dissimilar from reed warblers' and consistently elicit some of the highest rejection rates (often >60%) from this host despite little standardization of color or size across studies (Davies & Brooke, 1988; Dyrce & Halupka, 2007; Stokke et al., 1999, 2010).

Statistical analyses

All analyses were conducted in R v.4.3.2 (R Core Team, 2023). First, we used GLMs to check for bias of potential indirect cues (i.e., distance to potential cuckoo vantage points, see Supporting Information S1) and to confirm that initial responses to cuckoos (i.e., P1 in Finland, or P1|P2 in Sicily) were similar in the treatment groups and among years of the study (in Finland). We then used GLMMs with a random effect for nest identity when modeling change in response across multiple presentations (i.e., P1 vs. P3, or P1 vs. P2) as implemented in the package *glmmTMB* (Brooks et al., 2017). To investigate whether SI led to a change in mobbing propensity or intensity, we only included responses to the first and last presentation of a cuckoo at the nest (i.e., P1 and P3 in Finland). However, due to a large proportion of zeroes and lack of suitable fit for a zero-inflated model structure, we adopted a hurdle approach to model mobbing propensity (i.e., binary variable, binomial error structure, and logit-link function) and intensity (i.e., number of calls in at least one presentation > 0, quasi-Poisson (*nbinom1*) distribution) separately in Finland. For mobbing data from Sicily, we used Fisher's exact tests to compare treatment effects on mobbing propensity and quasi-binomial GLMMs to investigate changes in mobbing intensity across all pairs, due to the smaller sample size. To investigate whether SI affected egg rejection decisions, we used a GLM (binomial error structure, logit-link function) that included clutch size as a covariate predictor. Despite data being collected across 30 discrete sites in Finland, including location as a random effect did not improve model fit (Akaike information criterion > 2), so it was not part of our final models.

Finally, we investigated latency to approach the threat at the nest during experiments to assess more subtle differences in reaction to cuckoos (i.e., cryptic recognition without a mobbing response). We used separate GLMMs with Tweedie distributions to test if change in latency across presentations differed (i) between treatments and (ii) between mobbing vs. non-mobbing pairs as we lacked power to test a Presentation order*SI treatment*Mobbing status three-way interaction. Nest identity was included as a random effect, and pairwise contrasts were estimated using estimated marginal means with Tukey-adjusted p -values. We used QQ-normality plots, residuals vs. predicted plots, and dispersion and outlier tests using package *DHARMa* (Hartig, 2022) to assess assumptions and fits of all models, and we present estimated mean effect sizes ($\pm$ standard error), z -statistics, and p -values. Summary tables of all models are available in Supporting Information S2.

Results

Mobbing defenses

When first presented with a cuckoo at the nest, 36% (30/83) of pairs in Finland and 13% (5/40) of pairs in Sicily showed a mobbing response (Figure 1). This initial propensity to mob did not differ between SI treatment groups (Finland, GLM: estimate = 0.16 ± 0.46 , $z = -0.35$, $p = .73$; Sicily, Fisher's Exact test: $p = 1.0$; Table S2.1 in Supporting Information 2) or among years of the study in Finland (GLM: estimate = 0.25 ± 0.48 , $z = 0.52$, $p = .60$; Table S2.1).

Providing SI about the risk of parasitism in the local area did not induce significantly more reed warblers to mob cuckoos than pairs exposed to information about magpies (in Finland, GLMM Presentation order * SI treatment: estimate = 3.99 ± 2.39 , $z = 1.67$, $p = .10$, Figures 1B and 2A; Table S2.2) or when compared to SI about innocuous teals (in Sicily, across treatments $n = 10$ pairs with calls > 0, Fisher's exact test: $p = 1.00$, Figure 2A). This lack of effect was not because SI about controls also induced increased mobbing propensity (Finland: Figures 1C and 2A; Sicily: Figure 1F).

SI about cuckoos did, however, increase mobbing intensity (i.e., number of signals) but only for pairs who responded in Finland (across treatments $n = 36$ pairs with calls > 0, Figure 1B, C; GLMM Presentation order * SI treatment: estimate = 0.97 ± 0.45 , $z = 2.14$, $p = .03$, Figure 2B; Table S2.2). There was no significant change among presentations in the group who received SI about magpies (GLMM estimate = -0.28 ± 0.33 , $z = -0.84$, $p = .40$, Figure 2B; Table S2.2). Instead, these pairs were just as likely to increase or reduce mobbing intensity by the third presentation (P3; Figure 1C). In Sicily, SI had no effect on mobbing intensity (GLMM Presentation order * SI treatment: estimate = -1.20 ± 0.88 , $z = -1.36$, $p = .17$, Figures 2B and 1E, F; Table S2.3). Teal were mobbed at similar intensities between Sicily and Finland ($n = 96$, GLM: estimate = -0.39 ± 0.31 , $z = -1.27$, $p = .21$; Table S2.4).

Egg rejection defenses

Reed warblers did not reject "mimetic" brown-spotted eggs in either Finland (cuckoo treatment = 0/24; magpie treatment = 0/25; Figure 3) or Sicily (cuckoo treatment = 0/21; teal treatment = 0/19), and there was no rejection of non-mimetic blue eggs in Sicily (no cuckoo presentation = 0/11, after cuckoo presentation at nest = 0/12). Blue eggs were however rejected at low rates in Finland, although this did not differ between SI treatments (GLM: estimate = 0.11 ± 0.64 , $z = 0.17$, $p = .97$; Figure 3; Table S2.5).

Recognition of cuckoos vs. response

In Finland, non-mobbing birds were slower initially to approach a cuckoo at their nest than mobbing pairs (P1 data only, GLM: estimate = -0.42 ± 0.19 , $z = -2.29$, $p = .02$; Figure 4; Table S2.6) but became faster with subsequent presentations (GLMM Mobbing status * Presentation order, P1 and P3 only, interaction: estimate = 0.54 ± 0.28 , $z = 1.96$, $p = .049$; Figure 4; Table S2.7). However, this was not specific to SI because non-mobbers already reduced their latency to approach cuckoos by P2 (P1 vs. P2 non-mobbers, GLMM: estimate = -0.44 ± 0.21 , $z = -2.04$, $p = .10$; Table S2.8), and they decreased again after SI of magpies (P2 vs. P3 non-




Figure 2. (A) Propensity to mob cuckoos did not change after social information about cuckoos or a control in Finland or Sicily. (B) Mobbing pairs showed social responsiveness (i.e., increased intensity) to cuckoos, but only in Finland (distributions of changes in mobbing intensity, vertical lines = medians). (C) Mobbing reaction norms of reed warblers across parasitism risk (before and after social information of cuckoo activity) were more frequently positive in parasitized Wicken Fen (produced from [Davies & Welbergen, 2009](#) and [Thorogood & Davies, 2012](#) with permission) than in Finland and Sicily. Negative slopes are highlighted in red, and sample sizes represent the total number of pairs tested (top) and the number of slopes that remained at 0 (i.e., did not mob at all; bottom).

mobbers: estimate = -0.43 ± 0.24 , $z = -1.8$, $p = .17$; [Table S2.8](#); [Figure 4B](#)). Mobbing pairs showed no change in latency to approach cuckoos across the experiment (P1 vs. P3 Cuckoo, GLMM: estimate = 0.10 ± 0.29 , $z = 0.33$, $p = .94$; [Table S2.9](#); Magpie: estimate = 0.17 ± 0.28 , $z = 0.61$, $p = .81$; [Table S2.8](#)), and all were more wary of approaching a magpie at the nest than a cuckoo (P2 between treatments, GLM: estimate = 0.72 ± 0.26 , $z = 2.81$, $p = .005$; [Figure 4A](#); [Table S2.10](#)) even if they never mobbed (mobbing status * threat: estimate = 0.11 ± 0.38 , $z = 0.29$, $p = .78$; [Figure 4](#); [Table S2.10](#)).

Reed warblers' latencies in Sicily did not change over the experiment, either overall (GLMM Presentation order: estimate = 0.20 ± 0.21 , $z = 0.95$, $p = .34$; [Table S2.10](#)) or differently according to SI treatments (GLMM Presentation order * SI treatment: estimate = 0.67 ± 0.42 , $z = 1.58$, $p = .12$; [Table S2.11](#)). They also approached teal as quickly as

they did cuckoos (estimate = 0.07 ± 0.17 , $z = 0.39$, $p = .67$; [Table S2.12](#)).

Discussion

A wealth of prior research in parasitized areas has revealed that reed warblers' defenses predictably match local brood parasitism risk, including responding to SI of cuckoos by consistently increasing mobbing and egg rejection rates ([Campobello & Sealy, 2011](#); [Davies & Welbergen, 2009](#); [Thorogood & Davies, 2016](#)). But theory also predicts that the loss of defenses in non-parasitized populations is a key part of determining coevolutionary outcomes ([Soler, 2014](#); [Thompson, 2005](#)). Although behaviors are assumed hard to lose under relaxed selection, by using SI to simulate increased brood parasitism risk, we found that defenses and social responsiveness of a common host of cuckoos are de-




Figure 3. Rejection rates of experimental eggs in Finland (inset illustrations). Blue non-mimetic eggs (left) were rejected at the same rate regardless of whether social information was provided about cuckoos (24%) or magpies (22%). Mimetic brown-spotted eggs (right) were not rejected.




Figure 4. Latency of mobbing and non-mobbing pairs to approach presentations of models at their nest (indicated by icons) before and after social information (SI; dashed line) about (A) cuckoos or (B) a magpie control in Finland. Raw data are plotted as diamonds, with darker symbols indicating pairs that mobbed cuckoos. Predicted mean values ($\pm$ standard error) from a GLMM are plotted as circles and bars, with connecting lines indicating change across model presentation order (P1–P3).

grading in areas of allopatry. At the recent range front in Finland, only pairs that mobbed a cuckoo increased their mobbing intensity after receiving SI. The other two-thirds of birds largely remained quiet, although they potentially retained some recognition of cuckoos since both mobbing and non-mobbing pairs approached the nest more quickly than when presented with a magpie, suggesting the decrease in latency of non-mobbers was not purely driven by habituation to disturbance in general, but is specific to the threat. Still, this may reflect avoidance of magpies (Trnka & Požgayová, 2017) rather than awareness of cuckoos and requires further experimental work. Our findings in Finland are enhanced by our results from the Sicilian population, where reed warblers have been in allopatry for longer. There, they expressed defenses even more rarely and showed no evidence of SI use. Importantly, mimetic “cuckoo” eggs remained in nests, and although non-mimetic eggs were rejected at low rates in Finland, SI had no influence on this defense either. This suggests that both allopatric populations would be unlikely to express plasticity or mount significant resistance should cuckoos invade.

Our results are likely evidence for the decay of ancestral defenses rather than a methodological failure to detect responses, for three reasons. First, providing SI experimentally led to an increase in mobbing intensity in Finland, even if not in Sicily. Second, we followed very similar methods to previous studies conducted with reed warblers at parasitized sites, where SI heightened near-absent egg rejection of similarly manipulated eggs 10-fold (Thorogood & Davies, 2016) and induced mobbing of cuckoos (Campobello & Sealy, 2011; Davies & Welbergen, 2009; Thorogood & Davies, 2012), including for pairs that showed no initial response (e.g., 54% of quiet pairs mobbed cuckoos after SI: Davies & Welbergen, 2009). Although there were some differences in our use of controls and presentations of cuckoos at the nest, these represent similar variation to range-core experiments (e.g., Campobello & Sealy, 2011; Davies & Welbergen, 2009). Third, the differences we detected were specific to cuckoos as responses to a control teal and nest predator (magpie) were similar at our sites and to parasitized areas (Campobello & Sealy, 2010; Tolman et al., 2021; Welbergen & Davies, 2008; Welbergen & Davies, 2012) despite large ecological differences (i.e., climate, nesting density). Some level of cuckoo recognition might be retained in Finland as non-mobbers seemed slower initially to approach a cuckoo at their nest than mobbing pairs, but became faster with subsequent presentations. If non-mobbers perceived cuckoo models as hawks, we expected that they would instead become more wary (Thorogood & Davies, 2012), but mobbing pairs did not decrease their latency as expected, so it is difficult to conclude that this change in attentiveness is evidence of recognition after all. By using consistent experimental designs to previous studies, testing a large number of reed warblers nesting across a broad range of environmental variation, and considering social responsiveness explicitly, we can therefore be confident that the weak defenses in our allopatric study populations are not the result of cryptic plasticity (Scheiner & Levis, 2021). Instead, they provide compelling evidence that behavioral defenses may sometimes decay in allopatry from cuckoos.

Given that studies of animal culture across a range of taxa demonstrate that the spread of even arbitrary behaviors is possible (Whiten, 2021), it is surprising that we did not de-

tect evidence for social learning among “naïve” reed warblers (i.e., those that did not mob cuckoos). For example, juvenile superb fairy wrens begin to mob cuckoos after observing adults do so (Feeney & Langmore, 2013), and naïve individuals of several species can be trained to recognize novel predators (e.g., introduced species; Rowell et al., 2020) when paired with mobbing calls. Often social learning of threats requires only one exposure (i.e., similar to our experimental design; reviewed by Crane & Ferrari, 2013) and can even be effective when using arbitrary cues (Hämäläinen et al., 2022). Why then did so many birds not mob cuckoos or reject eggs? One possible explanation is that ontogenetic experience or repeated exposure may be necessary before behaviors are expressed, as in the antipredator responses of several fish and mammals (Kelley & Magurran, 2003; Mineka et al., 1984) and personal encounters with avian or piscine brood parasites (Blažek et al., 2018; Hauber et al., 2006; Stokke et al., 2007). Alternatively, but not mutually exclusively, multiple opportunities to learn socially may be needed to elicit a response. While we cannot exclude these possibilities, they seem unlikely to explain the lack of response we detected here: we provided three encounters with cuckoos at or near their nests, yet most birds still did not respond, and under natural conditions naïve pairs are unlikely to repeatedly encounter more responsive neighbors when parasitism rates are low to absent. Perhaps with relaxed selection, there is an initial loss of recognition that then limits hosts’ potential for observational learning.

Temporal and geographical variation in host defenses has typically been explained by plasticity (Langmore et al., 2012; Lindholm, 2000; Welbergen & Davies, 2012), particularly in reed warblers where individuals track local parasitism risk and adjust defenses accordingly (Thorogood & Davies, 2013). However, here we found only limited evidence for plasticity in our allopatric populations in response to either indirect environmental or social cues of heightened parasitism risk. This is surprising, given that in a previous study at another unparasitized site of similar length of allopatry to Finland (Llangorse Lake, Wales; Lindholm, 1999), reed warblers responded more aggressively to cuckoo models if nesting closer to vantage points for cuckoos (Welbergen & Davies, 2012). We found no evidence for this indirect cue use: instead, birds nesting further from vantage points in Finland were *more* likely to mob, and mobbing in Sicily was unaffected (Supporting Information S1). Nevertheless, of the birds that mobbed, their initial intensities were similar across all three allopatric sites (Finland: 225 ± 43 calls per 5 min, Sicily: 280 ± 120 , Llangorse: 281 ± 64 ; Welbergen & Davies, 2012), suggesting that while the frequency of birds retaining recognition of cuckoos may vary across allopatric populations, reaction norm intercepts have the potential to remain similar. However, comparing reactions to social cues experimentally presented in a similar manner across studies highlights a key difference in allopatry. Previous studies at Wicken Fen, a parasitized site, induced positive social plasticity (*sensu de Groot et al., 2023*) in almost all responding pairs (94.4% across two studies; Davies & Welbergen, 2009; Thorogood & Davies, 2012). At our allopatric sites, on the other hand, reed warblers appeared to be more often “confused,” with 30%–40% of responding pairs instead reducing mobbing intensity after SI of cuckoos (Figure 2C). Interestingly, a similar proportion of negative slopes was observed for birds presented

- eggs. *Biology Letters*, 2(2), 177–180. <https://doi.org/10.1098/rsbl.2005.0438>
- Järvinen, O., & Ulfstrand, S. (1980). Species turnover of a continental bird fauna: Northern Europe, 1850–1970. *Oecologia*, 46(2), 186–195. <https://doi.org/10.1007/BF00540125>
- Jolly, C. J., Webb, J. K., & Phillips, B. L. (2018). The perils of paradise: An endangered species conserved on an island loses antipredator behaviours within 13 generations. *Biology Letters*, 14(6), 20180222. <https://doi.org/10.1098/rsbl.2018.0222>
- Kelley, J. L., & Magurran, A. E. (2003). Effects of relaxed predation pressure on visual predator recognition in the guppy. *Behavioral Ecology and Sociobiology*, 54(3), 225–232. <https://doi.org/10.1007/s00265-003-0621-4>
- Kuehn, M. J., Peer, B. D., McCleery, R. A., & Rothstein, S. I. (2016). Yellow warbler defenses are retained in the absence of brood parasitism but enhanced by experience with cowbirds. *Behavioral Ecology*, 27(1), 279–286. <https://doi.org/10.1093/beheco/arv145>
- Lahti, D. C., Johnson, N. A., Ajie, B. C., Otto, S. P., Hendry, A. P., Blumstein, D. T., Coss, R. G., Donohue, K., & Foster, S. A. (2009). Relaxed selection in the wild. *Trends in Ecology & Evolution*, 24(9), 487–496. <https://doi.org/10.1016/j.tree.2009.03.010>
- Langmore, N. E., Feeney, W. E., Crowe-Riddell, J., Luan, H., Louwrens, K. M., & Cockburn, A. (2012). Learned recognition of brood parasitic cuckoos in the superb fairy-wren *Malurus cyaneus*. *Behavioral Ecology*, 23(4), 798–805. <https://doi.org/10.1093/beheco/ars033>
- Lawson, S. L., Enos, J. K., Fernandez-Duque, F., Kleindorfer, S., Ward, M. P., Gill, S. A., & Hauber, M. E. (2023). Absence of referential alarm calls in long-term allopatry from the referent: A case study with Galapagos yellow warblers. *Behavioral Ecology and Sociobiology*, 77(8), 99. <https://doi.org/10.1007/s00265-023-03372-0>
- Lawson, S. L., Leuschner, N., Gill, B. J., Enos, J. K., & Hauber, M. E. (2020). Loss of graded enemy recognition in a Whitehead population allopatric with brood-parasitic long-tailed cuckoos. *Avocetta*, 44(1), 4–10. <https://doi.org/10.30456/AVO.2020104>
- Lindholm, A. (2000). Tests of phenotypic plasticity in reed warbler defences against cuckoo parasitism. *Behaviour*, 137(1), 43–60. <https://doi.org/10.1163/156853900501863>
- Lindholm, A. K. (1999). Brood parasitism by the cuckoo on patchy reed warbler populations in Britain. *Journal of Animal Ecology*, 68(2), 293–309. <https://doi.org/10.1046/j.1365-2656.1999.00286.x>
- Lindholm, A. K., & Thomas, R. J. (2000). Differences between populations of reed warblers in defences against brood parasitism. *Behaviour*, 137(1), 25–42. <https://doi.org/10.1163/156853900501854>
- Marton, A., Fülöp, A., Ozogány, K., Moskát, C., & Bán, M. (2019). Host alarm calls attract the unwanted attention of the brood parasitic common cuckoo. *Scientific Reports*, 9(1), 1. <https://doi.org/10.1038/s41598-019-54909-1>
- Mineka, S., Davidson, M., Cook, M., & Keir, R. (1984). Observational conditioning of snake fear in rhesus monkeys. *Journal of Abnormal Psychology*, 93(4), 355–372. <https://doi.org/10.1037/0021-843X.93.4.355>
- Moskát, C., Szentpéteri, J., & Barta, Z. (2002). Adaptations by great reed warblers to brood parasitism: A comparison of populations in sympatry and allopatry with the common cuckoo. *Behaviour*, 139(10), 1313–1329. <https://doi.org/10.1163/156853902321104181>
- Nakamura, H. (1991). Brood parasitism by the cuckoo *Cuculus canorus* in Japan and the start of new parasitism on the azure-winged magpie *Cyanopica cyana*. *Japanese Journal of Ornithology*, 39(1), 1–18. <https://doi.org/10.3838/jjo.39.1>
- Nuismer, S. L., Thompson, J. N., & Gomulkiewicz, R. (2003). Coevolution between hosts and parasites with partially overlapping geographic ranges. *Journal of Evolutionary Biology*, 16(6), 1337–1345. <https://doi.org/10.1046/j.1420-9101.2003.00609.x>
- O'Connor, M. I., Pennell, M. W., Altermatt, F., Matthews, B., Melián, C. J., & Gonzalez, A. (2019). Principles of ecology revisited: Integrating information and ecological theories for a more unified science. *Frontiers in Ecology and Evolution*, 7(219). <https://www.frontiersin.org/articles/10.3389/fevo.2019.00219>
- Oien, I. J., Honza, M., Moksnes, A., & Roskaft, E. (1996). The risk of parasitism in relation to the distance from reed warbler nests to cuckoo perches. *The Journal of Animal Ecology*, 65(2), 147–153. <https://doi.org/10.2307/5717>
- R Core Team. (2023). R: A language and environment for statistical computing. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Rothstein, S. (2001). Relic behaviours, coevolution and the retention versus loss of host defences after episodes of avian brood parasitism. *Animal Behaviour*, 61, 95–107. <https://doi.org/10.1006/anbe.2000.1570>
- Rowell, T. A. A. D., Magrath, M. J. L., & Magrath, R. D. (2020). Predator-awareness training in terrestrial vertebrates: Progress, problems and possibilities. *Biological Conservation*, 252, 108740. <https://doi.org/10.1016/j.biocon.2020.108740>
- Scheiner, S. M., & Levis, N. A. (2021). The loss of phenotypic plasticity via natural selection: Genetic assimilation. In D. W. Pfennig, *Phenotypic plasticity & evolution*. CRC Press. <https://doi.org/10.1201/9780429343001>
- Smith, W. J., Rönkä, K., Bensch, S., Bergman, N. M., Dudaniec, R. Y., Eroukhanoff, F., Fossey, F., Hansson, B., Jezierski, M. T., Klun, E., Procházka, P., Sætre, C. L. C., Stokke, B. G., & Thorogood, R. (2025). Putative signatures of genomic adaptation to Cuckoo brood parasitism in reed warbler hosts across Europe. *BioRxiv*, 2025.01.06.631542. <https://doi.org/10.1101/2025.01.06.631542> bioRxiv.
- Soler, M. (2014). Long-term coevolution between avian brood parasites and their hosts. *Biological Reviews*, 89(3), 688–704. <https://doi.org/10.1111/brv.12075>
- Soler, M., & Møller, A. P. (1990). Duration of sympatry and coevolution between the great spotted cuckoo and its magpie host. *Nature*, 343(6260), 748–750. <https://doi.org/10.1038/343748a0>
- Stokke, B. G., Moksnes, A., Røskaft, E., Rudolfen, S., & Honza, M. (1999). Rejection of artificial cuckoo (*Cuculus canorus*) eggs in relation to variation in egg appearance among reed warblers (*Acrocephalus scirpaceus*). *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 266(1427), 1483–1488. <https://doi.org/10.1098/rspb.1999.0804>
- Stokke, B. G., Poláčiková, L., Dyrce, A., Hafstad, I., Moksnes, A., & Røskaft, E. (2010). Responses of reed warblers *Acrocephalus scirpaceus* to non-mimetic eggs of different sizes in a nest parasitism experiment. *Acta Ornithol*, 45(1), 98–104. <https://doi.org/10.3161/000164510x516137>
- Stokke, B. G., Takasu, F., Moksnes, A., & Røskaft, E. (2007). The importance of clutch characteristics and learning for antiparasite adaptations in hosts of avian brood parasites. *Evolution*, 61(9), 2212–2228. <https://doi.org/10.1111/j.1558-5646.2007.00176.x>
- Thompson, J. N. (200

- ing the breeding season, even when cuckoos are rare. *Frontiers in Ecology and Evolution*, 9, 547. <https://doi.org/10.3389/fevo.2021.725467>
- Trnka, A., & Požgayová, M. (2017). Does conspicuous colouration of Magpies *Pica pica* influence aggressive behaviour in nesting great reed warblers *Acrocephalus arundinaceus*? *Bird Study*, 64(1), 108–111. <https://doi.org/10.1080/00063657.2016.1273878>
- Welbergen, J. A., & Davies, N. B. (2008). Reed warblers discriminate cuckoos from sparrowhawks with graded alarm signals that attract mates and neighbours. *Animal Behaviour*, 76(3), 811–822. <https://doi.org/10.1016/j.anbehav.2008.03.020>
- Welbergen, J. A., & Davies, N. B. (2009). Strategic variation in mobbing as a front line of defense against brood parasitism. *Current Biology*, 19(3), 235–240. <https://doi.org/10.1016/j.cub.2008.12.041>
- Welbergen, J. A., & Davies, N. B. (2011). A parasite in wolf's clothing: Hawk mimicry reduces mobbing of cuckoos by hosts. *Behavioral Ecology*, 22(3), 574–579. <https://doi.org/10.1093/beheco/arr008>
- Welbergen, J. A., & Davies, N. B. (2012). Direct and indirect assessment of parasitism risk by a cuckoo host. *Behavioral Ecology*, 23(4), 783–789. <https://doi.org/10.1093/beheco/ars031>
- Whiten, A. (2021). The burgeoning reach of animal culture. *Science*, 372(6537), eabe6514. <https://doi.org/10.1126/science.abe6514>
- Yang, C., Huang, J., Liang, W., & Møller, A. P. (2021). Absence of anti-parasitic defenses in an Asian population of the magpie, a regular host of the great spotted cuckoo in Europe. *Current Zoology*, 67(3), 345–347. <https://doi.org/10.1093/cz/zoaa018>
- Yang, C., Wang, L., Cheng, S.-J., Hsu, Y.-C., Stokke, B. G., Roskaft, E., Moksnes, A., Liang, W., & Møller, A. P. (2015). Deficiency in egg rejection in a host species as a response to the absence of brood parasitism. *Behavioral Ecology*, 26(2), 406–415. <https://doi.org/10.1093/beheco/aru202>