



Entropy function from toric geometry

Antonio Amariti ^a, Ivan Garozzo ^{b,c,*}, Gabriele Lo Monaco ^{b,c}

^a INFN, Sezione di Milano, Via Celoria 16, I-20133 Milano, Italy

^b Dipartimento di Fisica, Università di Milano-Bicocca, Piazza della Scienza 3, I-20126 Milano, Italy

^c INFN, sezione di Milano-Bicocca, Piazza della Scienza 3, I-20126 Milano, Italy

Received 17 March 2020; received in revised form 21 September 2021; accepted 5 October 2021

Available online 11 October 2021

Editor: Clay Córdova

Abstract

It has recently been claimed that a Cardy-like limit of the superconformal index of 4d $\mathcal{N} = 4$ SYM accounts for the entropy function, whose Legendre transform corresponds to the entropy of the holographic dual AdS₅ rotating black hole. Here we study this Cardy-like limit for $\mathcal{N} = 1$ toric quiver gauge theories, observing that the corresponding entropy function can be interpreted in terms of the toric data. Furthermore, for some families of models, we compute the Legendre transform of the entropy function, comparing with similar results recently discussed in the literature.

© 2021 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

Contents

1. Introduction	2
2. The Cardy-like limit of toric quivers	4
3. The conifold	8
4. Other examples	13
4.1. SPP	14
4.2. \mathbb{F}_0	16
4.3. dP_1	17

* Corresponding author.

E-mail addresses: antonio.amariti@mi.infn.it (A. Amariti), ivangarozzo@gmail.com (I. Garozzo), g.lomonaco1@campus.unimib.it (G. Lo Monaco).

4.4.	dP ₂	18
4.5.	dP ₃	20
4.6.	(P)dP ₄	22
5.	Infinite families	23
5.1.	Y ^{pq}	24
5.2.	X ^{pq}	25
5.3.	L ^{pqr}	25
6.	Legendre transform and the entropy	26
6.1.	The Y ^{pp} family	27
6.2.	The L ^{aba} family	28
7.	Conclusions	29
	CRedit authorship contribution statement	30
	Declaration of competing interest	30
	Acknowledgements	30
	Appendix A. Saddle point analysis for the conifold at higher-rank	30
	References	33

1. Introduction

The possibility of counting black hole microstates using the CFT dual picture is one of the most attractive consequences of the AdS/CFT correspondence [1]. A recent result in this field is the relation between the entropy of AdS₅ rotating black holes [2–6] and the superconformal index (SCI) [7,8]. The black hole entropy is given by the Benekenstein-Hawking formula, $S_{BH} = \frac{A}{4G_5}$, being A the area of the black hole horizon and G_5 the five dimensional Newton constant. The problem has been for a long time how to take into account the gravitational exponential growing ensemble of states from the dual CFT perspective. The potential candidate, the SCI, corresponding to the partition function computed on the conformal boundary $S^3 \times \mathbb{R}$, led to a puzzle: the growth of asymptotic states was not large enough to reproduce the entropy obtained from the gravitational side; the final result was of order $\mathcal{O}(1)$ instead of the expected $\mathcal{O}(N^2)$ [7]. For long time it was believed that the main reason behind this phenomenon is that the index counts either bosonic or fermionic operators weighted by the fermion number operator $(-1)^F$, inducing a huge cancellation of states. However it has been recently pointed out that the problem actually comes from the fact that the fugacities were taken to be real. In spite of this, the saddle point approximation of the index is able to reproduce the predicted exponential growth of the states in the index when allowing for complex valued fugacities [9,10]. One may also perform some explicit calculations based on a different partition function, that can be obtained by manipulating the SCI. In this approach, the existence of a deconfinement transition due to the extended range of fugacities was observed in [10].

A breakthrough in the analysis has been recently given by [11], where the authors associated the black hole entropy to a CFT extremization problem. They focused on the maximally supersymmetric case with two angular momenta and three conserved global charges. Moreover, the authors conjectured an expression for the grand canonical BPS partition function such that the Legendre transform of its logarithm reproduces the entropy of the black hole (as in the cases of [1,12,13]). By reformulating the problem in term of a grand canonical BPS partition function, Z_{BPS} , they conjectured the black hole entropy as a Legendre transform of the logarithm of Z_{BPS} (as in the cases of [1,12,13]). A concrete proposal for such a BPS partition function has been ob-

tained on the field theory side [14]. Furthermore in [15] it was realized that Z_{BPS} can be obtained on the gravitational side by considering the complexified on-shell action.

Moreover the authors of [16] exploited a reformulation of the SCI of $4d \mathcal{N} = 1$ theories as a finite sum over the solution of the so-called Bethe Ansatz equation [17].

Using these ideas the authors of [9,14,16,18–20] have obtained the BPS entropy function $S_E(\Delta_1, \Delta_2, \Delta_3, \omega_1, \omega_2)$ of [11] from the SCI of $\mathcal{N} = 4$ SYM. At large N this function reads

$$S_E = -i\pi N^2 \frac{\Delta_1 \Delta_2 \Delta_3}{\omega_1 \omega_2} \tag{1.1}$$

where Δ_I and ω_a are the fugacities conjugated to the charges Q_I and J_a of the $SO(6)_R$ R-symmetry and the $SO(4) \subset SO(4, 2)$ conformal symmetry respectively. Furthermore these fugacities are constrained by the relation ¹ $\Delta_1 + \Delta_2 + \Delta_3 - \omega_1 - \omega_2 = 1$, that corresponds, on the supergravity dual, to a stability condition on the killing spinor [15].

A natural question regards the extension of this result to other families of $4d \mathcal{N} = 1$ SCFT with an holographic dual description. Recent attempts in this direction has been given in [18], for the case of necklace $\mathcal{N} = 2$ models, in [20] for the case of Y^{pp} family and in [14] for more general classes of superconformal quivers. In all these cases the authors considered a subgroup of the full global symmetry and found interesting extensions of the results, showing also that the Legendre transform led to the expected entropy of the dual black hole.

In this paper we focus on infinite families of models, denoted as toric quiver gauge theories, that include the cases considered so far. These models describe the low energy dynamics of a stack of N D3 branes probing the tip of a toric cone over a five dimensional Sasaki-Einstein manifold. We study the large N index in the Cardy-like limit with complex fugacities discussed above and we give evidences of a general relation of the form

$$S_E = -i\pi N^2 \frac{C_{IJK} \Delta_I \Delta_J \Delta_K}{6 \omega_1 \omega_2} \tag{1.2}$$

where the fugacities Δ_I are read from the toric data and they satisfy the constraint $\sum_{I=1}^d \Delta_I - \sum_{a=1}^2 \omega_a = 1$. This result has been already conjectured in [21,22], where it was proposed that the numerator of (1.2) has the functional structure of the conformal anomaly of the 4d theory extracted from the gravitational (or geometric) data. The coefficients C_{IJK} in (1.2) correspond to the Chern-Simons couplings of the holographic dual gravitational description. Under the AdS/CFT correspondence they are associated to the triangle anomalies of the SCFT as shown in [23].

The paper is organized as follows. In section 2 we review the main aspects of our calculation focusing on the Cardy-like limit of the superconformal index and on the relation between the toric data and the global symmetries of the dual field theory. In section 3 we study the case of the conifold, computing the Cardy-like limit of the SCI and giving some evidences for the general conjecture on the behavior of the gauge holonomies at the saddle point. In section 4 we study other simple examples of toric quiver gauge theories, showing the validity of (1.2) for each case. In section 5 we focus on some infinite families, Y^{pq} , L^{pqr} and X^{pq} theories, and also in these cases we give evidences of (1.2). In section 6 we discuss the Legendre transform of the formula for the entropy of $\mathcal{N} = 2$ necklace quivers and for quivers in the Y^{pp} family. In both cases we extend the results already computed in the literature by turning on all the global symmetries. In section 7 we conclude, discussing possible future lines of research.

¹ See [15] for a detailed explanation on the sign.

2. The Cardy-like limit of toric quivers

In this section we explain the general aspects of the calculation of the Cardy-like limit of the SCI with complex fugacities for toric quiver gauge theories.

Toric quiver gauge theories describe the low energy dynamics of a stack of N D3 branes probing the tip of a toric cone over a five dimensional Sasaki-Einstein manifold. The toric data describing the singularity can be associated with the field theory data obtained by studying the moduli space [24,25]. In order to obtain these data starting from a gauge theory one has to first embed the quiver in a two dimensional torus. In this way one obtains a planar diagram, that can be transformed in a dimer, by exchanging faces and nodes. On this structure one defines the notion of perfect matching (PM): the PMs are collections of fields that represent all the possible dimer covers. By weighting the PMs with respect to the one-cycles of the first homology group of the torus one defines two possible intersection numbers for each PM. One can then assign a vector $V_I = (\cdot, \cdot, 1)$ to each PM, such that the first two entries are the intersection numbers discussed above and the last one is fixed to 1. The toric diagram corresponds to the convex integral polygon constructed from the V_I vectors. Using this construction it is possible to assign a basis of global symmetries of the quiver directly from the toric diagram. This consists of assigning a $U(1)_I$ symmetry, denoted as Q_I , to each external point of the toric diagram. One can construct the R -symmetry and the flavor (and baryonic symmetries) by combining these $U(1)_I$ as follows.

First one assigns a set of coefficients $\mathbf{a}_I \equiv \{\mathbf{a}_I^{(R)}, \mathbf{a}_I^{(i)}\}$ to each PM. Then it is necessary to impose the constraints $\sum_{I=1}^d \mathbf{a}_I^{(R)} = 2$ and $\sum_{i=1}^d \mathbf{a}_I^{(i)} = 0 \forall i$, where d is the number of external points in the toric diagram. The charges of the fields are associated to the ones of the PM with the prescription of [26]. Furthermore, the areas of the triangles obtained by connecting three external points of the toric diagram coincide with the triangular anomalies between the three $U(1)_I$ symmetries associated to such points [23]

$$\frac{N^2}{2} |\det(V_I, V_J, V_K)| = \text{Tr}(Q_I Q_J Q_K) \equiv N^2 C_{IJK} \tag{2.1}$$


As an example let us discuss the simplest toric quiver gauge theory, corresponding to $\mathcal{N} = 4$ $SU(N)$ SYM. We look at this theory as an $\mathcal{N} = 1$ theory with superpotential

$$W = \Phi_1[\Phi_2, \Phi_3] \tag{2.2}$$

where Φ_I are in the adjoint gauge group. In this case we have three $U(1)$ trial R-symmetries, denoted as $2U(1)_{1,2,3}$, and each $U(1)_I$ assigns charge 1 to the I -th field and zero to the others:

	$U(1)_1$	$U(1)_2$	$U(1)_3$	
Φ_1	1	0	0	
Φ_2	0	1	0	
Φ_3	0	0	1	(2.3)

There are three PM as shown in (2.4), corresponding to the three fields Φ_I .



The toric diagram is then generated by the three vectors

$$V_1 = (0, 0, 1), \quad V_2 = (0, 1, 1), \quad V_3 = (1, 0, 1) \tag{2.5}$$

The three trial R -symmetries are associated to the three corners of the toric diagram generated by three vectors in (2.5). Combining these symmetries we can extract the $U(1)_R$ symmetry and the other two flavor symmetries associated to the Cartan of the $SU(4)_R$ symmetry group of $\mathcal{N} = 4$ SYM. For example we can choose as an R -symmetry the combination $\frac{2}{3}(U(1)_1 + U(1)_2 + U(1)_3)$. In this case this assigns R -charge $\frac{2}{3}$ to each fields and it gives accidentally also the exact R -symmetry of the model. More generally the exact R -symmetry is given by a -maximization [27], where the conformal anomaly in this language corresponds to the function [26,28]

$$a_{\text{geom}} \propto C_{IJK} \mathbf{a}_I^{(R)} \mathbf{a}_J^{(R)} \mathbf{a}_K^{(R)} \tag{2.6}$$

The other two global symmetries can be obtained by the combinations $U(1)'_1 = U(1)_1 - U(1)_3$ and $U(1)'_2 = U(1)_2 - U(1)_3$. In this way we assign the charges as

	$U(1)'_1$	$U(1)'_2$	$U(1)_R$
Φ_1	1	0	$\frac{2}{3}$
Φ_2	0	1	$\frac{2}{3}$
Φ_3	-1	-1	$\frac{2}{3}$

(2.7)

Using these ideas one can read the parameterization of the global symmetries entering in the superconformal index from the toric diagram. We just have to linearly combine the $U(1)_I$ symmetries in order to obtain the non- R , either flavor or baryonic symmetries. In the following we will choose the $d - 1$ combinations $U(1)_i - U(1)_d$, with $i = 1, \dots, d - 1$ as our basis of non R -global symmetries. Furthermore the R -symmetry (not necessarily the exact one) will correspond to the combination $\frac{2}{d} \sum_{I=1}^d U(1)_I$. Using this basis of charges and symmetries we can write the SCI of a toric quiver gauge theory in the form

$$I = \text{Tr}_{\text{BPS}} (-1)^F e^{-\beta H} p^{J_1 + \frac{1}{d} \sum_{i=1}^d Q_i} q^{J_2 + \frac{1}{d} \sum_{i=1}^d Q_i} \prod_{i=1}^{d-1} u_i^{Q_i - Q_d} \tag{2.8}$$

Then we shift the chemical potentials $u_i \rightarrow u_i(pq)^{-\frac{1}{d}}$ obtaining

$$I = \text{Tr}_{\text{BPS}} (-1)^F p^{J_1} q^{J_2} (pq)^{\frac{1}{d} \sum_{i=1}^d Q_i} \prod_{i=1}^{d-1} u_i^{Q_i - Q_d} (pq)^{\frac{Q_d - Q_i}{d}} \tag{2.9}$$

Then by defining $p = e^{2\pi i \omega_1}$, $q = e^{2\pi i \omega_2}$, $u_i = e^{2\pi i \Delta_i}$ (for $i = 1 \dots d - 1$) and $(-1)^F = e^{2i\pi Q_d}$ (using the fact that this is an R -symmetry as well) we can express the index as

$$I = \text{Tr}_{\text{BPS}} e^{2\pi i \omega_1 J_1} e^{2\pi i \omega_2 J_2} \prod_{I=1}^d e^{2\pi i \Delta_I Q_I} \tag{2.10}$$

with the constraint

$$\sum_{I=1}^d \Delta_I - \omega_1 - \omega_2 = 1 \tag{2.11}$$

The Cardy limit of the SCI [29] and its generalization in [30,31] are obtained by shrinking the circle on which the index is defined as a partition function $S^3 \times S^1$. This can be done with complex fugacities by taking the limit $|\omega_1|, |\omega_2| \rightarrow 0$ [9,18,19]

$$\lim_{|\omega_1|, |\omega_2| \rightarrow 0} I \simeq e^{-\frac{i\pi(\omega_1 + \omega_2)}{12\omega_1\omega_2} \text{Tr} R} \int \prod_{i=1}^{\text{rank } \mathcal{G}} da_i e^{V(a)} \tag{2.12}$$

where

$$V(a) = \frac{i\pi}{2\omega_1\omega_2} \left(V_1(a)(\omega_1 + \omega_2) + \frac{V_2(a)}{3} \right) \tag{2.13}$$

In this formula $\text{rank } \mathcal{G}$ refers to the dimension of the maximal abelian torus of the gauge group, that is parameterized by the gauge holonomies $e^{2\pi i a_i}$. The functions V_1 and V_2 are

$$\begin{aligned} V_1(a) &= \sum_{k=1}^G \sum_{m,n=1}^N \theta(a_m^{(k)} - a_n^{(k)}) + \sum_{k \rightarrow k'} \sum_{m,n=1}^N (R_{kk'} - 1) \theta(a_m^{(k)} - a_n^{(k')}) + \sum_{i=1}^{d-1} q_{kk'}^i \Delta_i \\ V_2(a) &= - \sum_{k \rightarrow k'} \sum_{m,n=1}^N \kappa(a_m^{(k)} - a_n^{(k')}) + \sum_{i=1}^{d-1} q_{kk'}^i \Delta_i \end{aligned} \tag{2.14}$$

Let us explain these formulas. In the first line G refers to the number of gauge groups. It is obtained from a toric diagram by the formula $G = 2\mathcal{I} + d - 2$, where \mathcal{I} is the number of internal points. In the formula for V_1 there are two contributions, the first comes from the vector multiplets while the second from each bifundamental multiplet connecting the k -th to the k' -th node. Adjoint matter fields have $k = k'$. The function V_2 takes contributions only from the matter fields. Each matter field has R -charge $R_{kk'}$ and global charges $q_{kk'}^i$. The fugacities Δ_i are the ones defined above. In this paper we will always refer to $SU(N)$ gauge theories, and this will impose the constraint $\sum_{i=1}^N a_m^{(k)} = 0$. Moreover, the functions $\theta(x)$ and $\kappa(x)$ are given by

$$\theta(x) = \{x\}(1 - \{x\}), \quad \kappa(x) = \{x\}(1 - \{x\})(1 - 2\{x\}), \tag{2.15}$$

with the fractional part $\{x\} = x - [x]$, and can be rewritten as

$$\theta(x) = |x| - x^2, \quad \kappa(x) = 2x^3 - 3x|x| + x \tag{2.16}$$

for $|x| \leq 1$. The next step consists of evaluating the integral (2.12). We start by ignoring the contribution of $\text{Tr } R$. This is because in this paper we always consider toric quivers with a weakly coupled gravity dual. It follows that the gravitational anomaly, proportional to $\text{Tr } R$, is order $\mathcal{O}(1)$, while we restrict to the leading large N contribution of the Cardy-like limit of the index. Furthermore, we focus on the regime $\text{Re} \left(\frac{i}{\omega_1\omega_2} \right) > 0$. This is the regime discussed in [9,14,18–20] where it was shown that there is a saddle point at vanishing holonomies when considering

$\mathcal{N} = 4$ SYM. Computing the Cardy-like limit of the SCI using the charges (2.7) we obtain the entropy function

$$S_E = -i\pi N^2 \frac{\Delta_1 \Delta_2 \Delta_3}{\omega_1 \omega_2} \tag{2.17}$$

where with a slight abuse of notation we defined a new set of charges $\Delta_I = \frac{2}{\omega_1 + \omega_2 - 1} \{\Delta_I\}$. The new charges Δ_I appearing in (2.17) are associated to the symmetries $U(1)_I$ with $I = 1, 2, 3$ and the result holds provided we impose the constraint $\Delta_1 + \Delta_2 + \Delta_3 - \omega_1 - \omega_2 = 1$.

Here we study more general classes of quiver gauge theories. The first problem corresponds to find arguments in favor of the existence on an universal saddle point with vanishing holonomies as already discussed in [9,14,18,19]. Here we will confirm this expectation, observing in examples on increasing complexity that there is always a regime of fugacities that allows the existence of such a universal saddle.

Furthermore in each example we compute the Cardy-like limit of the index at large N , and we observe that it is controlled by the function

$$S_E = -i\pi N^2 \frac{C_{IJK} \Delta_I \Delta_J \Delta_K}{6 \omega_1 \omega_2} \tag{2.18}$$

where Δ_I are the fugacities appearing in (2.10) and the constraint (2.11) is imposed. This result can be proved by considering the relation obtained in [15,20] for the Cardy-like limit of the SCI of a generic $\mathcal{N} = 1$ gauge theory in presence of flavor fugacities. The relation is

$$S_E = -i\pi N^2 \frac{\text{Tr}(\Delta R + x_i F_i)^3}{6 \omega_1 \omega_2} \tag{2.19}$$

that holds imposing the constraint

$$2\Delta - \omega_1 - \omega_2 = 1 \tag{2.20}$$

where Δ represents the R -symmetry fugacity, while x_i are the flavor symmetry fugacities. We can express the R -symmetry and the flavor symmetries F_i as

$$R = \sum_{I=1}^d Q_I \mathbf{a}_I^{(R)}, \quad F_i = \sum_{I=1}^d Q_I \mathbf{a}_I^{(i)} \quad \forall i \tag{2.21}$$

with the constraints $\sum_{I=1}^d \mathbf{a}_I^{(R)} = 2$ and $\sum_{I=1}^d \mathbf{a}_I^{(i)} = 0, \forall i$. The combination appearing in (2.19) can be expressed in terms of these redefinitions as

$$\Delta R + x_i F_i = \sum_{I=1}^d Q_I \left(\Delta \mathbf{a}_I^{(R)} + \sum_{i=1}^{d-1} x_i \mathbf{a}_I^{(i)} \right) \equiv \sum_{I=1}^d Q_I \Delta_I \tag{2.22}$$

where in the last equality we defined the new fugacities Δ_I . These fugacities are constrained as

$$\sum_{I=1}^d \Delta_I = \sum_{I=1}^d \left(\Delta \mathbf{a}_I^{(R)} + \sum_{i=1}^{d-1} x_i \mathbf{a}_I^{(i)} \right) = 2\Delta = \omega_1 + \omega_2 + 1 \tag{2.23}$$

where in the last equality we used the constraint (2.20). In terms of the Q_I symmetries the entropy function reads

$$\begin{aligned}
 S_E &= -\frac{i\pi N^2}{6\omega_1\omega_2} \text{Tr} \left(\sum_{I=1}^d Q_I \Delta_I \right)^3 = -\frac{i\pi N^2}{6\omega_1\omega_2} \text{Tr} (Q_I Q_J Q_K) \Delta_I \Delta_J \Delta_K \\
 &= -\frac{i\pi N^2}{6\omega_1\omega_2} C_{IJK} \Delta_I \Delta_J \Delta_K
 \end{aligned}
 \tag{2.24}$$

with the constraint (2.23) and the last equality follows from the relation (2.1). We are going to verify (2.24) in the rest of the paper by explicitly studying the Cardy-like limit of the SCI for many toric quiver gauge theories.

3. The conifold

The conifold represents an ideal arena where testing, at finite rank, the Cardy formula and show the agreement with our general proposal (2.18), once the charges are parametrized from a geometric point of view.

The theory we are going to study has been proposed originally in [32] as the theory living on a stack of N D3-branes probing the tip of the conical singularity $xy - zt = 0$; taking the near-horizon limit, the theory turns out to be holographically dual to $\text{AdS}_5 \times T^{1,1}$ background where $T^{1,1}$ is what is properly named conifold. $T^{1,1}$ can be seen as an $U(1)$ fibration over $\mathbb{CP}^1 \times \mathbb{CP}^1$ with the $U(1)$ fiber playing the role of Reeb vector; the manifold admits a Sasaki-Einstein structure and has the topology of $S^2 \times S^3$. More importantly for our discussion, $T^{1,1}$ is also toric, with the toric diagram identified by the following four vectors:

$$V_1 = (1, 0, 0), \quad V_2 = (1, 1, 0), \quad V_3 = (1, 1, 1), \quad V_4 = (1, 0, 1).
 \tag{3.1}$$

The dual theory can be summarized by the following quiver and superpotential:

$W \propto \epsilon_{ij} \epsilon_{kl} \text{Tr} (A^i B^k A^j B^l)$

$\tag{3.2}$

The isometries of $T^{1,1}$ suggest the global symmetries of the CFT: a $U(1)_R$ factor (the R -symmetry generated by action of the Reeb vector) and two $SU(2)$ factors to be identified with the isometries of $\mathbb{CP}^1 \times \mathbb{CP}^1$; finally, we need to add a $U(1)_B$ baryonic symmetry associated to the unique non-trivial three-cycle of the geometry². The charges of the fields under $U(1)_R$, $U(1)_B$ and (a combination of) the Cartan generators $U(1)_{1,2}$ of the $SU(2)$ factors are summarized in the table below:

	$U(1)_R$	$U(1)_B$	$U(1)_1$	$U(1)_2$
A_1	1/2	1	1	1
A_2	1/2	1	-1	-1
B_1	1/2	-1	1	-1
B_2	1/2	-1	-1	1

$\tag{3.3}$

² As we said, the topology of $T^{1,1}$ is actually the same of $S^2 \times S^3$. The unique three-cycle can be understood as this S^3 .

We will turn on fugacities $\Delta_{F_{1,2}}$ for the flavor symmetries $U(1)_{1,2}$ and fugacity Δ_B for the baryonic symmetry $U(1)_B$.

We want to study now the Cardy formula in the rank-1 case, *i.e.* for $SU(2)$ gauge groups; in fact, a crucial point is understanding the behavior of the saddle points with respect to the holonomies. In low-rank cases it is possible to prove the main conjecture, *i.e.* it is possible to find charge configurations where the dominant saddle-point contribution is unique and corresponds to putting to zero all the holonomies; then, we will generalize to arbitrary N assuming the conjecture to be true at any rank. This fits with the discussions on the existence of such and universal saddle point in [14,19,20]. Moreover, we want to show that the choice of range for the fugacities is crucial and not all of them are suitable for our purpose. Let us start evaluating:

$$V_2 = - \sum_{m,n=1}^N \left(\kappa \left[a_m^{(1)} - a_n^{(2)} + \Delta_{F_1} + \Delta_{F_2} + \Delta_B \right] + \kappa \left[a_m^{(1)} - a_n^{(2)} - \Delta_{F_1} - \Delta_{F_2} + \Delta_B \right] + \kappa \left[a_m^{(2)} - a_n^{(1)} + \Delta_{F_1} - \Delta_{F_2} - \Delta_B \right] + \kappa \left[a_m^{(2)} - a_n^{(1)} - \Delta_{F_1} + \Delta_{F_2} - \Delta_B \right] \right), \tag{3.4}$$

where $a_m^{(1)}$ and $a_m^{(2)}$ are the holonomies for the first and second gauge group respectively. In the $SU(2)$ case we also need to enforce the condition $a_2^{(k)} = -a_1^{(k)}$ so that we are actually left with just two independent variables; in the following it will be more convenient to use the combinations:

$$a_{\pm} = a_1^{(1)} \pm a_1^{(2)}. \tag{3.5}$$

After some algebraic manipulation, (3.4) can be reduced to

$$V_2 = - (f[a_+] + f[a_-]), \tag{3.6}$$

where the function f is defined as follows:

$$f[x] = \kappa[x + \Delta_{F_1} + \Delta_{F_2} + \Delta_B] - \kappa[x - \Delta_{F_1} - \Delta_{F_2} - \Delta_B] + \kappa[x - \Delta_{F_1} - \Delta_{F_2} + \Delta_B] - \kappa[x - \Delta_{F_1} - \Delta_{F_2} - \Delta_B] + \kappa[x + \Delta_{F_1} - \Delta_{F_2} - \Delta_B] - \kappa[x - \Delta_{F_1} + \Delta_{F_2} + \Delta_B] + \kappa[x - \Delta_{F_1} + \Delta_{F_2} + \Delta_B] - \kappa[x + \Delta_{F_1} - \Delta_{F_2} + \Delta_B]. \tag{3.7}$$

Extremizing V_2 amounts to find extrema of $f[x]$. Observe that this function is invariant under permutations of fugacities Δ_{F_1} , Δ_{F_2} and Δ_B . It follows that we can choose an ordering of the charges without loss of generality, let us say $0 \leq \Delta_{F_1} \leq \Delta_{F_2} \leq \Delta_B$. Furthermore, using the property $\kappa[x] = \kappa[x + 1]$ we can move to a region where $-1/2 \leq \Delta_{F_1} + \Delta_{F_2} + \Delta_B \leq 1/2$. We want to focus for simplicity on a particular “chamber”, where we fix $0 \leq \Delta_B, \Delta_{F_1}, \Delta_{F_2} \leq 1/2$; this choice almost fixes completely the chamber and an ordering for all possible combinations $\Delta_{F_1} \pm \Delta_{F_2} \pm \Delta_B$. We are left with two possibilities:

$$\Delta_{F_1} + \Delta_{F_2} \geq \Delta_B \quad \text{or} \quad \Delta_{F_1} + \Delta_{F_2} \leq \Delta_B. \tag{3.8}$$

Now we are able to analytically evaluate $f[x]$ in the “fundamental regions” $|x| < 1 - \Delta_{F_1} - \Delta_{F_2} + \Delta_B$ and $|x| < 1 + \Delta_{F_1} + \Delta_{F_2} - \Delta_B$ respectively, where we can use the simplified expression (2.16). We proceed to study the behavior of $f[x]$ and we will see that these regimes are physically different and do not share the same properties.




Fig. 1. Plot of $f[x]$ for $\{\Delta_{F_1}, \Delta_{F_2}, \Delta_B\} = \{0.04, 0.9, 0.27\}$

- $\Delta_B \geq \Delta_{F_1} + \Delta_{F_2}$: In this case the fundamental region is $|x| < 1 + \Delta_{F_1} + \Delta_{F_2} - B$ and $f[x]$ reads:

$$f[x] = \begin{cases} 48\Delta_{F_1}\Delta_{F_2}(2\Delta_B - 1) & 0 < x < x_1 \\ 6((x + \Delta_{F_1} + \Delta_{F_2} - \Delta_B)^2 + 8(2\Delta_B - 1)\Delta_{F_1}\Delta_{F_2}) & x_1 < x < x_2 \\ 24\Delta_{F_1}((4\Delta_{F_2} - 1)\Delta_B + x - \Delta_{F_2}) & x_2 < x < x_3 \\ 6(16\Delta_B\Delta_{F_1}\Delta_{F_2} - (x - \Delta_B - \Delta_{F_1} - \Delta_{F_2})^2) & x_3 < x < x_4 \\ 96\Delta_B\Delta_{F_1}\Delta_{F_2} & x > x_4 \end{cases} \quad (3.9)$$

where $x_1 = \Delta_B - \Delta_{F_1} - \Delta_{F_2}$, $x_2 = \Delta_B + \Delta_{F_1} - \Delta_{F_2}$, $x_3 = \Delta_B - \Delta_{F_1} + \Delta_{F_2}$, and $x_4 = \Delta_B + \Delta_{F_1} + \Delta_{F_2}$

We can observe that in a whole neighborhood of $x = 0$ the function is constant; thus, for vanishing holonomies, $f[x]$ exhibits a plateau of extrema, rather than a unique minimum or maximum, as we can see from its plot in Fig. 1. As a consequence, in order to evaluate the index, we should perform an integration over the whole plateaux, making the study harder. For this reason, we exclude this case from our analysis but we will comment more about this point in the conclusions.

- $\Delta_B \leq \Delta_{F_1} + \Delta_{F_2}$: In this case the fundamental domain is $|x| < 1 - \Delta_{F_1} - \Delta_{F_2} + \Delta_B$ and $f[x]$ reads:

$$f[x] = \begin{cases} 12(\Delta_B^2 + (\Delta_{F_1} - \Delta_{F_2})^2 - 2\Delta_B\Delta_{F_2} + \Delta_B\Delta_{F_1}(8\Delta_{F_2} - 2) + x^2) & 0 < x < x_1 \\ 6((x + \Delta_{F_1} + \Delta_{F_2} - \Delta_B)^2 - 8(1 - 2\Delta_B)\Delta_{F_1}\Delta_{F_2}) & x_1 < x < x_2 \\ 24\Delta_{F_1}(x - \Delta_{F_2} + \Delta_B(4\Delta_{F_2} - 1)) & x_2 < x < x_3 \\ 6(16\Delta_B\Delta_{F_1}\Delta_{F_2} - (x - \Delta_{F_1} - \Delta_{F_2} - \Delta_B)^2) & x_3 < x < x_4 \\ 96\Delta_B\Delta_{F_1}\Delta_{F_2} & x > x_4 \end{cases} \quad (3.10)$$





Fig. 2. Plot of $f[x]$ for $\{\Delta_{F_1}, \Delta_{F_2}, \Delta_B\} = \{0.05, 0.15, 0.19\}$

where $x_1 = \Delta_{F_1} + \Delta_{F_2} - \Delta_B$, $x_2 = \Delta_B + \Delta_{F_1} - \Delta_{F_2}$, $x_3 = \Delta_B - \Delta_{F_1} + \Delta_{F_2}$, and $x_4 = \Delta_B + \Delta_{F_1} + \Delta_{F_2}$. This time, we can observe that $f[x]$ is manifestly non-constant in a neighborhood of $x = 0$, where a unique minimum is located (see Fig. 2). Reminding that V_2 is actually, $-f[x] - f[y]$, we discover that this extremal point $a_+ = a_- = 0$, *i.e.* vanishing holonomies, dominates the Cardy-like limit in the regime:

$$\text{Re}\left(\frac{i}{\omega_1 \omega_2}\right) > 0. \tag{3.11}$$

We showed that not all the chambers lead to honest isolated extrema and a similar analysis must be always performed. A numerical analysis for higher ranks still validates the work-hypothesis of vanishing holonomies and from now on we will consider arbitrary rank N following this assumption³.

In order to exploit the geometric insight, from now on we prefer to take a suitable basis of field charges that are directly suggested by the toric diagram, as discussed in section 2; in this basis flavor and baryonic symmetries get mixed and can be considered on equal footing. We will label the three $U(1)$ global symmetries simply as $U(1)_{1,2,3}$ and the associated fugacities as $\Delta_{1,2,3}$; R -symmetry will be denoted by $U(1)_R$ instead. Following the discussion in section 2 we can parameterize the global charges from the geometry using perfect matchings.



(3.12)

³ In appendix A we perform a similar analysis for the conifold at rank 2; we show that it is reasonable to extend to this case the results of our rank-1 study

In this case there are four perfect matchings associated to the four external points of the toric diagram. They are listed in (3.12), where it is possible to observe that in this case each PM corresponds to a bifundamental chiral field. The charges of these fields can be parameterized in terms of the PMs as

$$\begin{array}{c|c|c|c}
 A_1 & B_1 & A_2 & B_2 \\
 \hline
 \mathbf{a}_1 & \mathbf{a}_2 & \mathbf{a}_3 & \mathbf{a}_4
 \end{array} \tag{3.13}$$

The charges of the fields with respect to the symmetries suggested by the geometric data can be taken then as follows

$$\begin{array}{c|cccc}
 & U(1)_1 & U(1)_2 & U(1)_3 & U(1)_R \\
 \hline
 A_1 & 1 & 0 & 0 & 1/2 \\
 A_2 & 0 & 0 & 1 & 1/2 \\
 \hline
 B_1 & 0 & 1 & 0 & 1/2 \\
 B_2 & -1 & -1 & -1 & 1/2
 \end{array} \tag{3.14}$$

We want to stress that, as in the $\mathcal{N} = 4$ case discussed in section 2, the R -symmetry that we consider here accidentally coincides with the exact R -symmetry at the conformal fixed point. However this will not be the case in the models that we are going to discuss in the next section. Observe that the new conjugated fugacities Δ_i can thought as linear combination of the flavor and baryonic ones:

$$\Delta_1 = \Delta_B + \Delta_{F_1} + \Delta_{F_2}, \quad \Delta_2 = -\Delta_B + \Delta_{F_1} - \Delta_{F_2}, \quad \Delta_3 = \Delta_B - \Delta_{F_1} - \Delta_{F_2}. \tag{3.15}$$

V_2 for arbitrary rank can be now expressed as:

$$\begin{aligned}
 V_2 = - \sum_{m,n=1}^N & \left(\kappa \left[a_m^{(1)} - a_n^{(2)} + \Delta_1 \right] + \kappa \left[a_m^{(2)} - a_n^{(1)} + \Delta_2 \right] + \right. \\
 & \left. \kappa \left[a_m^{(1)} - a_n^{(2)} + \Delta_3 \right] + \kappa \left[a_m^{(2)} - a_n^{(1)} - \Delta_1 - \Delta_2 - \Delta_3 \right] \right). \tag{3.16}
 \end{aligned}$$

In this basis we fix the fugacities such that $0 \leq \Delta_{1,2,3} < 1/2$ and $0 \leq \Delta_1 + \Delta_2 + \Delta_3 \leq 1/2$. In this chamber, $f[x]$ enjoys a local maximum for vanishing holonomies and thus V_2 exhibits a minimum⁴. The extremum dominates the Cardy-like limit if

$$\text{Re} \left(\frac{i}{\omega_1 \omega_2} \right) < 0. \tag{3.17}$$

⁴ The range we fixed for $\Delta_{1,2,3}$ leads to slightly different features with respect to the one we chose for $\Delta_{F_{1,2}}, \Delta_B$ in our previous discussion; in the former case, V_2 enjoys a local minimum while in the latter V_2 possesses a local maximum, in both cases for vanishing holonomies. For this reason, the two extrema dominates the Cardy-like limit in different regimes, (3.17) and (3.11) respectively. Both fugacity ranges can be chosen, up to minimal changes to be performed in going from one regime to the other, as carefully shown in [19].

In the fixed regime, we can evaluate the dominant saddle contribution:

$$\begin{aligned}
 V &= \frac{i\pi}{2\omega_1\omega_2} \left(\frac{V_2}{3} + (\omega_1 + \omega_2)V_1 \right) \Big|_{a_m^{(l)}=0} = \\
 &= \frac{\pi i N^2}{2\omega_1\omega_2} \left\{ (\omega_1 + \omega_2) \left(\Delta_3(\Delta_3 - 1) + \Delta_1(\Delta_2 + \Delta_3 - 1) + \Delta_1^2 + \Delta_2^2 + \Delta_2(\Delta_3 - 1) \right) + \right. \\
 &\quad \left. - \left(\Delta_2\Delta_3(\Delta_2 + \Delta_3 - 1) + \Delta_1^2(\Delta_2 + \Delta_3 + \Delta_1(\Delta_2 + \Delta_3 - 1)(\Delta_2 + \Delta_3)) \right) \right\}.
 \end{aligned}
 \tag{3.18}$$

This equation gets further simplified performing a suitable shift of the fugacities

$$\Delta_{1,2,3} \rightarrow \Delta_{1,2,3} - \frac{\omega_1 + \omega_2}{4}
 \tag{3.19}$$

and taking the leading order in the Cardy-like limit $|\omega_1|, |\omega_2| \rightarrow 0$; let us stress that the shift (3.19) is actually dictated by the geometry: as we will test for other toric models in the next sections, the fugacities get always shifted by $-\frac{\omega_1 + \omega_2}{d}$ where d is the number of external points in the toric diagram. After the shift, the entropy function can be expressed as

$$S_E = -\frac{i\pi N^2}{\omega_1\omega_2} (\Delta_1\Delta_2\Delta_3 + \Delta_1\Delta_2\Delta_4 + \Delta_1\Delta_3\Delta_4 + \Delta_2\Delta_3\Delta_4),
 \tag{3.20}$$

where we have defined:

$$\Delta_4 \equiv \omega_1 + \omega_2 + 1 - \Delta_1 - \Delta_2 - \Delta_3.
 \tag{3.21}$$

The entropy function (3.20) enjoys the expected scaling behavior $S_E \propto N^2$; moreover, it is in perfect agreement with our general proposal

$$S_E = -\frac{i\pi N^2}{6\omega_1\omega_2} C_{IJK} \Delta_I \Delta_J \Delta_K,
 \tag{3.22}$$

where I, J, K run from 1 to 4 and C_{IJK} is defined as in (2.1).

4. Other examples

In this section we test our proposal (2.18) in various cases of growing complexity. In each case we assign the charges using the prescription discussed in section 2 and we assume that the Cardy-like limit is dominated by a unique minimum where all the holonomies vanish. We have tested the last conjecture in the rank-1 cases of dP_1 , dP_2 , $(P)dP_4$ and \mathbb{F}_0 , finding evidence of its validity. In each case the minimum is found in a chamber where fugacities Δ_i of the $d - 1$ $U(1)$ global symmetries are taken such that:

$$0 \leq \Delta_i \leq \frac{1}{2} \quad \forall i, \quad 0 \leq \sum_{i=1}^{d-1} \Delta_i \leq 1.
 \tag{4.1}$$

Since in this range V_2 enjoys a minimum, we restrict to the regime (3.17).

As a general remark let us stress that some of the theories that we are going to discuss admit more Seiberg dual realizations, denoted as phases. We specify for each model the Seiberg phase that we focus on. Finally, observe that we are not necessarily fixing the R -charges of the fields at the conformal fixed point, but we refer to a trial R -current, using the uniform prescription for all the models under investigation, as explained in section 2.

4.1. SPP

The suspended pinch point (SPP) gauge theory corresponds to the near horizon limit of a stack of N D3 branes probing the tip of the conical singularity, $x^2y = wz$. This is the simplest example of a larger class of models, defined by the equation $x^a y^b = wz$, denoted as L^{aba} models. In the SPP case the toric Sasaki-Einstein base is described by the following vectors:

$$V_1 = (1, 1, 1), \quad V_2 = (1, 0, 1), \quad V_3 = (0, 0, 1), \quad V_4 = (2, 0, 1), \quad V_5 = (1, 0, 1). \quad (4.2)$$

The vector V_5 represents a point on the perimeter of the toric diagram and it turns out that two different perfect matchings can be associated to it and, consequently, we can get two different possible charge sets. Following the prescription of [26] we can associate a non vanishing set of charges to just one of them.

The theory living on a stack of N D3-branes at the SPP conical singularity is described by the following quiver:



with superpotential

$$W = \text{Tr}[X_{21}X_{12}X_{23}X_{32} - X_{32}X_{23}X_{31}X_{13} + X_{13}X_{31}\phi - X_{12}X_{21}\phi]. \quad (4.4)$$

Each X_{ij} transforms in the \mathbf{N} representation of the i node and in the $\overline{\mathbf{N}}$ of the j -th node; the field transforming in the adjoint of the first node is named, instead, ϕ . The charges of the fields can be parameterized in terms of the PMs using the assignation

ϕ	X_{12}	X_{21}	X_{23}	X_{32}	X_{31}	X_{13}
$\mathbf{a}_1 + \mathbf{a}_2$	\mathbf{a}_4	$\mathbf{a}_3 + \mathbf{a}_5$	\mathbf{a}_2	\mathbf{a}_1	$\mathbf{a}_4 + \mathbf{a}_5$	\mathbf{a}_3

(4.5)

It follows that the charge assignment for $U(1)_R$ and the extra four $U(1)_i$ global can be taken as follows:

	$U(1)_1$	$U(1)_2$	$U(1)_3$	$U(1)_4$	$U(1)_R$
ϕ	1	1	0	0	4/5
X_{12}	0	0	0	1	2/5
X_{21}	-1	-1	0	-1	4/5
X_{23}	0	1	0	0	2/5
X_{32}	1	0	0	0	2/5
X_{31}	-1	-1	-1	0	4/5
X_{13}	0	0	1	0	2/5

(4.6)

Observe that as usual we took a combination of $U(1)$ natural for toric geometry and we have not done any distinction between flavor and baryonic symmetries. We denote Δ_i the fugacity associated to $U(1)_i$. With this assignment, V_2 admits a minimum for vanishing holonomies in a chamber where $0 \leq \Delta_i \leq 1/2$ for each $U(1)_i$ and $0 < \Delta_1 + \Delta_2 + \Delta_3 + \Delta_4 < 1$; we can evaluate

$$V = \frac{i\pi}{2\omega_1\omega_2} \left(V_1(\omega_1 + \omega_2) + \frac{V_2}{3} \right) \tag{4.7}$$

and, after performing a shift of the charges by a factor $-\frac{\omega_1+\omega_2}{5}$ and taking the leading order in $|\omega_1|, |\omega_2| \rightarrow 0$, we get:

$$V_{\text{leading}} = \frac{i\pi N^2}{\omega_1\omega_2} \left(\Delta_1((\Delta_2 + \Delta_3)(\Delta_2 + \Delta_3 - \omega_1 - \omega_2 - 1) + \Delta_4^2 - \Delta_4(1 + \omega_1 + \omega_2 - 2\Delta_2)) + \Delta_2(\Delta_3^2 - \Delta_3(1 + \omega_1 + \omega_2 - \Delta_2) + \Delta_4(\Delta_2 + \Delta_4 - 1 - \omega_1 - \omega_2)) + \Delta_1^2(\Delta_2 + \Delta_3 + \Delta_4) \right). \tag{4.8}$$

If we now define a new constrained fugacity:


$$\Delta_5 = 1 + \omega_1 + \omega_2 - \Delta_1 - \Delta_2 - \Delta_3 - \Delta_4 \tag{4.9}$$

we obtain the following expression for the entropy:

$$S_E = -\frac{i\pi N^2}{\omega_1\omega_2} (\Delta_1\Delta_2\Delta_3 + \Delta_1\Delta_2\Delta_4 + 2\Delta_1\Delta_3\Delta_4 + 2\Delta_2\Delta_3\Delta_4 + \Delta_1\Delta_2\Delta_5 + \Delta_1\Delta_3\Delta_5 + \Delta_2\Delta_3\Delta_5 + \Delta_1\Delta_4\Delta_5 + \Delta_2\Delta_4\Delta_5). \tag{4.10}$$

This result is in agreement with our expectation from toric geometry, encoded in formula (2.18).

Finally, as discussed at the beginning of this section, the SPP singularity can be thought as a particular case of a larger class of toric models, denoted as L^{aba} , for $a = 1, b = 2$. The toric diagram of an L^{aba} singularity is depicted in (4.11).



In this case there are $a + b$ gauge groups, and two flavor symmetries and $a + b - 1$ non anomalous baryonic symmetries. This huge amount of baryonic symmetries reflects in the toric diagram onto the large number of external point lying on the perimeter. Each of these points contribute with triangle areas to reproduce the correct entropy function, following the general prescription (2.18). Observe that for $a = 0$ the models become $\mathcal{N} = 2$ necklace quivers, corresponding to \mathbb{Z}_b orbifolds of $\mathcal{N} = 4$ SYM. The entropy for these models has been studied in [18], by turning off the baryonic fugacities. Here in section 6 we will study the most general situation.

4.2. \mathbb{F}_0

The complex cone over the first Hirzebruch surface \mathbb{F}_0 is a \mathbb{Z}_2 orbifold of the conifold; the toric diagram is parametrized by the four vectors

$$V_1 = (0, 0, 1), \quad V_2 = (1, 0, 1), \quad V_3 = (0, 2, 1), \quad V_4 = (-1, 2, 1). \tag{4.12}$$

The corresponding theory in its phase I is described by the following quiver and superpotential:

$$W = \epsilon_{\alpha\beta\gamma\delta} \text{Tr}[X_{12}^{(\alpha)} X_{34}^{(\beta)} X_{23}^{(\gamma)} X_{41}^{(\delta)}]. \tag{4.13}$$

One can assign charges to the fields in the theory directly from the geometry. The charges of the fields can be parameterized in terms of the PMs using the assignation

X_{12}	X_{34}	X_{23}	X_{41}
\mathbf{a}_1	\mathbf{a}_2	\mathbf{a}_3	\mathbf{a}_4

(4.14)

The model has three global $U(1)$ symmetries in addition to one $U(1)_R$ and in this case one gets the following global charges

	multiplicity	$U(1)_1$	$U(1)_2$	$U(1)_3$	$U(1)_R$
X_{12}	2	1	0	0	1/2
X_{23}	2	0	1	0	1/2
X_{34}	2	0	0	1	1/2
X_{41}	2	-1	-1	-1	1/2

(4.15)

We now compute the Cardy-like limit of the superconformal index for this theory; if we denote the fugacities for the symmetries $U(1)_{1,2,3}$ as $\Delta_{1,2,3}$ respectively, the expression that we find after shifting each fugacity by $-\frac{\omega_1+\omega_2}{4}$ is

$$V_{\text{leading}} = -\frac{2\pi i N^2}{\omega_1 \omega_2} \left(\Delta_2(1 + \omega_1 + \omega_2 - \Delta_2 - \Delta_3)\Delta_3 - \Delta_1^2(\Delta_2 + \Delta_3) - \Delta_1(\Delta_2 + \Delta_3)(1 + \omega_1 + \omega_2 - \Delta_2 - \Delta_3) \right) \tag{4.16}$$

The entropy function in this case can be written as:

$$S_E(\Delta, \omega) = -\frac{2\pi i N^2}{\omega_1 \omega_2} (\Delta_1 \Delta_2 \Delta_3 + \Delta_1 \Delta_2 \Delta_4 + \Delta_1 \Delta_3 \Delta_4 + \Delta_2 \Delta_3 \Delta_4), \tag{4.17}$$

and it exactly reproduces (4.16) by using the constraint

$$\Delta_4 = 1 + \omega_1 + \omega_2 - \Delta_1 - \Delta_2 - \Delta_3. \tag{4.18}$$

Observe that the entropy function just obtained is twice the one for the conifold, as one should expect from the fact that we are dealing with a \mathbb{Z}_2 orbifold of the latter⁵.

4.3. dP_1

Let us consider the theory arising from a stack of N D3 branes at the tip of the complex Calabi-Yau cone whose base is the first del Pezzo surface. The toric diagram is generated by

$$V_1 = (1, 0, 1), \quad V_2 = (0, 1, 1), \quad V_3 = (-1, 0, 1), \quad V_4 = (-1, -1, 1). \tag{4.19}$$

The corresponding quiver is as follows



and the superpotential for this theory reads

$$W = \epsilon_{\alpha\beta} \text{Tr}[X_{34}^{(\alpha)} X_{41}^{(\beta)} X_{13} - X_{34}^{(\alpha)} X_{23}^{(\beta)} X_{42} + X_{12} X_{34}^{(3)} X_{41}^{(\alpha)} X_{23}^{(\beta)}]. \tag{4.21}$$

The charges of the fields can be parameterized in terms of the PMs using the assignation

$X_{34}^{(3)}, X_{13}, X_{42}$	$X_{23}^{(1)}, X_{41}^{(1)}$	X_{12}	$X_{23}^{(2)}, X_{41}^{(2)}$	$X_{34}^{(1)}$	$X_{34}^{(2)}$
\mathbf{a}_1	\mathbf{a}_2	\mathbf{a}_3	\mathbf{a}_4	$\mathbf{a}_2 + \mathbf{a}_3$	$\mathbf{a}_3 + \mathbf{a}_4$

(4.22)

The charges for $U(1)_R$ and the three global $U(1)$ symmetries of the model coming from the perfect matching are the following

⁵ To be more precise, the entropy function reproduces twice the conifold one because the orbifold action does not introduce new singularities or, equivalently, new symmetries. A non-chiral \mathbb{Z}_2 orbifold of the conifold like the L^{222} model does not have this property.

	$U(1)_1$	$U(1)_2$	$U(1)_3$	$U(1)_R$
X_{12}	0	0	1	1/2
$X_{23}^{(1)}$	0	1	0	1/2
$X_{23}^{(2)}$	-1	-1	-1	1/2
$X_{34}^{(1)}$	0	1	1	1
$X_{34}^{(2)}$	-1	-1	0	1
$X_{34}^{(3)}$	1	0	0	1/2
$X_{41}^{(1)}$	0	1	0	1/2
$X_{41}^{(2)}$	-1	-1	-1	1/2
X_{13}	1	0	0	1/2
X_{42}	1	0	0	1/2

(4.23)

The expression for the entropy function in this case gives:

$$S_E(\Delta, \omega) = -\frac{i\pi N^2}{\omega_1\omega_2} (2\Delta_1\Delta_2\Delta_3 + 2\Delta_1\Delta_2\Delta_4 + 2\Delta_1\Delta_3\Delta_4 + \Delta_2\Delta_3\Delta_4). \tag{4.24}$$

Again, the same result can be obtained by taking the Cardy-like limit of the superconformal index. The leading order of the function

$$V = \frac{i\pi N^2}{2\omega_1\omega_1} \left(V_1(\omega_1 + \omega_2) + \frac{V_2}{3} \right), \tag{4.25}$$

taken after shifting the charges by a factor $-\frac{\omega_1+\omega_2}{4}$, is given by

$$V_{\text{leading}} = \frac{i\pi N^2}{\omega_1\omega_2} (3\Delta_1\Delta_2(\Delta_1 + \Delta_2 - \omega_1 - \omega_2 - 1) - 2\Delta_1\Delta_3(1 + \omega_1 + \omega_2 - \Delta_1) - (1 + \omega_1 + \omega_2 - 4\Delta_1)\Delta_2\Delta_3 + \Delta_2^2\Delta_3 + (2\Delta_1 + \Delta_2)\Delta_3^2). \tag{4.26}$$

If we now take the expression of the entropy function (4.24) and impose the constraint on the fugacities

$$\Delta_4 = 1 + \omega_1 + \omega_2 - \Delta_1 - \Delta_2 - \Delta_3, \tag{4.27}$$

we obtain the expression (4.26) for the entropy function.

4.4. dP_2

The toric diagram for the complex cone over the dP_2 surface is generated by the following vectors

$$V_1=(1, 1, 1), \quad V_2=(0, 1, 1), \quad V_3=(-1, 0, 1), \quad V_4=(-1, -1, 1), \quad V_5 = (0, -1, 1). \tag{4.28}$$

The charges of the fields can be parameterized in terms of the PMs using the assignation

X_{13}	X_{24}	$X_{51}^{(1)}$	X_{23}	X_{41}	$X_{51}^{(2)}$	$X_{12}^{(2)}$	X_{45}	$X_{12}^{(1)}$	X_{35}	X_{34}
$\mathbf{a}_4 + \mathbf{a}_5$	\mathbf{a}_5	\mathbf{a}_5	\mathbf{a}_2	$\mathbf{a}_1 + \mathbf{a}_2$	$\mathbf{a}_2 + \mathbf{a}_3$	$\mathbf{a}_3 + \mathbf{a}_4$	\mathbf{a}_4	\mathbf{a}_1	\mathbf{a}_1	\mathbf{a}_3

(4.29)

The theory arising from a stack of N D3 branes put at the tip of this toric Calabi-Yau cone admits two phases. The phase I can be described by a quiver with five nodes



and superpotential

$$\begin{aligned}
 W = \text{Tr} [& X_{13} X_{34} X_{41} - X_{12}^{(2)} X_{24} X_{41} + X_{12}^{(1)} X_{24} X_{45} X_{51}^{(2)} - X_{13} X_{35} X_{51}^{(2)} \\
 & + X_{12}^{(2)} X_{23} X_{35} X_{51}^{(1)} - X_{12}^{(1)} X_{23} X_{34} X_{45} X_{51}^{(1)}].
 \end{aligned}$$

(4.31)

Here X_{ij} denotes a bifundamental field connecting the i -th and j -th nodes. The charges assigned to the various fields in the quiver directly from the perfect matchings are

	$U(1)_1$	$U(1)_2$	$U(1)_3$	$U(1)_4$	$U(1)_R$
X_{13}	-1	-1	-1	0	4/5
X_{24}	-1	-1	-1	-1	2/5
$X_{51}^{(1)}$	-1	-1	-1	-1	2/5
X_{23}	0	1	0	0	2/5
X_{41}	1	1	0	0	4/5
$X_{51}^{(2)}$	0	1	1	0	4/5
$X_{12}^{(2)}$	0	0	1	1	4/5
X_{45}	0	0	0	1	2/5
$X_{12}^{(1)}$	1	0	0	0	2/5
X_{35}	1	0	0	0	2/5
X_{34}	0	0	1	0	2/5

(4.32)

The entropy function obtained from toric geometry is:

$$\begin{aligned}
 S_E(\Delta, \omega) = & -\frac{i\pi N^2}{\omega_1 \omega_2} (\Delta_1 \Delta_2 \Delta_3 + 2\Delta_1 \Delta_2 \Delta_4 + 2\Delta_1 \Delta_3 \Delta_4 + \Delta_2 \Delta_3 \Delta_4 \\
 & + 2\Delta_1 \Delta_2 \Delta_5 + 3\Delta_1 \Delta_3 \Delta_5 + 3\Delta_1 \Delta_3 \Delta_5 + 2\Delta_2 \Delta_3 \Delta_5 \\
 & + 2\Delta_1 \Delta_4 \Delta_5 + 2\Delta_2 \Delta_4 \Delta_5 + \Delta_3 \Delta_4 \Delta_5)
 \end{aligned}$$

(4.33)

The leading order of the Cardy-like limit of the superconformal index gives, after shifting the charges by a factor $-\frac{\omega_1+\omega_2}{5}$

$$V_{\text{leading}} = \frac{i\pi}{\omega_1\omega_2} \left[(2\Delta_2^2(\Delta_3 + \Delta_4) + (\Delta_3\Delta_4 + 2\Delta_2(\Delta_3 + \Delta_4))(\Delta_3 + \Delta_4 - 1 - \omega_1 - \omega_2) + \Delta_1^2(2\Delta_2 + 3\Delta_3 + 2\Delta_4) + 2\Delta_1\Delta_2^2 - 3\Delta_1\Delta_3(1 + \omega_1 + \omega_2 - \Delta_3) - 2\Delta_1\Delta_2(1 + \omega_1 + \omega_1 - 3\Delta_3 - 2\Delta_4) - 2\Delta_1\Delta_4(1 + \omega_1 + \omega_2 - 2\Delta_3) + 2\Delta_1\Delta_4^2) \right]. \tag{4.34}$$

This is the same expression that one gets by taking (4.33) and using the constraint on the fugacities

$$\Delta_5 = -\Delta_1 - \Delta_2 - \Delta_3 - \Delta_4 + \omega_1 + \omega_2 + 1. \tag{4.35}$$

4.5. dP_3

The toric diagram for the Calabi-Yau cone over the dP_3 surface is generated by the following vectors

$$\begin{aligned} V_1 &= (1, 1, 1), & V_2 &= (0, 1, 1), & V_3 &= (-1, 0, 1), \\ V_4 &= (-1, -1, 1), & V_5 &= (0, -1, 1), & V_6 &= (1, 0, 1). \end{aligned} \tag{4.36}$$

The charges of the fields can be parameterized in terms of the PMs using the assignation

X_{12}	X_{13}	X_{23}	X_{24}	X_{34}	X_{35}	X_{45}	X_{46}	X_{56}	X_{51}	X_{61}	X_{62}
\mathbf{a}_6	$\mathbf{a}_2 + \mathbf{a}_3$	\mathbf{a}_5	$\mathbf{a}_1 + \mathbf{a}_2$	\mathbf{a}_4	$\mathbf{a}_1 + \mathbf{a}_6$	\mathbf{a}_3	$\mathbf{a}_5 + \mathbf{a}_6$	\mathbf{a}_2	$\mathbf{a}_4 + \mathbf{a}_5$	\mathbf{a}_1	$\mathbf{a}_3 + \mathbf{a}_4$

(4.37)

The theory associated to this cone admits three phases; in its phase I , it can be described by the following quiver:



with superpotential

$$W = \text{Tr} \left[X_{12}X_{24}X_{45}X_{51} - X_{24}X_{46}X_{62} + X_{23}X_{35}X_{56}X_{62} - X_{35}X_{51}X_{13} + X_{34}X_{46}X_{61}X_{13} - X_{12}X_{23}X_{34}X_{45}X_{56}X_{61} \right], \tag{4.39}$$

where X_{ij} denotes a bifundamental field connecting the i -th and j -th nodes. The charges assigned from the toric diagram to the various fields in the theory are

	$U(1)_1$	$U(1)_2$	$U(1)_3$	$U(1)_4$	$U(1)_5$	$U(1)_R$
X_{12}	-1	-1	-1	-1	-1	1/3
X_{13}	0	1	1	0	0	2/3
X_{23}	0	0	0	0	1	1/3
X_{24}	1	1	0	0	0	2/3
X_{34}	0	0	0	1	0	1/3
X_{35}	0	-1	-1	-1	-1	2/3
X_{45}	0	0	1	0	0	1/3
X_{46}	-1	-1	-1	-1	0	2/3
X_{56}	0	1	0	0	0	1/3
X_{51}	0	0	0	1	1	2/3
X_{61}	1	0	0	0	0	1/3
X_{62}	0	0	1	1	0	2/3

(4.40)

The entropy function for this theory is

$$S_E(\Delta, \omega) = -\frac{i\pi N^2}{\omega_1\omega_2} (\Delta_1\Delta_2\Delta_3 + 2\Delta_1\Delta_2\Delta_4 + 2\Delta_1\Delta_3\Delta_4 + \Delta_2\Delta_3\Delta_4 + 2\Delta_1\Delta_2\Delta_5 + 3\Delta_1\Delta_3\Delta_5 + 2\Delta_2\Delta_3\Delta_5 + 2\Delta_1\Delta_4\Delta_5 + 2\Delta_2\Delta_4\Delta_5 + \Delta_3\Delta_4\Delta_5 + \Delta_1\Delta_2\Delta_6 + 2\Delta_1\Delta_3\Delta_6 + 2\Delta_2\Delta_3\Delta_6 + 2\Delta_1\Delta_4\Delta_6 + 3\Delta_2\Delta_4\Delta_6 + 2\Delta_3\Delta_4\Delta_6 + \Delta_1\Delta_5\Delta_6 + 2\Delta_2\Delta_5\Delta_6 + 2\Delta_3\Delta_5\Delta_6 + \Delta_4\Delta_5\Delta_6).$$

(4.41)

In fact, evaluating V at the leading order in $|\omega_1|, |\omega_2| \rightarrow 0$ we get

$$V_{\text{leading}} = -\frac{i\pi N^2}{\omega_1\omega_2} (2\Delta_3\Delta_4(1+\omega_1+\omega_2-\Delta_3-\Delta_4) - 2\Delta_3^2\Delta_5 + 2\Delta_3\Delta_5(1+\omega_1+\omega_2-2\Delta_4) + \Delta_4\Delta_5(1+\omega_1+\omega_2-\Delta_4) + \Delta_5^2(2\Delta_3+\Delta_4) - \Delta_1^2(\Delta_2+2\Delta_3+2\Delta_4+\Delta_5) - \Delta_2^2(2\Delta_3+3\Delta_4+2\Delta_5) - 2\Delta_2\Delta_3^2 + 3\Delta_2\Delta_4(1+\omega_1+\omega_2-\Delta_4) + 2\Delta_2\Delta_3(1+\omega_1+\omega_2-3\Delta_4-2\Delta_5) + 2\Delta_2\Delta_5(1+\omega_1+\omega_2-2\Delta_4) - 2\Delta_2\Delta_5^2 - \Delta_1\Delta_2^2 - 2\Delta_1(\Delta_3+\Delta_4)(\Delta_3+\Delta_4-1-\omega_1-\omega_2) + \Delta_1\Delta_2(1+\omega_1+\omega_2-4\Delta_3-4\Delta_4-2\Delta_5) - \Delta_1\Delta_5^2 + \Delta_1\Delta_5(1+\omega_1+\omega_2-2\Delta_3-2\Delta_4)),$$

(4.42)

where we also shifted the fugacities by $-\frac{\omega_1+\omega_2}{6}$, consistently with the general prescription. Again, by imposing the constraint

$$\Delta_6 = -\Delta_1 - \Delta_2 - \Delta_3 - \Delta_4 - \Delta_5 + \omega_1 + \omega_2 + 1,$$

(4.43)

on (4.42) we obtain (4.41).

4.6. (P)dP₄

The fourth del Pezzo surface is defined as the blow-up of \mathbb{P}^2 at four generic⁶ points. The (complex) cone over it possesses a Calabi-Yau structure and the theory living on N D3-branes probing the conical singularity is known; however the superpotential of the dual gauge theory is such that no non-anomalous flavor symmetries except $U(1)_R$ are admitted, so that the model is non-toric. Blowing-up \mathbb{P}^2 at non-generic points, however, it is possible to build models where more symmetries are preserved.

One choice can be the toric model whose diagram is generated by the following vectors:

$$\begin{aligned} V_1 &= (0, 0, 1), V_2 = (1, 0, 1), V_3 = (2, 0, 1), V_4 = (2, 1, 1), \\ V_5 &= (1, 2, 1), V_6 = (0, 2, 1), V_7 = (0, 1, 1), \end{aligned} \tag{4.44}$$

and that we will denote as pseudo dP₄ or (P)dP₄. The dual gauge theory can be described by the following quiver:



with superpotential:

$$\begin{aligned} W &= \text{Tr} [X_{61} X_{17} X_{74} X_{46} + X_{21} X_{13} X_{35} X_{52} + X_{27} X_{73} X_{36} X_{62} + X_{14} X_{45} X_{51} + \\ &\quad - X_{51} X_{17} X_{73} X_{35} - X_{21} X_{14} X_{46} X_{62} - X_{27} X_{74} X_{45} X_{52} - X_{13} X_{36} X_{61}], \end{aligned} \tag{4.46}$$

where each X_{ij} must be understood as a field transforming in the bifundamental representation with respect to the i -th and j -th nodes. The charges of the fields can be parameterized in terms of the PMs using the assignation

X_{17}	X_{21}	X_{27}	X_{73}	X_{14}	X_{74}	X_{13}	X_{62}
$\mathbf{a}_1 + \mathbf{a}_6 + \mathbf{a}_7$	\mathbf{a}_7	$\mathbf{a}_3 + \mathbf{a}_4$	\mathbf{a}_2	$\mathbf{a}_1 + \mathbf{a}_2 + \mathbf{a}_3$	\mathbf{a}_5	$\mathbf{a}_4 + \mathbf{a}_5 + \mathbf{a}_6$	$\mathbf{a}_5 + \mathbf{a}_6$
X_{51}	X_{61}	X_{52}	X_{36}	X_{45}	X_{46}	X_{35}	
$\mathbf{a}_4 + \mathbf{a}_5$	$\mathbf{a}_2 + \mathbf{a}_3$	$\mathbf{a}_1 + \mathbf{a}_2$	$\mathbf{a}_1 + \mathbf{a}_7$	$\mathbf{a}_6 + \mathbf{a}_7$	\mathbf{a}_4	\mathbf{a}_3	

Thus, the set of charges suitable for the underlying geometry is:

⁶ Meaning that none of the possible triples of points lies on a line.

	$U(1)_1$	$U(1)_2$	$U(1)_3$	$U(1)_4$	$U(1)_5$	$U(1)_6$	$U(1)_R$
X_{17}	0	-1	-1	-1	-1	0	6/7
X_{21}	-1	-1	-1	-1	-1	-1	2/7
X_{27}	0	0	1	1	0	0	4/7
X_{73}	0	1	0	0	0	0	2/7
X_{14}	1	1	1	0	0	0	6/7
X_{74}	0	0	0	0	1	0	2/7
X_{13}	0	0	0	1	1	1	6/7
X_{62}	0	0	0	0	1	1	4/7
X_{51}	0	0	0	1	1	0	4/7
X_{61}	0	1	1	0	0	0	4/7
X_{52}	1	1	0	0	0	0	4/7
X_{36}	0	-1	-1	-1	-1	-1	4/7
X_{45}	-1	-1	-1	-1	-1	0	4/7
X_{46}	0	0	0	1	0	0	2/7
X_{46}	0	0	1	0	0	0	2/7

(4.48)

We assign a fugacity Δ_i to each global $U(1)_i$ and, assuming V_2 has a local minimum for vanishing holonomies, we computed the entropy function following the same guide-line as before. By considering the $|\omega|_1, |\omega|_2 \rightarrow 0$ and shifting each fugacity by $-\frac{\omega_1+\omega_2}{7}$ we obtain the following expression:

$$\begin{aligned}
 S_E = -\frac{i\pi N^2}{\omega_1\omega_2} & (\Delta_1\Delta_2\Delta_3 + 2\Delta_1\Delta_2\Delta_4 + 2\Delta_1\Delta_3\Delta_4 + \Delta_2\Delta_3\Delta_4 + 2\Delta_1\Delta_2\Delta_5 + \\
 & + 2\Delta_2\Delta_3\Delta_5 + 2\Delta_1\Delta_4\Delta_5 + 2\Delta_2\Delta_4\Delta_5 + \Delta_3\Delta_4\Delta_5 + \Delta_1\Delta_2\Delta_6 + \\
 & + 2\Delta_1\Delta_3\Delta_6 + 2\Delta_2\Delta_3\Delta_6 + 2\Delta_1\Delta_4\Delta_6 + 3\Delta_2\Delta_4\Delta_6 + 2\Delta_3\Delta_4\Delta_6 + \\
 & + \Delta_1\Delta_3\Delta_7 + 2\Delta_2\Delta_3\Delta_7 + 2\Delta_1\Delta_4\Delta_7 + 4\Delta_2\Delta_4\Delta_7 + 3\Delta_3\Delta_4\Delta_7 + \\
 & + 2\Delta_1\Delta_5\Delta_7 + 4\Delta_2\Delta_5\Delta_7 + 4\Delta_3\Delta_5\Delta_7 + 2\Delta_4\Delta_5\Delta_7 + \Delta_1\Delta_6\Delta_7 + \\
 & + 3\Delta_1\Delta_3\Delta_5 + \Delta_1\Delta_5\Delta_6 + 2\Delta_2\Delta_5\Delta_6 + 2\Delta_3\Delta_5\Delta_6 + \Delta_4\Delta_5\Delta_6 + \\
 & + 2\Delta_2\Delta_6\Delta_7 + 2\Delta_3\Delta_6\Delta_7 + \Delta_4\Delta_6\Delta_7) .
 \end{aligned}$$

(4.49)

This is again in agreement with (2.18).

5. Infinite families

In this section we compute the Cardy-like limit of the superconformal index at large N with complex fugacities, for infinite families of quiver gauge theories. We assume that the fugacities are in the regime $0 \leq \Delta_1, \dots, \Delta_{d-1} \leq \frac{1}{2}$ and $0 \leq \sum_{i=1}^{d-1} \Delta_i \leq 1$. In this regime we assume the validity of the conjecture on the existence of a universal saddle point associated to the vanishing of the holonomies. For each family we extract the entropy function and we verify the validity of the relation (2.18).

5.1. Y^{pq}

We start our analysis with the Y^{pq} quiver gauge theories, introduced in [33]. They correspond to quiver gauge theories with $2p$ gauge groups and a chiral field content of bifundamental fields. When p and q are generic the models enjoy an $SU(2) \times U(1)$ flavor symmetry and in addition one non-anomalous $U(1)_B$.

The toric diagram is parameterized by the four vectors

$$V_1 = (0, 0, 1), \quad V_2 = (1, 0, 1), \quad V_3 = (0, p, 1) \quad V_4 = (-1, p - q, 1) \tag{5.1}$$

As discussed in section 2 we can parameterize the global symmetries using the toric data. In this case there are four perfect matchings associated to the four external points of the toric diagram. The charges of the fields can be parameterized in terms of the PMs using the assignation

Y	U_1	Z	U_2	V_1	V_2	
\mathbf{a}_1	\mathbf{a}_2	\mathbf{a}_3	\mathbf{a}_4	$\mathbf{a}_2 + \mathbf{a}_3$	$\mathbf{a}_3 + \mathbf{a}_4$	

(5.2)

we can use this parameterization to construct the basis of $U(1)_i$ symmetries that we will use in the calculation of the index. Following the discussion in section 2 we have

	multiplicity	$U(1)_1$	$U(1)_2$	$U(1)_3$	$U(1)_R$	
Y	$p + q$	1	0	0	$\frac{1}{2}$	
U_1	p	0	1	0	$\frac{1}{2}$	
Z	$p - q$	0	0	1	$\frac{1}{2}$	(5.3)
U_2	p	-1	-1	-1	$\frac{1}{2}$	
V_1	q	-1	0	0	1	
V_2	q	-1	-1	0	1	

We can assign a fugacity Δ_i to the i -th $U(1)$ in this table. Furthermore we assign an equal R -symmetry to each perfect matching, such that the R -charges of the fields are given in the table. Then we shift each fugacity by $-\frac{\omega_1 + \omega_2}{4}$, where 4 refers to the number of points in the toric diagram. After this shift we compute the Cardy-like limit of the index at the universal saddle point, *i.e.* by setting all the gauge holonomies to zero. In this way, at large N , the leading contribution to the index, corresponding to the entropy function

$$S_E = -\frac{i\pi N^2}{\omega_1 \omega_2} \left(p\Delta_1 \Delta_2 \Delta_3 + ((p + q)\Delta_1 \Delta_2 + p\Delta_1 \Delta_3 + (p - q)\Delta_2 \Delta_3)(1 + \omega_1 + \omega_2 - \Delta_1 - \Delta_2 - \Delta_3) \right) \tag{5.4}$$

Defining $\Delta_4 \equiv 1 + \omega_1 + \omega_2 - \Delta_1 - \Delta_2 - \Delta_3$ the entropy function in (5.4) becomes

$$S_E = -\frac{i\pi N^2}{\omega_1 \omega_2} \left(p\Delta_1 \Delta_2 \Delta_3 + (p + q)\Delta_1 \Delta_2 \Delta_4 + p\Delta_1 \Delta_3 \Delta_4 + (p - q)\Delta_2 \Delta_3 \Delta_4 \right) \tag{5.5}$$

It is straightforward to check that the final form of the entropy function is then given by (2.18), where the coefficients C_{IJK} , are computed from (2.1).

5.2. X^{pq}

These models have been introduced in [34]. For generic values of p and q there are $2p + 1$ gauge groups, there is a $U(1)^2$ flavor symmetry and two non anomalous baryonic $U(1)$ symmetries. The toric diagram is parameterized by the five vectors

$$V_1 = (1 - q, 1, 1), V_2 = (-1, 0, 1), V_3 = (q - p, -1, 1), V_4 = (0, -1, 1), V_5 = (p, 1, 1) \tag{5.6}$$

As discussed in section 2 we can parameterize the global charges as

multiplicity	$U(1)_1$	$U(1)_2$	$U(1)_3$	$U(1)_4$	$U(1)_R$
$p + q - 1$	1	0	0	0	$\frac{2}{5}$
1	0	1	0	0	$\frac{2}{5}$
1	0	0	1	0	$\frac{2}{5}$
$p - q$	0	0	0	1	$\frac{2}{5}$
p	-1	-1	-1	-1	$\frac{2}{5}$
$p - 1$	0	1	1	0	$\frac{4}{5}$
1	0	0	1	1	$\frac{4}{5}$
$q - 1$	0	1	1	1	$\frac{6}{5}$
1	1	1	0	0	$\frac{4}{5}$
q	-1	-1	-1	0	$\frac{4}{5}$

(5.7)

Then we shift each charge by $-\frac{\omega_1 + \omega_2}{5}$, where 5 refers to the number of points in the toric diagram. We also assign a trial R-symmetry to each field as in the table above. Then we compute the Cardy-like limit of the index by setting all the holonomies to zero and supposing that there exists a regime of charges such that a minimum exists. In this way, at large N , the entropy function is

$$S_E = -\frac{i\pi N^2}{\omega_1 \omega_2} \left(\Delta_1 \Delta_5 \Delta_3 (p + q) + \Delta_4 \Delta_5 \Delta_3 (p - q) + \Delta_1 \Delta_2 \Delta_5 (p + q - 1) \right. \\ \left. + \Delta_2 \Delta_4 \Delta_5 (p - q + 1) + \Delta_1 \Delta_4 \Delta_5 p + \Delta_1 \Delta_4 \Delta_3 p + \Delta_1 \Delta_2 \Delta_4 p \right. \\ \left. + \Delta_1 \Delta_2 \Delta_3 + \Delta_2 \Delta_4 \Delta_3 + 2\Delta_2 \Delta_5 \Delta_3 \right) \tag{5.8}$$

where we defined $\Delta_5 = 1 + \omega_1 + \omega_2 - \Delta_1 - \Delta_2 - \Delta_3 - \Delta_4$.

This formula can be interpreted in terms of the geometric data by assigning the charges Δ_I the four corners of the toric diagram. The final form of the entropy function is then given by (2.18), where the coefficients C_{IJK} are computed from (2.1).

5.3. L^{pqr}

These models have been introduced in [35–37]. The toric diagram is parameterized by the four vectors

$$V_1 = (0, 0, 1), \quad V_2 = (1, 0, 1), \quad V_3 = (P, s, 1) \quad V_4 = (-k, q, 1) \tag{5.9}$$

where $Pq = r - ks$. If $p \neq r$ we can parameterize the global charges as ⁷

	multiplicity	$U(1)_1$	$U(1)_2$	$U(1)_3$	$U(1)_R$
Y	q	1	0	0	$\frac{1}{2}$
W_2	s	0	1	0	$\frac{1}{2}$
Z	p	0	0	1	$\frac{1}{2}$
X_2	r	-1	-1	-1	$\frac{1}{2}$
W_1	$q - s$	0	1	1	1
X_1	$q - r$	-1	-1	0	1

(5.10)

where $p + q = r + s$.

Then we shift each charge by $-\frac{\omega_1 + \omega_2}{4}$, where 4 refers to the number of points in the toric diagram. We also assign a trial R -symmetry to each field as in the table above. Then we compute the Cardy-like limit of the index by setting all the holonomies to zero and supposing that there exists a regime of charges such that a minimum exists. In this way, at large N , we can show that the entropy function is equivalent to

$$S_E = -\frac{i\pi N^2}{\omega_1 \omega_2} \left(s \Delta_1 \Delta_2 \Delta_3 + q \Delta_1 \Delta_2 \Delta_4 + r \Delta_1 \Delta_3 \Delta_4 + p \Delta_2 \Delta_4 \Delta_3 \right) \tag{5.11}$$

where $\Delta_4 = 1 + \omega_1 + \omega_2 - \Delta_1 - \Delta_2 - \Delta_3$.

This formula can be interpreted in terms of the geometric data by assigning the charges Δ_I the four corners of the toric diagram. The final form of the entropy function is then given by (2.18), where the coefficients C_{IJK} are computed from (2.1).

6. Legendre transform and the entropy

In this section we obtain the entropy associated to some of the families discussed above. We focus on the Y^{pp} and on the L^{0b0} cases. These two cases are similar to the $\mathcal{N} = 4$ case because they can be constructed by an orbifold projection of \mathbb{C}^3 . At the level of the toric diagram this reflects in the fact that there are three corners. The other external points are on the perimeter, signaling the presence of non smooth horizons induced by the orbifold. These models are anyway richer, because they have a higher amount of global, baryonic, symmetries. In this section we compute the Legendre transform of the entropy function for these models, by turning on all of the possible global symmetries. The ones discussed in this section are the only cases where we have found an expression for the entropy by computing the Legendre transform. For other geometries with more than three corners in the toric diagram and all the global symmetries turned on, we have not found a systematic way to compute the Legendre transform of the entropy function.

Let us also comment on the C_{IJK} coefficients for the theories that we discuss below with respect to the multi-charge AdS_5 black holes obtained from gauged supergravity in [6]. On the supergravity side the condition

⁷ In the case $p = r$ the toric diagram gains a large amount of external points lying on the perimeter. It induces a large set of non-anomalous baryonic symmetries in the quiver.

$$C_{IJK}C_{J'(LM}C_{PQ)K'}\delta^{JJ'}\delta^{KK'} = \frac{4}{3}\delta_{I(L}C_{MPQ)} \tag{6.1}$$

was imposed, while here we have explicitly checked that the C_{IJK} coefficients discussed in this section do not satisfy (6.1). This is a common feature of all the SE_5 theories except the case of S^5 . On the other hand, in these cases there are not truncations available in the literature with all the global symmetries of the dual field theory appearing as massless vector multiplets in the gravitational description. This implies that we cannot compare the results discussed in this section with the ones expected from gauged supergravity. The only partial exception is the case of the conifold, where a truncation with one vector multiplet was found in [38]. This vector multiplet, denoted as Betti multiplet, corresponds to the baryonic symmetry of the dual field theory. In this case there is also an hypermultiplet, signaling the presence of a massive vector multiplet as well. The entropy for this model has been recently discussed in [39], and the result is in agreement with our result.

6.1. The Y^{PP} family

In this section we study the Legendre transform of the entropy function obtained in the case of Y^{PP} models. The entropy function is given by

$$S_E = -\frac{i\pi p N^2}{\omega_1\omega_2}(\Delta_1\Delta_2\Delta_3 + \Delta_1\Delta_3\Delta_4 + 2\Delta_1\Delta_2\Delta_4) \tag{6.2}$$

with $\sum_I \Delta_I = \omega_1 + \omega_2 - 1$. The Legendre transform is computed in terms of the conjugate charges Q_I and angular momenta J_a and it corresponds to

$$S(Q, J) = S_E(X, \omega) + 2\pi i \left(\sum_{I=1}^4 \Delta_I Q_I + \sum_{a=1}^2 \omega_a J_a \right) + 2\pi i \Lambda \left(\sum_{I=1}^4 \Delta_I - \sum_{a=1}^2 \omega_a - 1 \right) \tag{6.3}$$

Observing that

$$S_E(X, \omega) = \left(\sum_{I=1}^4 \Delta_I \frac{\partial S_E}{\partial \Delta_I} + \sum_{a=1}^2 \omega_a \frac{\partial S_E}{\partial \omega_a} \right) \tag{6.4}$$

we have that $S(Q, J) = 2\pi i \Lambda$. The Lagrange multiplier can be obtained from the equation

$$(\Lambda + Q_1)(2(2(\Lambda + Q_3) + (\Lambda + Q_4))(\Lambda + Q_2) - (\Lambda + Q_2)^2 - (2(\Lambda + Q_3) - (\Lambda + Q_4))^2) + 4N^2 p (\Lambda - J_1)(\Lambda - J_2) = 0 \tag{6.5}$$

Reorganizing the polynomial on the LHS of this equation in the form $\Lambda^3 + \Lambda^2 p_2 + \Lambda p_1 + p_0$ we have two imaginary solutions if $p_0 = p_1 p_2$. The coefficients p_i in this case are

$$\begin{aligned} p_2 &= N^2 p + Q_1 + Q_2 + Q_4 \\ p_1 &= (Q_1 + Q_3)(Q_2 + Q_4) + \frac{Q_4 Q_2}{2} - \frac{Q_2^2}{4} - Q_3^2 - \frac{Q_4^2}{4} - N^2 p (J_1 + J_2) \\ p_0 &= N^2 p J_1 J_2 - \frac{1}{4} Q_1 Q_2^2 + Q_1 Q_3 Q_2 + \frac{1}{2} Q_1 Q_4 Q_2 - Q_1 Q_3^2 - \frac{1}{4} Q_1 Q_4^2 + Q_1 Q_3 Q_4 \end{aligned} \tag{6.6}$$

and the entropy corresponds to

$$S(Q, J) = 2\pi \sqrt{(Q_1 + Q_3)(Q_2 + Q_4) + \frac{Q_4 Q_2}{2} - \frac{Q_2^2}{4} - Q_3^2 - \frac{Q_4^2}{4} - N^2 p(J_1 + J_2)} \tag{6.7}$$

Furthermore this model has been recently analyzed by [20]. The authors studied the entropy by turning off the fugacity for the $SU(2)_L$ symmetry. It corresponds here to turn off the variable Δ_2 in (6.2). In this case the entropy function becomes

$$S_E(X, \omega) = -\frac{i\pi p N^2}{\omega_1 \omega_2} (\Delta_1 \Delta_3 \Delta_4) \tag{6.8}$$

with the constraint $\Delta_1 + \Delta_3 + \Delta_4 = \omega_1 + \omega_2 - 1$. One can repeat the analysis discussed above. The relevant point of the discussion is that in this case the cubic equation for the Lagrange multiplier is

$$(\Lambda + Q_1)(\Lambda + Q_3)(\Lambda + Q_4) + \frac{N^2 p}{2} (\Lambda - J_1)(\Lambda - J_2) = 0 \tag{6.9}$$

and the entropy becomes

$$S(Q, J) = 2\pi \sqrt{Q_1 Q_3 + Q_1 Q_4 + Q_3 Q_4 - \frac{N^2 p}{2} (J_1 + J_2)} \tag{6.10}$$

This result can be mapped to the one of [20] by mapping the charges Q_I to the ones discussed there.

6.2. The L^{aba} family

Here we compute the Legendre transform of the entropy function of a family of necklace quivers, that correspond to the L^{0b0} family studied above. This class of models has already been discussed by [18], where it has been shown that the orbifold modifies just an overall contribution to the entropy function. This was proven by just studying the effect of the flavor symmetries, but in this case there are in addition $G - 1$ non-anomalous baryonic symmetries, being G the number of gauge groups in the necklace. Here we study the entropy function for a generic parameterization of the charges, taking care of all the non anomalous baryonic symmetries as well.

As discussed in sub-section 4.1 the entropy function can be expressed in terms of the areas of the toric diagram, contracted with the fugacities Δ_I , with $I = 1, \dots, d$ runs over all the external points of the toric diagram. In this case the formula can be expressed as

$$S_E = -i\pi N^2 \frac{\Delta_1 \sum_{i,j=2}^d |i - j| \Delta_i \Delta_j}{2\omega_1 \omega_2} \tag{6.11}$$

The entropy is given by the Legendre transform

$$S(Q, J) = S_E(\Delta, \omega) + 2\pi i \left(\sum_{I=1}^d \Delta_I Q_I + \sum_{a=1}^2 \omega_a J_a \right) + 2\pi i \Lambda \left(\sum_{I=1}^d \Delta_I - \sum_{a=1}^2 \omega_a - 1 \right) \tag{6.12}$$

The relation

$$S_E = \sum_{I=1}^d \Delta_I \frac{\partial S_E}{\partial \Delta_I} + \sum_{a=1}^2 \omega_a \frac{\partial S_E}{\partial J_a} \tag{6.13}$$

guarantees that

$$S_Q = 2\pi i \Lambda \tag{6.14}$$

By using the equations of motion we have induced a cubic relation satisfied by the Lagrange multiplier Λ . The cubic relation for Λ is

$$\begin{aligned} & (Q_1 + \Lambda) \left(\sum_{i=2}^{d-1} (Q_i + \Lambda)(Q_{i+1} + \Lambda) - \sum_{i=2}^{d-1} (Q_i + \Lambda)^2 + \frac{(Q_2 + \Lambda)(Q_d + \Lambda)}{d-2} \right. \\ & \left. + \frac{(d-1)((Q_2 + \Lambda)^2 + (Q_d + \Lambda)^2)}{2(d-2)} \right) + N^2(\Lambda - J_1)(\Lambda - J_2) = 0 \end{aligned} \tag{6.15}$$

This equation is of the form $\Lambda^3 + p_2\Lambda^2 + p_1\Lambda + p_0 = 0$, with

$$\begin{aligned} p_2 &= (d-2)\frac{N^2}{2} + Q_1 + Q_2 + Q_d, \\ p_1 &= 2Q_1(Q_2 + Q_d) + Q_2Q_d - \frac{d-1}{2}(Q_2^2 + Q_d^2) \\ &+ (d-2) \left(\sum_{i=2}^{d-1} Q_i Q_{i+1} - \sum_{i=2}^d Q_i^2 \right) - (d-2)\frac{N^2}{2}(J_1 + J_2) \\ p_0 &= (d-2)\frac{N^2}{2}J_1J_2 + \frac{1}{4}Q_1 \left((d-1)(Q_2^2 + Q_d^2) + 2Q_2Q_d \right. \\ &\left. - 2(d-2)\sum_{i=2}^d Q_i^2 + 2(d-2)\sum_{i=2}^{d-1} Q_i Q_{i+1} \right). \end{aligned} \tag{6.16}$$

There is an imaginary solution for $p_2p_1 = p_0$, and in this case the entropy is given by $\Lambda = -i\sqrt{p_1}$, or more explicitly

$$\begin{aligned} S(Q, J) &= 2\pi \left(2Q_1(Q_2 + Q_d) + Q_2Q_d - \frac{d-1}{2}(Q_2^2 + Q_d^2) \right. \\ &\left. + (d-2) \left(\sum_{i=2}^{d-1} Q_i Q_{i+1} - \sum_{i=2}^d Q_i^2 \right) - (d-2)\frac{N^2}{2}(J_1 + J_2) \right)^{\frac{1}{2}} \end{aligned} \tag{6.17}$$

Observe that such expression correctly gives us back the entropy for $\mathcal{N} = 4$ SYM case for $d = 3$.

7. Conclusions

In this paper we studied the Cardy-like behavior of the SCI in presence of complex fugacities. This quantity has been recently observed to reproduce, in the case of $\mathcal{N} = 4$ SYM, the entropy of an AdS₅ rotating black hole, through a Legendre transform. Here we focused on infinite families of 4d $\mathcal{N} = 1$ quiver gauge theories, describing stacks of D3 branes probing the tip of a toric CY₃ cones over a 5d SE₅ base. We showed that the general formula (2.18) for the entropy function of the models under investigation can be obtained from the Cardy-like limit of the SCI with complex fugacities. Furthermore we computed the Legendre transform for some of the models analyzed here, giving a prediction for the entropy of the dual black hole.

In the analysis we left many open questions that deserve further investigation. First we conjectured that it is always possible to find a regime of charges such that the holonomies are vanishing at the saddle point. This conjecture is consistent with the ones given in [9,18,19]. Further arguments in favor of this idea has been recently given by [14]. In addition we have obtained the expected result in a regime of fugacities corresponding to the choices $0 \leq \Delta_1, \dots, \Delta_{d-1} \leq \frac{1}{2}$ and $0 \leq \sum_{i=1}^{d-1} \Delta_i \leq 1$. In other regimes we have not found a minimum of the potential $V_2(a)$ but a plateau. Similar plateaux have been discussed in [18], but in that case they appeared for the choice $\text{Re}\left(\frac{i}{\omega_1\omega_2}\right) < 0$ and they were associated to the Stokes lines discussed also in [16]. Here the plateaux appear also in the regime $\text{Re}\left(\frac{i}{\omega_1\omega_2}\right) > 0$ and it should be interesting to have a deeper understanding of them and of their holographic dual interpretation. Furthermore, we did not find a general way to obtain the Legendre transform for entropy functions associated to toric diagram with more than three external corners if all the global symmetries are turned on. It should be interesting to see if this is just a technical obstruction or if is there a deeper physical reason. We conclude observing that a similar geometric relation between the Cardy-like limit of the SCI and the entropy function can be expected for non-toric cases, as the one discussed in [40]. It should be interesting to investigate along this line.

CRedit authorship contribution statement

Antonio Amariti: Conceptualization, Formal analysis, Methodology, Software, Validation, Writing – original draft, Writing – review & editing. **Ivan Garozzo:** Formal analysis, Methodology, Software, Validation, Writing – original draft, Writing – review & editing. **Gabriele Lo Monaco:** Formal analysis, Methodology, Software, Validation, Writing – original draft, Writing – review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We are grateful to Alberto Zaffaroni, Alessandra Gnechchi, Francesco Benini and Noppadol Mekareeya for useful comments. This work has been supported in part by Italian Ministero dell’Istruzione, Università e Ricerca (MIUR) and Istituto Nazionale di Fisica Nucleare (INFN) through the “Gauge Theories, Strings, Supergravity” (GSS) research project. We gratefully acknowledge the ICTP where some of the research for this paper was performed during the 2019 “Spring School on Superstring Theory and Related Topics”. A.A. thanks CERN for the hospitality during the completion of this project.

Appendix A. Saddle point analysis for the conifold at higher-rank

In section 3 we studied the behavior of the minima of V_2 with respect to the fugacity range in the case of $SU(2)$ gauge groups. In this appendix we want to collect some evidence about the possibility of extending those considerations to higher ranks. Let us briefly remind the main results: V_2 can be expressed as




Fig. 3. V_2 for $SU(3)$ conifold with cyclically identified holonomies: on the left we fixed $\Delta_{F_1} = 0.12, \Delta_{F_2} = 0.15, \Delta_B = 0.22$ and we can observe that the function enjoys a local maximum at the origin; on the right we fixed $\Delta_{F_1} = 0.05, \Delta_{F_2} = 0.1, \Delta_B = 0.3$ and we can observe that V_2 only possesses plateaux.

$$V_2 = - \sum_{m,n=1}^N \left(\kappa \left[a_m^{(1)} - a_n^{(2)} + \Delta_{F_1} + \Delta_{F_2} + \Delta_B \right] + \kappa \left[a_m^{(1)} - a_n^{(2)} - \Delta_{F_1} - \Delta_{F_2} + \Delta_B \right] + \kappa \left[a_m^{(2)} - a_n^{(1)} + \Delta_{F_1} - \Delta_{F_2} - \Delta_B \right] + \kappa \left[a_m^{(2)} - a_n^{(1)} - \Delta_{F_1} + \Delta_{F_2} - \Delta_B \right] \right), \tag{A.1}$$

where $a_m^{(k)}$ are holonomies for k -th gauge group and $\Delta_{F_1}, \Delta_{F_2}, \Delta_B$ are fugacities for flavor and baryonic symmetries. At rank $N - 1$ we have to enforce the constraint:

$$a_N^{(k)} = - \sum_{m=1}^{N-1} a_m^{(k)}, \quad k = 1, 2, \tag{A.2}$$

so that in rank-1 case we are left with just two independent variables. We fixed a chamber where

$$0 \leq \Delta_{F_1} \leq \Delta_{F_2} \leq \Delta_B \leq 1/2, \quad 0 \leq \Delta_{F_1} + \Delta_{F_2} + \Delta_B \leq 1/2, \tag{A.3}$$

finding two possible behaviors for V_2 :

- $\Delta_B > \Delta_{F_1} + \Delta_{F_2}$: V_2 admits only plateaux of minima and thus the index is hard to evaluate.
- $\Delta_B < \Delta_{F_1} + \Delta_{F_2}$: V_2 admits a local maximum for vanishing holonomies that dominates in the Cardy-like limit.

Performing a similar analysis for higher rank is more complicated, because more variables are involved; however we can use the high symmetry of the model to simplify the computation: a natural expectation is that at high temperature, *i.e.* in the Cardy limit, all the global symmetries are preserved and no gauge symmetries are broken, so that no Higgs mechanisms are involved. In other words, we want to count the degrees of freedom of the theory in the deconfining phase. A symmetry that we expect to be preserved at high temperature is a \mathbb{Z}_2 discrete symmetry of the quiver, exchanging the two nodes and the two couples of bifundamental fields; in order to keep this symmetry, we impose the following cyclic condition:

$$a_m^{(1)} = a_m^{(2)} \quad \forall m, \tag{A.4}$$

as already suggested in [18].




Fig. 4. V_2 in the $a_1^{(1)} - a_2^{(1)}$ plane for $SU(4)$ conifold with cyclically identified holonomies and $\Delta_{F_1} = 0.1, \Delta_{F_2} = 0.15, \Delta_B = 0.2$; from the top left in clockwise sense we fixed $a_3^{(1)} = 0, a_3^{(1)} = 0.08$ and $a_3^{(1)} = 0.16$. We notice a minimum whose lowest value is reached for $a_3^{(1)} = 0$.




Fig. 5. V_2 in the $a_1^{(1)} - a_2^{(1)}$ plane for $SU(4)$ conifold with cyclically identified holonomies and $\Delta_{F_1} = 0.05, \Delta_{F_2} = 0.1, \Delta_B = 0.3, a_3^{(1)} = 0$; only plateaux are present.

For the rank-2 case this is enough in order to study numerically V_2 , that is again a function of two variables only, $a_1^{(1)}$ and $a_2^{(1)}$. The plots of the function in Fig. 3 shows that V_2 still shares the same properties as before.

We can perform a similar analysis at rank 3. In this case we have three independent variables, $a_1^{(1)}, a_2^{(1)}$ and $a_3^{(1)}$ and thus we cannot make a single plot; we need to use a slightly different technique: we can make a plot V_2 in the $a_1^{(1)} - a_2^{(1)}$ plane at fixed $a_3^{(1)}$ and then vary the value of this last holonomy. If a minimum is located at the origin, V_2 restricted to the $a_1^{(1)} - a_2^{(1)}$ plane

should have a minimum, as deep as we get closer to $a_3^{(1)} = 0$. This is the kind of behavior that we can observe in Fig. 4, where we fixed the fugacity in such a way the condition $\Delta_{F_1} + \Delta_{F_2} > \Delta_B$ to hold. When $\Delta_{F_1} + \Delta_{F_2} < \Delta_B$, instead, the presence of plateaux is already evident from a plot of V_2 at $a_3^{(1)} = 0$, as shown in Fig. 5. Let us stress finally stress that, even relaxing the assumption (A.4), we can use other numerical tools such as **FindMaximum/FindMinimum** of Mathematica in order to understand the behavior of V_2 ; this kind of study still returns the same results as before.

References

- [1] A. Strominger, C. Vafa, Microscopic origin of the Bekenstein-Hawking entropy, *Phys. Lett. B* 379 (1996) 99–104, arXiv:hep-th/9601029 [hep-th].
- [2] J.B. Gutowski, H.S. Reall, General supersymmetric AdS(5) black holes, *J. High Energy Phys.* 04 (2004) 048, arXiv:hep-th/0401129 [hep-th].
- [3] J.B. Gutowski, H.S. Reall, Supersymmetric AdS(5) black holes, *J. High Energy Phys.* 02 (2004) 006, arXiv:hep-th/0401042 [hep-th].
- [4] Z.W. Chong, M. Cvetič, H. Lu, C.N. Pope, Five-dimensional gauged supergravity black holes with independent rotation parameters, *Phys. Rev. D* 72 (2005) 041901, arXiv:hep-th/0505112 [hep-th].
- [5] Z.W. Chong, M. Cvetič, H. Lu, C.N. Pope, General non-extremal rotating black holes in minimal five-dimensional gauged supergravity, *Phys. Rev. Lett.* 95 (2005) 161301, arXiv:hep-th/0506029 [hep-th].
- [6] H.K. Kunduri, J. Lucietti, H.S. Reall, Supersymmetric multi-charge AdS(5) black holes, *J. High Energy Phys.* 04 (2006) 036, arXiv:hep-th/0601156 [hep-th].
- [7] J. Kinney, J.M. Maldacena, S. Minwalla, S. Raju, An index for 4 dimensional super conformal theories, *Commun. Math. Phys.* 275 (2007) 209–254, arXiv:hep-th/0510251 [hep-th].
- [8] C. Romelsberger, Counting chiral primaries in $N = 1, d=4$ superconformal field theories, *Nucl. Phys. B* 747 (2006) 329–353, arXiv:hep-th/0510060 [hep-th].
- [9] S. Choi, J. Kim, S. Kim, J. Nahmgoong, Large AdS black holes from QFT, arXiv:1810.12067 [hep-th].
- [10] S. Choi, J. Kim, S. Kim, J. Nahmgoong, Comments on deconfinement in AdS/CFT, arXiv:1811.08646 [hep-th].
- [11] S.M. Hosseini, K. Hristov, A. Zaffaroni, An extremization principle for the entropy of rotating BPS black holes in AdS₅, *J. High Energy Phys.* 07 (2017) 106, arXiv:1705.05383 [hep-th].
- [12] A. Sen, Quantum entropy function from AdS(2)/CFT(1) correspondence, *Int. J. Mod. Phys. A* 24 (2009) 4225–4244, arXiv:0809.3304 [hep-th].
- [13] A. Sen, Arithmetic of quantum entropy function, *J. High Energy Phys.* 08 (2009) 068, arXiv:0903.1477 [hep-th].
- [14] A. Cabo-Bizet, D. Cassani, D. Martelli, S. Murthy, The asymptotic growth of states of the 4d $N=1$ superconformal index, arXiv:1904.05865 [hep-th].
- [15] A. Cabo-Bizet, D. Cassani, D. Martelli, S. Murthy, Microscopic origin of the Bekenstein-Hawking entropy of supersymmetric AdS₅ black holes, arXiv:1810.11442 [hep-th].
- [16] F. Benini, P. Milan, Black holes in 4d $\mathcal{N} = 4$ Super-Yang-Mills, arXiv:1812.09613 [hep-th].
- [17] F. Benini, P. Milan, A Bethe Ansatz type formula for the superconformal index, arXiv:1811.04107 [hep-th].
- [18] M. Honda, Quantum black hole entropy from 4d supersymmetric Cardy formula, arXiv:1901.08091 [hep-th].
- [19] A. Arabi Ardehali, Cardy-like asymptotics of the 4d $\mathcal{N} = 4$ index and AdS₅ blackholes, arXiv:1902.06619 [hep-th].
- [20] J. Kim, S. Kim, J. Song, A 4d $N=1$ Cardy formula, arXiv:1904.03455 [hep-th].
- [21] S.M. Hosseini, K. Hristov, A. Zaffaroni, A note on the entropy of rotating BPS AdS₇ \times S⁴ black holes, *J. High Energy Phys.* 05 (2018) 121, arXiv:1803.07568 [hep-th].
- [22] A. Zaffaroni, Lectures on AdS black holes, holography and localization, arXiv:1902.07176 [hep-th], 2019.
- [23] S. Benvenuti, L.A. Pando Zayas, Y. Tachikawa, Triangle anomalies from Einstein manifolds, *Adv. Theor. Math. Phys.* 10 (3) (2006) 395–432, arXiv:hep-th/0601054 [hep-th].
- [24] A. Hanany, K.D. Kennaway, Dimer models and toric diagrams, arXiv:hep-th/0503149 [hep-th].
- [25] S. Franco, A. Hanany, K.D. Kennaway, D. Vegh, B. Wecht, Brane dimers and quiver gauge theories, *J. High Energy Phys.* 01 (2006) 096, arXiv:hep-th/0504110 [hep-th].
- [26] A. Butti, A. Zaffaroni, R-charges from toric diagrams and the equivalence of a-maximization and Z-minimization, *J. High Energy Phys.* 11 (2005) 019, arXiv:hep-th/0506232 [hep-th].
- [27] K.A. Intriligator, B. Wecht, The exact superconformal R symmetry maximizes a, *Nucl. Phys. B* 667 (2003) 183–200, arXiv:hep-th/0304128 [hep-th].

- [28] D. Martelli, J. Sparks, S.-T. Yau, The geometric dual of a-maximisation for toric Sasaki-Einstein manifolds, *Commun. Math. Phys.* 268 (2006) 39–65, arXiv:hep-th/0503183 [hep-th].
- [29] L. Di Pietro, Z. Komargodski, Cardy formulae for SUSY theories in $d = 4$ and $d = 6$, *J. High Energy Phys.* 12 (2014) 031, arXiv:1407.6061 [hep-th].
- [30] L. Di Pietro, M. Honda, Cardy formula for 4d SUSY theories and localization, *J. High Energy Phys.* 04 (2017) 055, arXiv:1611.00380 [hep-th].
- [31] A. Arabi Ardehali, High-temperature asymptotics of supersymmetric partition functions, *J. High Energy Phys.* 07 (2016) 025, arXiv:1512.03376 [hep-th].
- [32] I.R. Klebanov, E. Witten, Superconformal field theory on three-branes at a Calabi-Yau singularity, *Nucl. Phys. B* 536 (1998) 199–218, arXiv:hep-th/9807080 [hep-th].
- [33] S. Benvenuti, S. Franco, A. Hanany, D. Martelli, J. Sparks, An Infinite family of superconformal quiver gauge theories with Sasaki-Einstein duals, *J. High Energy Phys.* 06 (2005) 064, arXiv:hep-th/0411264 [hep-th].
- [34] A. Hanany, P. Kazakopoulos, B. Wecht, A new infinite class of quiver gauge theories, *J. High Energy Phys.* 08 (2005) 054, arXiv:hep-th/0503177 [hep-th].
- [35] S. Benvenuti, M. Kruczenski, From Sasaki-Einstein spaces to quivers via BPS geodesics: $L^{p,q|r}$, *J. High Energy Phys.* 04 (2006) 033, arXiv:hep-th/0505206 [hep-th].
- [36] A. Butti, D. Forcella, A. Zaffaroni, The dual superconformal theory for $L^{p,q,r}$ manifolds, *J. High Energy Phys.* 09 (2005) 018, arXiv:hep-th/0505220 [hep-th].
- [37] S. Franco, A. Hanany, D. Martelli, J. Sparks, D. Vegh, B. Wecht, Gauge theories from toric geometry and brane tilings, *J. High Energy Phys.* 01 (2006) 128, arXiv:hep-th/0505211 [hep-th].
- [38] D. Cassani, A.F. Faedo, A supersymmetric consistent truncation for conifold solutions, *Nucl. Phys. B* 843 (2011) 455–484, arXiv:1008.0883 [hep-th].
- [39] F. Benini, E. Colombo, S. Soltani, A. Zaffaroni, Z. Zhang, Superconformal indices at large N and the entropy of $\text{AdS}_5 \times \text{SE}_5$ black holes, *Class. Quantum Gravity* 37 (21) (2020) 215021, arXiv:2005.12308 [hep-th].
- [40] A. Butti, A. Zaffaroni, D. Forcella, Deformations of conformal theories and non-toric quiver gauge theories, *J. High Energy Phys.* 02 (2007) 081, arXiv:hep-th/0607147 [hep-th].