

# A global research priority agenda to advance public health responses to fatty liver disease

Jeffrey V. Lazarus<sup>1,2,3,\*†</sup>, Henry E. Mark<sup>4,5,†</sup>, Alina M. Allen<sup>6,†</sup>, Juan Pablo Arab<sup>7,8,9,†</sup>, Patrizia Carrieri<sup>10,†</sup>, Mazen Nouredin<sup>11,†</sup>, William Alazawi<sup>12</sup>, Naim Alkhouri<sup>13</sup>, Saleh A. Alqahtani<sup>14</sup>, Marco Arrese<sup>9</sup>, Ramon Bataller<sup>15</sup>, Thomas Berg<sup>16</sup>, Paul N. Brennan<sup>17</sup>, Patrizia Burra<sup>18</sup>, Graciela E. Castro-Narro<sup>19,20,21</sup>, Helena Cortez-Pinto<sup>22</sup>, Kenneth Cusi<sup>23</sup>, Nikos Dedes<sup>24</sup>, Ajay Duseja<sup>25</sup>, Sven M. Francque<sup>26,27</sup>, Hannes Hagström<sup>28</sup>, Terry T.-K. Huang<sup>3,29</sup>, Dana Ivancovsky Wajcman<sup>1</sup>, Achim Kautz<sup>30</sup>, Christopher J. Kopka<sup>31</sup>, Aleksander Krag<sup>32</sup>, Veronica Miller<sup>33</sup>, Philip N. Newsome<sup>34</sup>, Mary E. Rinella<sup>35</sup>, Diana Romero<sup>36</sup>, Shiv Kumar Sarin<sup>37</sup>, Marcelo Silva<sup>38</sup>, C. Wendy Spearman<sup>39</sup>, Emmanuel A. Tsochatzis<sup>40</sup>, Luca Valenti<sup>41,42</sup>, Marcela Villota-Rivas<sup>1</sup>, Shira Zelber-Sagi<sup>43,44</sup>, Jörn M. Schattenberg<sup>45,†</sup>, Vincent Wai-Sun Wong<sup>46,†</sup>, Zobair M. Younossi<sup>47,†</sup>, on behalf of the Healthy Livers, Healthy Lives Collaborators

Journal of Hepatology 2023. vol. 79 | 618–634



**Background & aims:** An estimated 38% of adults worldwide have non-alcoholic fatty liver disease (NAFLD). From individual impacts to widespread public health and economic consequences, the implications of this disease are profound. This study aimed to develop an aligned, prioritised fatty liver disease research agenda for the global health community.

**Methods:** Nine co-chairs drafted initial research priorities, subsequently reviewed by 40 core authors and debated during a three-day in-person meeting. Following a Delphi methodology, over two rounds, a large panel (R1 n = 344, R2 n = 288) reviewed the priorities, via Qualtrics XM, indicating agreement using a four-point Likert-scale and providing written feedback. The core group revised the draft priorities between rounds. In R2, panellists also ranked the priorities within six domains: epidemiology, models of care, treatment and care, education and awareness, patient and community perspectives, and leadership and public health policy.

**Results:** The consensus-built fatty liver disease research agenda encompasses 28 priorities. The mean percentage of 'agree' responses increased from 78.3 in R1 to 81.1 in R2. Five priorities received unanimous combined agreement ('agree' + 'somewhat agree'); the remaining 23 priorities had >90% combined agreement. While all but one of the priorities exhibited at least a supermajority of agreement (>66.7% 'agree'), 13 priorities had <80% 'agree', with greater reliance on 'somewhat agree' to achieve >90% combined agreement.

**Conclusions:** Adopting this multidisciplinary consensus-built research priorities agenda can deliver a step-change in addressing fatty liver disease, mitigating against its individual and societal harms and proactively altering its natural history through prevention, identification, treatment, and care. This agenda should catalyse the global health community's efforts to advance and accelerate responses to this widespread and fast-growing public health threat.

© 2023 The Author(s). Published by Elsevier B.V. on behalf of European Association for the Study of the Liver. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Over the past three decades, the fatty liver disease burden has increased drastically. An estimated 38% (95% CI 33.71–42.49) of the global adult population<sup>1,2</sup> and around 13% of children and adolescents,<sup>3</sup> now have the disease. Left unmanaged, the disease can progress through increasing stages of hepatic fibrosis, leading to cirrhosis and associated complications, including hepatocellular carcinoma (HCC) (Box 1).<sup>4,5</sup> Fatty liver disease causes quality of life impairments, which worsen with disease progression and are compounded by comorbidities.<sup>6–8</sup> Fatty liver disease is a leading cause of HCC, which is the second leading cause of years of life lost amongst all cancers.<sup>9</sup> Beyond the human toll, the disease has wide-reaching social

and economic implications and yet it remains under-recognised and under-evaluated.<sup>10,11</sup>

Through cardiometabolic risk factors, fatty liver disease shares a complex bi-directional relationship with other common diseases, including cardiovascular disease, the leading cause of death in those with fatty liver disease.<sup>13</sup> Type 2 diabetes mellitus, cancer, sarcopenia, and chronic kidney disease are all commonly associated with fatty liver disease,<sup>14,15</sup> with the risk of extrahepatic complications increasing in parallel with liver disease severity.<sup>15</sup>

The multisystem nature of fatty liver disease has important implications for patient management, including the development of multi-disciplinary care models.<sup>16</sup> A lack of specific

Keywords: Delphi method; Global health; Non-communicable disease (NCD); NAFLD/NASH; Steatotic liver disease.

Received 5 April 2023; received in revised form 21 April 2023; accepted 26 April 2023; available online 20 June 2023

\* Corresponding author. The City University of New York Graduate School of Public Health and Health Policy (CUNY SPH), New York, NY, USA.

E-mail address: [Jeffrey.Lazarus@sph.cuny.edu](mailto:Jeffrey.Lazarus@sph.cuny.edu) (J.V. Lazarus).

† Contributed equally as first authors

‡ Contributed equally as senior authors

<https://doi.org/10.1016/j.jhep.2023.04.035>



symptoms often leads to a clinically relevant delay in diagnosis.<sup>17,18</sup> Non-invasive tests (NITs) provide a practical and safe way of assessing fibrosis severity<sup>19</sup> and the risk of future liver-related events.<sup>20,21</sup> NITs have been used in the development of pathways to identify and stratify patients based on care needs,<sup>22–25</sup> yet such pathways are not implemented in the majority of healthcare settings.<sup>16</sup>

Several pharmacological treatments are under development, with some in late-stage clinical trials.<sup>26,27</sup> However, India is the only country where regulators have approved a pharmacological treatment for routine use in fatty liver disease (saroglitazar).<sup>28</sup> In the absence of pharmacological treatments approved for NASH, management is focused on improving insulin resistance and weight loss, when needed, and on attenuating the pro-inflammatory milieu of obesity, which are the predominant disease drivers.<sup>29,30</sup>

Few outside of the liver community recognise the need to deliver whole-of-society public health responses to fatty liver disease.<sup>31–33</sup> In 2021, a global consortium of experts set out consensus recommendations on how to accelerate public health action on this issue.<sup>34</sup> The negative impact of the COVID-19 pandemic on fatty liver disease risk factors<sup>35,36</sup> further reinforces the urgent attention this public health threat requires.

The liver health community (*i.e.*, the people and organisations who are largely working to improve liver health) must now build on past efforts to develop a clear vision and pathway to reduce the burden and address the individual and societal impacts of this growing challenge. Research is the central pivot for these efforts, accelerating the pace of knowledge creation and its translation into policy and practice.<sup>37–39</sup>

## Materials and methods

### Delphi expert panel member sample

The study's nine co-chairs used an iterative approach involving purposive, snowball, and targeted sampling to generate a large, global panel for this Delphi study. Based on publication record and engagement with the fatty liver disease agenda, the co-chairs identified 31 experts in clinical care (*e.g.*, liver, diabetes, obesity, and nutrition), public health, policy, advocacy, and patient representation, who collectively formed the core author group ( $n = 40$ ) (Table S1). The proposed survey panel ( $n = 473$ ) was created by

#### Box 1. The evolution of fatty liver disease nomenclature.<sup>12</sup>

The relationship between fatty liver disease, obesity, and diabetes was first described in the mid-1900s. Until then, the distinction between alcohol-related and non-alcoholic fatty liver disease (NAFLD) had been uncommon. In 1980, Ludwig *et al.* coined the term non-alcoholic steatohepatitis (NASH),<sup>12</sup> with NAFLD being a widely used umbrella term describing a histological spectrum ranging from steatosis without inflammation, to steatosis with varying degrees of inflammation and hepatocellular ballooning, which can lead to fibrosis, cirrhosis, and hepatocellular carcinoma. Recently, several proposals have been made to change the disease name, moving away from the construct of 'non-alcoholic' and better reflecting the metabolic foundation of the disease's aetiology. In 2023, a global consensus process is ongoing to reach agreement on the disease nomenclature. In this paper, we refer to fatty liver disease which arises in the absence of heavy alcohol intake and independently of other liver diseases.

compiling a list of known fatty liver disease experts from around the world with input from the core group (Fig. 1). Through this process, an expert panel diverse in demographic, disciplinary, and geographical characteristics was obtained (Table 1).

### Delphi statement domains


The development of the research priorities started with the core author group leading the development of evidence notes around seven topics, summarising the current knowledge base, envisioning what 'success' would look like in the next decade, identifying key questions, and suggesting research priorities for: (1) the human and economic burden, (2) defining and implementing models of care, (3) treatment and care, (4) education and awareness, (5) patient and community perspectives, (6) policy strategies and a societal approach, and (7) leadership for the fatty liver disease public health agenda. Twenty-six of the core-group members and 11 co-authors held a three-day meeting at Wilton Park, UK, in October 2022, co-chaired by H.E.M and opened by T.B and J.V.L, as part of the process. The research priorities were subsequently revised by J.V.L and H.E.M to reflect the Wilton Park discussions, and topics 6 and 7 were combined. The full core group received revised priorities for review in December 2022, with further revisions made based on core group feedback ahead of the first Delphi survey round (21 December 2022 to 15 January 2023).

### Delphi method data collection and analysis

The study design consisted of an in-person Wilton Park meeting (Table S2) and two survey rounds (R1 and R2) wherein panellists reviewed and voted on the research priorities. The study used the Qualtrics XM platform to develop and distribute the surveys (round duration ranged from 2 to 3.5 weeks), which included four-point Likert-type response categories for measuring the level of agreement with the draft research priorities (*i.e.*, 'agree', 'somewhat agree', 'somewhat disagree', and 'disagree'); the survey included a fifth 'not qualified to respond' option to accommodate the broad range of knowledge and expertise across panel participants. Panellists could provide comments and suggest edits to individual priorities in text boxes, which followed each of the statements. Both R1 and R2 included a text box allowing for overall comments at the end of each survey. Demographic data were collected from participants in R1.

An analytic team comprised of a sub-set of the core group (J.V.L, H.E.M, P.N.B, C.J.K, D.R, D.I.W, and M.V-R) reviewed the R1 data, including 600 open-ended comments from the panellists, and initiated draft revisions. The full core group then reviewed the revised priorities. In R2 (8–21 February 2023), panellists voted on the revised priorities, which were accompanied by text boxes summarising changes made based on panellist and core group input from R1. Panellists also ranked at least half of the priorities within each of the six domains; for domains with only three priorities panellists ranked all three.

Quantitative analysis of the R1 and R2 results included frequencies and proportions of the four response categories spanning 'agree' to 'disagree' for all research priority statements, as well as those selecting 'not qualified to respond'. For the final R2 Delphi results, we assigned each research priority statement with a grade to indicate the level of combined



**Fig. 1. Delphi panel generation and data collection.** Study methodology, including sample and data collection. Top, the iterative sampling approach used to generate a large and diverse Delphi panel (n = 344): nine co-chairs identified a core group of 31 experts in clinical care, public health, policy, and advocacy, who collectively formed the core authorship group (n = 40), working across 20 countries; the core group identified individuals with expertise in the fatty liver disease field. Bottom, the iterative digital data-collection process, including priority drafting (by the study co-chairs) and revision (by the co-chairs and other core authors) of the statements; one survey round (R1) of draft statements; analysis and revision of the statements by the core group; and a revised and final survey round (R2) of the consensus statements. R1 included text boxes for panellists to provide comments and suggest edits to individual statements; the final round (R2) allowed for overall comments at the end of each domain. For the final set of statements in R2, panellists ranked at least the top half of recommendations in each of the six domains. \*For domains with three priorities, panellists were required to rank them all. RR, response rate.

agreement (‘agree’ + ‘somewhat agree’), using a system that has been used in other Delphi studies<sup>34,40,41</sup> in which ‘U’ denotes unanimous (100%) agreement, ‘A’ denotes 90–99% combined agreement, ‘B’ denotes 78–89% combined agreement, and ‘C’ denotes 67–77% combined agreement. The data tables report the proportion who selected ‘not qualified to respond’, who were removed from the denominator to calculate the levels of agreement/disagreement from the relevant sample.

For the ranking, scores were calculated and normalised in Microsoft Excel (v.16.70) to compare rankings within each domain. Demographic data were analysed descriptively, including frequencies and proportions. No data were excluded from any analyses. Instances of missing data were totalled, and denominators were adjusted as applicable, for any calculations involving missing datapoints.

**Ethical considerations**

As this study does not include patients, patient data, or biological human samples, it received ethical review exemption from the Hospital Clínic of Barcelona, Spain, ethics committee on 19 December 2022. Each participant was asked to consent to participating in the study, prior to their inclusion. Adequate measures to ensure personal data protection and confidentiality have been taken and data were deidentified for all analyses.

**Results**

A total of 473 individuals were invited to participate in R1 and 344 (72.7%) completed the survey. These 344 respondents

were invited to participate in R2, of whom 288 (83.7%) completed the survey. Table 1 details the demographics of all expert panellists involved in the study. The mean age of respondents was 53.8 (standard deviation: 10.1). Most respondents were male (64.8%), worked in high-income countries (69.9%) and in the Europe and Central Asia region (42.2%), were primarily employed in the academic sector (66.6%), and worked in the clinical research field (79.4%). A total of 94 countries were represented in terms of respondent country of origin and 91 in terms of respondent country of work.

In R1, the study presented 31 initial research priorities to the panel. During revisions ahead of R2, three priorities were removed, with key components of these original statements being merged with existing priorities, leaving 28 priorities for the panel to review in R2. Across the two Delphi rounds, consensus increased for all six domains. The mean percentage of ‘agree’ responses across domains increased from 78.3 in R1 to 81.1 in R2, following the consideration of substantive comments received in R1.

Table 2 presents the final priorities, agreement grades, and rankings for each of the six domains. Within the final priorities in R2 (Fig. 2), the panel reached a unanimous combined agreement (‘agree’ + ‘somewhat agree’) with five priorities and >90% combined agreement with the remaining 23; the mean level of combined agreement across all priorities was 97.7%. For 13 priorities, ‘agree’ answers were below 80%, with higher reliance on ‘somewhat agree’ to achieve the high rate of overall combined agreement (Table S3).

**Table 1. Delphi panel characteristics (n = 344).**

Characteristic	n (%)
<b>Gender</b>	
Woman	115 (33.7)
Man	221 (64.8)
Non-binary or gender diverse	3 (0.9)
Prefer not to say	2 (0.6)
No response	3 (0.9)
<b>Age</b>	
All, mean [SD]	53.8 [10.1]
No response	12 (3.5)
<b>Country of origin, by income level (n = 94)</b>	
Low or middle	124 (36.9)
High	212 (63.1)
No response	8 (2.3)
<b>Global region<sup>a</sup> of origin</b>	
East Asia and Pacific	37 (11.0)
Europe and Central Asia <sup>b</sup>	142 (42.3)
Latin America and Caribbean	41 (12.2)
Middle East and North Africa	28 (8.3)
North America	52 (15.5)
South Asia	19 (5.7)
Sub-Saharan Africa	17 (5.1)
No response	8 (2.3)
<b>Country of work, by income level (n = 91)</b>	
Low or middle	102 (30.1)
High	237 (69.9)
No response	5 (1.5)
<b>Global region<sup>a</sup> of work</b>	
East Asia and Pacific	36 (10.6)
Europe and Central Asia <sup>c</sup>	143 (42.2)
Latin America and Caribbean	34 (10.0)
Middle East and North Africa	24 (7.1)
North America	76 (22.4)
South Asia	12 (3.5)
Sub-Saharan Africa	14 (4.1)
No response	5 (1.5)
<b>Primary sector of employment<sup>d</sup></b>	
Academic	229 (66.6)
Public	62 (18.0)
Private	38 (11.0)
Civil society	9 (2.6)
Other	3 (0.9)
No response	3 (0.9)
<b>Field(s) of employment<sup>d,e</sup></b>	
Clinical research	273 (79.4)
Non-clinical research	81 (23.5)
Healthcare provider	180 (52.3)
Patient/policy advocacy	36 (10.5)
Education	10 (2.9)
Other	7 (2.0)
No response	3 (0.9)
<b>Years working in fatty liver disease field</b>	
1 to 11	148 (43.7)
12 to 22	132 (38.9)
23 to 33	49 (14.5)
34 to 44	8 (2.4)
45 to 55	2 (0.6)
No response	5 (1.5)
<b>Publications authored focused on fatty liver disease</b>	
<6	103 (30.9)
6 to 25	95 (28.5)
26 to 50	54 (16.2)
51 to 100	42 (12.6)
>100	39 (11.7)
No response	11 (3.2)

(continued)

**Table 1. (continued)**

Characteristic	n (%)
<b>International or regional liver association membership(s)<sup>e</sup></b>	
AASLD	165 (48.0)
APASL	34 (9.9)
ALEH	30 (8.7)
EASL	191 (55.5)
Other	18 (5.2)
No membership	152 (44.2)
<b>Area of national professional association/society membership(s) in country of work<sup>e</sup></b>	
Liver disease	254 (73.8)
Gastroenterology	184 (53.5)
Obesity	42 (12.2)
Diabetes/endocrinology	45 (13.1)
Heart disease	11 (3.2)
Cancer	15 (4.4)
Primary care	5 (1.5)
Other	26 (7.6)
No membership	25 (7.3)

Percentages may sum to >100 due to rounding. Percentages for 'no response' are based on the total number of participants; all other percentages are calculated after excluding n of no response, unless otherwise indicated. AASLD, American Association for the Study of Liver Diseases; APASL, Asian Pacific Association for the Study of the Liver; ALEH, Asociación Latinoamericana para el Estudio del Hígado (Latin American Association for the Study of the Liver); EASL, European Association for the Study of the Liver.

<sup>a</sup>Based on World Bank regions.

<sup>b</sup>n = 3 participants are originally from Central Asia.

<sup>c</sup>n = 3 participants work in Central Asia.

<sup>d</sup>Denominator includes n of no response.

<sup>e</sup>Sum may exceed sample size as participants could choose >1 response.

*Patient and community perspectives* was one of two domains within the study where more than half of the research priority statements had <80% of the panel 'agree.' Five of six statements (5.1 and 5.3-5.6) illustrated higher reliance on 'somewhat agree' to achieve >90% combined agreement. *Leadership and public health policies* was the second domain where more than half of the research priority statements had <80% of the panel 'agree.' Two of three statements (6.1 and 6.3) illustrated higher reliance on 'somewhat agree' to achieve >90% combined agreement.

The priority rankings are explored in the discussion, alongside a summary of current evidence within each area.

## Discussion

This study engaged a multi-disciplinary group of experts and leaders from around the world to develop a consensus research agenda that is ambitious and transformational in nature and can deliver a step-change in how fatty liver disease is prevented and managed. To achieve this change, the study not only puts forward a shared research vision, but also illuminates degrees of agreement within the fatty liver disease community of practice, underscoring the benefit of continued discussion throughout the community. Here, we explore the importance of the research priorities for advancing the field with a focus on the highest ranked priorities across the six fatty liver disease research domains. The high response rates across both survey rounds and the substantial, often near-unanimous agreement of the panellists on all priorities suggests that agreeing on research priorities, for the first time in the fatty liver disease field, was a meaningful and important undertaking that builds on early priority-setting efforts.<sup>42,43</sup>



### Building the case for action: a better understanding of the human and economic burden

Knowledge of fatty liver disease has advanced tremendously over the past three decades, including on the predominant risk factors and disease drivers, yet gaps remain in our understanding of its natural history.<sup>44</sup> Most studies on its natural history and clinical progression emerge from tertiary centres, registries, or are based on biopsy availability,<sup>45–47</sup> which introduces disease-spectrum bias, while few are from prospective unbiased cohorts. As the field seeks to advance understanding, the panel specifically prioritised cohort studies that prospectively monitor outcomes in patients with defined liver disease (priority 1.2; ranked 1st in its domain); such cohorts will ideally be sampled from the general population.

In a separate but related priority, the panel unanimously supported the development and validation of risk prediction models to forecast progressive hepatic and extrahepatic outcomes (priority 1.4; ranked 3rd in its domain). The few risk-prediction models currently in use focus on broad risk factors, such as diabetes and body mass index. Future efforts will lead toward more nuanced predictors of outcomes, including novel biomarkers, as part of the quest towards precision medicine.<sup>43,48</sup>

While notable efforts have been made to establish the disease prevalence in both adults and children,<sup>1–3,48</sup> there are critical knowledge gaps in most countries and regions. The panel prioritised additional studies to better quantify the overall burden, including the disease prevalence and the quality of life impairment, in the general population and in high-risk groups (priority 1.1; ranked 2nd in its domain).

The panel almost universally agreed that additional studies are necessary to better quantify both the direct and indirect costs of fatty liver disease (priority 1.3). Available studies, while showing substantial costs associated with the disease,<sup>10,11,49,50</sup> are limited to a small number of high-income countries. Of equal importance to the panel was to advance understanding of the factors driving inequities in fatty liver disease (priority 1.5). The limited data in existence highlight large inequities by social group<sup>51</sup> and mirror what is seen more widely with non-communicable diseases (NCDs).<sup>52</sup> Further elucidating these factors will support the development of more targeted approaches to prevent and manage the burden of fatty liver disease.

### Advancing health system responses to fatty liver disease

#### *Defining and implementing multidisciplinary models of care*

One of the greatest challenges in clinical practice remains the identification of those with fatty liver disease and subsequent risk-stratification of those needing aggressive intervention and close monitoring by a specialist, from those who can be managed in primary care. Great advances have been made on the use of NITs in the past decades<sup>48,53,54</sup> and there is increasingly robust evidence of the value of current NITs for prognostication,<sup>20,21</sup> yet substantial challenges remain.<sup>55</sup> There is a critical unmet need when it comes to non-invasive approaches for monitoring disease progression and assessing disease resolution or meaningful improvements in fibrosis.<sup>48,56,57</sup> The panel unanimously recognised this imperative and gave a high priority to further validating NITs, with a focus on enabling early diagnosis, prognosis, and monitoring of

liver disease progression (priority 2.4; ranked 1st in its domain). Within this, emphasis will be needed on the cost-effectiveness of different approaches within different resource environments<sup>19,58,59</sup> and the appropriate cut-offs to be used in different settings and population groups.<sup>60</sup>

The multisystem nature of fatty liver disease, and the fact that many patients present with a range of comorbidities, requires a multi-disciplinary approach to management and care;<sup>16,61,62</sup> however, multi-disciplinary care models have not been widely adopted in most healthcare settings.<sup>16</sup> The panel unanimously called for further studies to determine the effectiveness of different models of care, including the impact on patient outcomes and their cost-effectiveness (priority 2.1; ranked 2nd in its domain). As this work advances, emphasis should be placed on care models that can be adaptable and implementable based on local resources. Specific considerations are needed around care models for paediatric populations (priority 2.2).<sup>63–66</sup>

Alongside the development of effective care models, the heterogeneity of patient presentation, coupled with the large burden of disease, means that tools are needed to support clinicians to identify those at highest risk of disease progression and adverse outcomes. The panel stressed the need to validate risk prediction models in different population groups, enabling them to be tailored to specific groups (priority 2.3; ranked 3rd in its domain).

The use of digital technologies in healthcare settings holds great potential for supporting service delivery,<sup>67</sup> yet this is a relatively new area of research within the field of fatty liver disease.<sup>68–71</sup> The panel supported the exploration of how novel digital technologies can be utilised within healthcare settings (priority 2.5) and further exploration of the potential for artificial intelligence methods to improve diagnosis of fatty liver disease (priority 2.7). The panel also supported further understanding how digital health approaches can support patients to achieve lifestyle behavioural change (priority 2.6). This work can build upon and complement broader efforts within the NCD and mental health fields.<sup>72,73</sup>

#### *Accelerating advances in fatty liver disease treatment*

In advancing treatment and care for affected populations, the panel highly ranked the importance of understanding the role of NITs in guiding treatment indication, response, and discontinuation, as well as predicting outcomes (priority 3.2; ranked 1st in its domain). This priority speaks to the current gap in evidence on the use of NITs to gain information on therapeutic responses.<sup>48,74</sup> As specific therapeutics are approved, NITs which can guide treatment decisions will be critical, especially given the likely long duration and high cost of treatment.

Currently, the central focus of treatment for fatty liver disease has been lifestyle interventions (e.g., nutrition, exercise, and weight loss), pharmacological treatment of comorbidities such as obesity and diabetes, and liver-directed therapies. The panel acknowledged that the prevention of fatty liver disease-related cirrhosis or HCC will require multi-pronged strategies which address an array of risk factors (e.g., social, environmental, behavioural, biological, and genetic) and called for studies to assess the efficacy and cost-effectiveness of such strategies (priority 3.1; ranked 2nd in its domain). As fatty liver disease treatment options expand, patient-centred decision

Table 2. Consensus statements for a fatty liver disease research priorities agenda.

Statement	Grade	Rank	A (%)	SA (%)	A+SA (%)	SD (%)	D (%)	NQ (%)	N	
<b>Domain 1: The human and economic burden</b>										
1.1	Implement studies to better quantify the fatty liver disease burden, including health-related quality of life, in the general population and in specific high-risk groups.	A	2	92.7	6.3	99.0	0.7	0.3	0.0	288
1.2	Conduct cohort studies to prospectively monitor outcomes in patients with defined liver disease phenotypes (e.g., NASH, NASH with fibrosis, cirrhosis, hepatocellular carcinoma).	U	1	93.0	6.6	99.7	0.3	0.0	0.3	287
1.3	Conduct additional studies on the socio-economic costs of fatty liver disease, capturing direct and indirect costs.	A		84.3	13.2	97.6	1.7	0.7	0.3	287
1.4	Develop and validate risk prediction models to forecast progressive hepatic and extrahepatic outcomes, to inform clinical decision making.	U	3	91.6	8.4	100.0	0.0	0.0	0.7	286
1.5	Report all data disaggregated by sex, race, ethnicity, age, socioeconomic status, education level, and other variables related to inequities.	A		74.2	23.3	97.6	2.1	0.3	0.3	287
<b>Domain 2: Defining and implementing models of care</b>										
2.1	Determine the effectiveness of different models of care for fatty liver disease, including their impact on patient outcomes and their cost-effectiveness.	U	2	90.6	9.0	99.7	0.0	0.3	0.0	288
2.2	Validate multidisciplinary models of care for fatty liver disease in paediatric populations.	A		80.7	17.9	98.6	1.4	0.0	2.8	280
2.3	Evaluate how risk prediction models for fatty liver disease perform in different populations, so that they can be tailored to specific populations and groups.	A	3	85.1	12.8	97.9	1.7	0.3	0.0	288
2.4	Validate non-invasive tests to enable early diagnosis, prognosis, and monitoring of liver disease progression.	U	1	93.4	6.3	99.7	0.3	0.0	0.0	288
2.5	Explore how novel digital technologies (e.g., artificial intelligence, data-based analytics, digital health applications and therapeutics) can be utilised within healthcare settings.	A		73.2	21.3	94.4	4.9	0.7	0.3	287
2.6	Assess how digital health (e.g., applications, interventions, therapeutics) can support patients to achieve lifestyle behavioural change.	A		69.8	26.7	96.5	2.8	0.7	0.0	288
2.7	Further explore the use of artificial intelligence to improve diagnostics for fatty liver disease.	A		63.9	31.2	95.1	3.5	1.4	1.0	285
<b>Domain 3: Treatment and care</b>										
3.1	Assess the efficacy and cost-effectiveness of multi-faceted strategies (e.g., social, environmental, behavioural, biological) to prevent fatty liver-related cirrhosis and hepatocellular carcinoma.	A	2	84.7	14.3	99.0	0.7	0.3	0.3	287
3.2	Study the role of non-invasive tests in guiding treatment indication, response, and discontinuation, as well as predicting outcomes.	A	1	94.1	5.2	99.3	0.3	0.3	0.0	288
3.3	Evaluate patient-centred decision making in relation to fatty liver disease treatment and care outcomes.	A		79.5	16.3	95.8	3.5	0.7	0.0	288
3.4	Evaluate the efficacy and cost-effectiveness of the optimal management of related diseases (e.g., diabetes, obesity) on fatty liver disease and other liver-related outcomes.	U	3	90.3	9.4	99.7	0.3	0.0	0.0	288
<b>Domain 4: Education and awareness</b>										
4.1	Conduct comparative population-based surveys to understand fatty liver disease knowledge amongst the general population and high-risk groups specifically, to inform the development of awareness-raising approaches.	A	2	82.2	14.6	96.9	2.1	1.0	0.3	287

(continued on next page)

Table 2. (continued)

Statement	Grade	Rank	A (%)	SA (%)	A+SA (%)	SD (%)	D (%)	NQ (%)	N
4.2 Conduct research to identify the educational needs of healthcare providers in key areas, such as primary care, diabetes/endocrinology, obesity medicine, and cardiology, about fatty liver disease.	A	1	89.6	9.4	99.0	0.7	0.3	0.0	288
4.3 Study the effectiveness of strategies to impact fatty liver disease knowledge, attitudes, beliefs, and practices (KABPs), prioritising KABPs among healthcare professionals and high-risk groups.	A	3	72.8	25.4	98.3	1.0	0.7	0.3	287
<b>Domain 5: Patient and community perspectives</b>									
5.1 Conduct research to understand the needs and experiences of fatty liver disease patients and at-risk communities (e.g., perspectives around prevention, treatment, and care, including mental health).	A	2	78.0	19.2	97.2	2.4	0.3	0.3	287
5.2 Study the impact of treatment and care on overall quality of life, including functional status (physical, psychological, social), in fatty liver disease patients.	A	1	81.9	17.4	99.3	0.7	0.0	0.0	288
5.3 Assess if published patient guidelines result in patients having an improved understanding of fatty liver disease and taking a more active role in their disease management.	A		71.9	25.7	97.6	2.1	0.3	0.0	288
5.4 Evaluate the efficacy of patient-led self-care programmes in improving fatty liver disease outcomes.	A		74.6	22.0	96.5	3.5	0.0	0.3	287
5.5 Explore the potential of new technologies (e.g., digital health applications and therapeutics, mobile interventions) to foster patient engagement in treatment and care.	A	3	75.5	21.3	96.9	2.4	0.7	0.7	286
5.6 Evaluate the effect of interventions to reduce liver disease stigma among patients, the public, and healthcare providers.	A		74.7	21.2	95.8	2.8	1.4	0.0	288
<b>Domain 6: Leadership and policies for the fatty liver disease public health agenda</b>									
6.1 Conduct periodic studies of national and sub-national policies and guidelines for the prevention and management of fatty liver disease, to identify trends and gaps, and assess their implementation.	A	1	79.4	18.8	98.3	1.0	0.7	0.3	287
6.2 Analyse policy successes and failures in addressing non-communicable diseases, to inform the development of fatty liver disease-specific strategies.	A	2	82.6	14.6	97.2	2.4	0.3	0.3	287
6.3 Monitor, study, and report mentions of fatty liver disease within patient groups and professional societies outside of the field of hepatology (e.g., at events, in publications).	A	3	66.8	28.0	94.8	3.8	1.4	0.7	286
<b>Mean % agreement</b>			81.1	16.6	97.7				

Percentages may add up to >100 due to rounding. Grades are based on the percentage of combined agreement ('agree' + 'somewhat agree'). U, unanimous (100%) agreement; A, 90–99% agreement. Responses to each statement are presented as percentages of the total responses. A, agree; SA, somewhat agree; SD, somewhat disagree; D, disagree; NQ, the percentage of participants that indicated that they were not qualified to respond; N, total number of responses; NASH, non-alcoholic steatohepatitis.

making will also become increasingly important (priority 3.3), as seen more broadly in the NCD field.<sup>75,76</sup>

Recognising the shared metabolic risks inherent in both fatty liver disease and other highly prevalent co-morbidities, the panel unanimously highlighted the importance of further evaluating the efficacy and cost-effectiveness of optimal management of related diseases (e.g., diabetes and obesity) on liver-related outcomes (priority 3.4; ranked 3rd in its domain). These factors increase risk of fibrosis progression, which in turn increases all-cause mortality, with the majority of patients succumbing to cardiovascular disease and solid organ malignancy,<sup>46,77</sup> but further evidence is needed that optimal management of these comorbidities has a beneficial impact on

hepatic outcomes.<sup>78</sup> This evidence will be critical in ongoing efforts to engage and involve primary care and non-liver specialities within the multidisciplinary management of fatty liver disease.

*Improving knowledge and awareness*

Despite fatty liver disease being highly prevalent, awareness is generally low amongst non-liver health specialists – most importantly primary care physicians and diabetologists – with critical knowledge gaps around risk-factors, diagnosis, and management approaches<sup>79–81</sup> and a lack of tools to support clinical decision making.<sup>79</sup> While important progress is being made in this area,<sup>82</sup> the panel gave prominence to research

aimed at identifying the educational needs of healthcare providers in key areas, including primary care, diabetes, and obesity (priority 4.2; ranked 1st in its domain), to inform targeted educational strategies. Studies will also be needed to assess the effectiveness of such strategies to impact knowledge, attitudes/beliefs, and practices, starting with key healthcare professionals (priority 4.3; ranked 3rd in its domain).

Many people living with fatty liver disease are unaware of their fibrosis stage, which has important implications for adherence to management approaches.<sup>83</sup> Equally, within population groups at high risk of disease progression, including people with type 2 diabetes mellitus and other metabolic diseases, awareness of fatty liver disease, the health risks posed by it, or how it interacts with their other diseases, is low.<sup>84,85</sup> To inform the development of awareness-raising approaches, the panel suggests that comparative population-based surveys are implemented to understand knowledge amongst the general population and high-risk groups (priority 4.1; ranked 2nd in its domain).

### Delivering whole-of-society responses

#### *Patient and community perspectives*

The large prevalence of fatty liver disease and the less severe effects of steatosis potentially masks that, in advanced stages, the disease causes substantial impairment in quality of life which is often compounded in those with multiple morbidities.<sup>6–8</sup> Alongside liver-related outcomes, fatigue and depression are important contributors to reduced quality of life in people living with fatty liver disease.<sup>86,87</sup>

While combined agreement on patient-centred orientations was high, the study's results noted above suggest that further discussion within the liver health community on research

priorities for patient and community engagement is warranted. The panel felt strongly about the need to study the impact of treatment and care on overall quality of life, including functional status (priority 5.2; ranked 1st in its domain). Knowledge gaps remain about patient needs and experiences,<sup>88</sup> including perspectives around prevention, treatment, and care. The panel prioritised studies that can advance this understanding (priority 5.1; ranked 2nd in its domain).

In chronic disease management, engaged patients are shown to have better outcomes,<sup>89</sup> while lower engagement levels are associated with more adverse events.<sup>90</sup> Digital approaches have proven effective at improving patient engagement in NCD management<sup>91,92</sup> and initial efforts have been made to understand the role of technologies in fostering patient engagement in fatty liver disease treatment and care.<sup>93,94</sup> The panel prioritised further exploration in this novel area of research (priority 5.5; ranked 3rd in its domain). In the area of patient engagement, the panel also agreed that assessing the impact of patient care guidelines<sup>95</sup> will be important (priority 5.3).

In other areas, the panel agreed on further evaluating the efficacy of patient-led self-care programmes in improving fatty liver disease outcomes (priority 5.4) and evaluating the effect of interventions to reduce liver disease stigma among patients, the public, and healthcare providers (priority 5.6).

### Leadership and public health policies

From the local to the global level, public health policy responses to fatty liver disease have, to date, not stemmed the increase in fatty liver disease morbidity or mortality.<sup>31,34,96</sup> A global review of policies, strategies, and guidelines conducted in 2020 found that of 102 countries assessed, around one-third




Fig. 2. Research priorities agenda for fatty liver disease.



of countries scored zero on an associated preparedness index.<sup>96</sup> Fatty liver disease is also absent in otherwise broad, global strategies and guidelines, including the World Health Organization's NCD strategies.<sup>97,98</sup> Building on past efforts, the panel near unanimously called for periodic studies to assess national and sub-national policies and guidelines for the prevention and management of fatty liver disease, to identify trends and gaps and assess their implementation (priority 6.1; ranked 1st in its domain).

Agreement was also reached on the need to analyse policy successes and failures in addressing NCDs, to inform the development of fatty liver disease-specific strategies (priority 6.2; ranked 2nd in its domain). This will require a greater focus within the liver health community on the commercial determinants driving the increasing burden of fatty liver disease.<sup>99</sup>

The multidisciplinary and multi-sectoral nature of the public health responses to fatty liver disease require further engagement and collaboration with those outside of the field of hepatology. In an effort to guide and inform such engagements, and assess their impact, the panel recommends efforts to monitor, study, and report mentions of fatty liver disease within patient groups and professional societies (priority 6.3; ranked 3rd in its domain).

### Study strengths and limitations

The major strength of this study is its novelty as the first global effort to propose a comprehensive research agenda for fatty liver disease utilising the rigorous Delphi consensus methodology. Additionally, within this methodology, the ability to illustrate degrees of agreement by breaking-out 'agree' and 'somewhat agree' responses may assist decision makers and researchers. We suggest that the breadth of issues covered, combined with the relatively focused priorities, makes the

outcome both aspirational and practical. While this process did not consider how to operationalise these research priorities, including the resource requirements to do so, the findings can guide the investment decisions of research funders.

The Delphi methodology used in this study is an effective approach in consensus building, yet building consensus is not without challenges. In this study we used purposive sampling to develop a core group. To mitigate the biases of purposive sampling, we used snowballing and targeted sampling to yield a panel of 344 people diverse in both expertise and geographical representation. We do, however, acknowledge that the characteristics of the final group (e.g., predominantly based in high-income countries and employed in the academic sector), will have been reflected within the agreement levels of the research priorities. For instance, the lower levels of agreement on the more patient centric and policy-oriented priorities likely reflects the smaller proportion of the panel working in patient/policy advocacy. While 10.5% (n = 36) of the panel reported some engagement in patient/policy advocacy, this was the primary area of work for only 4.7% (n = 16). Conducting the survey in the English language may have also influenced who accepted the invitation to contribute.

### Conclusions

Delivering comprehensive health system and public health responses to fatty liver disease will require the global health community to re-envision the landscape, grow the fatty liver disease community of practice, and place greater emphasis on collective and collaborative thinking and action. This global multidisciplinary effort has, for the first time, developed a consensus fatty liver disease research agenda that can serve as the foundation for turning the tide on this silent public health threat.

### Affiliations

<sup>1</sup>Barcelona Institute for Global Health (ISGlobal), Hospital Clínic, University of Barcelona, Barcelona, Spain; <sup>2</sup>Faculty of Medicine and Health Sciences, University of Barcelona, Barcelona, Spain; <sup>3</sup>CUNY Graduate School of Public Health and Health Policy (CUNY SPH), New York, NY, USA; <sup>4</sup>European Association for the Study of the Liver (EASL), Geneva, Switzerland; <sup>5</sup>Independent Consultant, Nottingham, UK; <sup>6</sup>Division of Gastroenterology and Hepatology, Department of Medicine, Mayo Clinic, Rochester, MN, USA; <sup>7</sup>Division of Gastroenterology, Department of Medicine, Schulich School of Medicine, Western University & London Health Sciences Centre, London, Ontario, Canada; <sup>8</sup>Department of Epidemiology and Biostatistics, Schulich School of Medicine, Western University, London, Ontario, Canada; <sup>9</sup>Department of Gastroenterology, School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile; <sup>10</sup>Aix Marseille Univ, Inserm, IRD, SESSTIM, Sciences Economiques & Sociales de la Santé & Traitement de l'Information Médicale, ISSPAM, Marseille, France; <sup>11</sup>Houston Methodist Hospital, Houston Research Institute, Houston, TX, USA; <sup>12</sup>Barts Liver Centre, Blizard Institute, Queen Mary University of London, London, UK; <sup>13</sup>Fatty Liver Program, Arizona Liver Health, Phoenix, AZ, USA; <sup>14</sup>King Faisal Specialist Hospital & Research Centre, Riyadh, Saudi Arabia; <sup>15</sup>Liver Unit, Hospital Clínic, University of Barcelona, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain; <sup>16</sup>Division of Hepatology, Department of Medicine II, Leipzig University Medical Center, Leipzig, Germany; <sup>17</sup>Division of Hepatology, University of Dundee, Dundee, Scotland, UK; <sup>18</sup>Multivisceral Transplant Unit-Gastroenterology, Department of Surgery, Oncology and Gastroenterology at the Padua University Hospital, Padua, Italy; <sup>19</sup>Department of Hepatology and Transplant, Hospital Médica Sur, Mexico City, Mexico; <sup>20</sup>Department of Gastroenterology, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico; <sup>21</sup>Asociación Latinoamericana para el Estudio del Hígado (ALEH), Santiago, Chile; <sup>22</sup>Clinica Universitária de Gastrenterologia, Laboratório de Nutrição, Faculdade de Medicina, Universidade de Lisboa, Lisbon, Portugal; <sup>23</sup>Division of Endocrinology, Diabetes & Metabolism, Department of Medicine, University of Florida, Gainesville, FL, USA; <sup>24</sup>Greek Patients Association, Athens, Greece; <sup>25</sup>Department of Hepatology, Postgraduate Institute of Medical Education and Research, Chandigarh, India; <sup>26</sup>Department of Gastroenterology and Hepatology, Antwerp University Hospital, Edegem, Belgium; <sup>27</sup>InflaMed Centre of Excellence, Laboratory for Experimental Medicine and Paediatrics, Translational Sciences in Inflammation and Immunology, Faculty of Medicine and Health Sciences, University of Antwerp, Wilrijk, Belgium; <sup>28</sup>Department of Medicine, Huddinge, Karolinska Institutet, Stockholm, Sweden; <sup>29</sup>CUNY Center for Systems and Community Design and NYU-CUNY Prevention Research Center, New York, NY, USA; <sup>30</sup>Kautz 5 gUG, Köln, Germany; <sup>31</sup>Independent Researcher, Ponte de Lima, Portugal; <sup>32</sup>Department of Gastroenterology and Hepatology, Odense University Hospital, Odense, Denmark; <sup>33</sup>University California Berkeley School of Public Health, Berkeley, CA, USA; <sup>34</sup>National Institute for Health Research Birmingham Biomedical Research Centre at University Hospitals Birmingham NHS Foundation Trust and the University of Birmingham, Birmingham, UK; <sup>35</sup>Department of Medicine, University of Chicago, Chicago, IL, USA; <sup>36</sup>Department of Community Health and Social Sciences, CUNY Graduate School of Public Health and Health Policy, New York, NY, USA; <sup>37</sup>Department of Hepatology, Institute of Liver and Biliary Sciences, New Delhi, India; <sup>38</sup>Hepatology and Clinical Research Units, Hospital Universitario Austral, Buenos Aires, Argentina; <sup>39</sup>Division of Hepatology, Department of Medicine, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa; <sup>40</sup>UCL Institute for Liver and Digestive Health, Royal Free Hospital, London, UK; <sup>41</sup>Precision Medicine, Biological Resource Center, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy; <sup>42</sup>Department of Pathophysiology and Transplantation, University of Milan, Milan, Italy; <sup>43</sup>School of Public Health, Faculty of Social Welfare and Health Sciences, University of Haifa, Haifa, Israel; <sup>44</sup>Department of Gastroenterology, Tel Aviv Medical Centre, Tel Aviv, Israel; <sup>45</sup>Metabolic Liver Research Program, I. Department of Medicine, University Medical

Centre Mainz, Mainz, Germany;<sup>46</sup>The Chinese University of Hong Kong, Hong Kong;<sup>47</sup>Center for Liver Disease, Inova, Falls Church, VA, USA

## Abbreviations

HCC, hepatocellular carcinoma; NAFLD, non-alcoholic fatty liver disease; NASH, non-alcoholic steatohepatitis; NCD, non-communicable disease; NIT, non-invasive test; R, round.

## Financial support

The data collection and analysis were funded by the European Association for the Study of the Liver (EASL), with support from Takeda, MSD, Bristol-Myers-Squibb Company, and Apollo Endo.

## Conflict of interest

J.V.L acknowledges grants and speaker fees from AbbVie, Gilead Sciences, MSD, and Roche Diagnostics to his institution, speaker fees from Intercept, Janssen, Novo Nordisk, and ViiV, and consulting fees from Novavax, outside of the submitted work. He also acknowledges support to ISGlobal from the grant CEX2018-000806-S funded by MCIN/AEI/10.13039/501100011033 and the Generalitat de Catalunya through the CERCA Programme, outside of the submitted work. H.E.M acknowledges consultancy fees from the European Association for the Study of The Liver (EASL) related to the current manuscript. A.M.A acknowledges grant support to her institution from the National Institutes of Health (NIH) (DK128127), Novo Nordisk, Pfizer, and Target Pharma and advisory board participation for Novo Nordisk, outside of the submitted work. M.N acknowledges consulting fees from and advisory board participation for Altimune, BI, BMS, Cytodyn, 89BIO, EchoSens, Gilead, GSK, Madrigal, Merck, Novo Nordisk, OWL, Prespecturm, Pfizer, Roche Diagnostics, Siemens, Terns, and Takeda, payment or honoraria from Madrigal, Echosens, and Sonic Incytes, patents from AI in NAFLD, roles as AASLD Vice-chair of NAFLD SIG and AE at CGH, and stock from Anaetos Rivus Pharma, CIMA, ChronWell, and Viking, outside of the submitted work. W.A has received fees for consulting and teaching or competitively awarded grant funding from Gilead Sciences, GlaxoSmithKline, and AstraZeneca, outside of the submitted work. N.A acknowledges grants or contracts from 89Bio, AbbVie/Allergan, Akero, Better Therapeutics, Boehringer Ingelheim, Bristol-Myers Squibb, Corcept, DSM, Galectin, Genentech, Genfit, Gilead, Hepagene, Healio, Intercept, Inventiva, Ionis, Madrigal, Merck, NGM, Noom, NorthSea, Novo Nordisk, Perspectrum, Pfizer, Poxel, Viking, and Zydus, consulting fees from AbbVie/Allergan, Echosens, Fibronostics, Gilead, Intercept, Madrigal, Novo Nordisk, Perspectrum, Pfizer, and Zydus, and payment or honoraria from AbbVie/Allergan, Alexion, Echosens, Eisai, Exelixis, Gilead, Intercept, Perspectrum, Salix, and Theratechnologies, outside of the submitted work. R.B acknowledges payment or honoraria and support for attending meetings and/or travel from AbbVie, outside of the submitted work. T.B acknowledges grants to or contracts with his institution from AbbVie, BMS, Gilead, MSD/Merck, Humedics, Intercept, Merz, Norgine, Novartis, Orphanal, and Sequana Medical, consulting fees from AbbVie, Alexion, Bayer, Gilead, GSK, Eisai, Enyo Pharma, HepaRegeniX GmbH, Humedics, Intercept, Ipsen, Janssen, MSD/Merck, Novartis, Orphanal, Roche, Sequana Medical, SIRTEX, SOBI, and Shionogi, payment or honoraria from AbbVie, Alexion, Bayer, Gilead, Eisai, Falk Foundation, Intercept, Ipsen, Janssen, MedUpdate GmbH, MSD/Merck, Novartis, Orphanal, Sequana Medica, SIRTEX, and SOBI, and support for attending meetings and/or travel from Gilead, AbbVie, Intercept, and Janssen, outside of the submitted work. P.N.B acknowledges consulting fees from Resolution Therapeutics and payment or honoraria from Takeda, outside of the submitted work. P.B acknowledges payment or honoraria from Gilead and Chiesi Farmaceutici, support for attending meetings and/or travel from Alpha Wasserman, advisory board participation for Biotech, Kedrion, Astellas, Gilead, Sandoz, and Chiesi Farmaceutici, and a preceptorship for Novartis, outside of the submitted work. H.C-P acknowledges consulting fees from Orphanal and Novo Nordisk and lecture fees from and advisory board participation for Roche Portugal and EISAI, outside of the submitted work. K.C acknowledges grant support to his institution from Echosens, Inventiva, Poxel, Novo, Labcorp, and Zydus, consulting fees from BMS, Lilly, Madrigal, Merck, Myovant, Novo Nordisk, Prosciento, Quest, Sagimet, Sonic Incytes, and Terns, and other support from Echosens, Inventiva, Poxel, Labcorp, and Zydus, outside of the submitted work. N.D acknowledges roles as Chair of the Greek Patients Association and the Positive Voice (PLHIV Association), outside of the submitted work. A.D acknowledges an unpaid role as Secretary-General of the Indian National Association for the Study of the Liver (INASL), outside of the submitted work. S.M.F acknowledges grants or contracts from Research Foundation Flanders, Astellas, Falk Pharma, Genfit, Gilead Sciences, GlympsBio, Janssens Pharmaceutica, Inventiva, Merck Sharp & Dome, Pfizer, and Roche, consulting fees from AbbVie, Actelion, Aelin Therapeutics, Aligos Therapeutics, Allergan, Astellas, Astra Zeneca, Bayer, Boehringer

Ingelheim, Bristol-Meyers Squibb, CSL Behring, Coherus, Echosens, Eisai, Enyo, Galapagos, Galmed, Genetech, Genfit, Gilead Sciences, Intercept, Inventiva, Janssens Pharmaceutical, Julius Clinical, Madrigal, Medimmune, Merck Sharp & Dome, NGM Bio, Novartis, Novo Nordisk, Promethera, and Roche, and lecture fees from AbbVie, Allergan, Bayer, Eisai, Genfit, Gilead Sciences, Janssens Cilag, Intercept, Inventiva, Merck Sharp & Dome, Novo Nordisk, and Promethera, outside of the submitted work. H.H acknowledges grants or contracts from Astra Zeneca, Pfizer, MSD, EchoSens, and Gilead, consulting fees from Astra Zeneca, and payment or honoraria from MediPlast, outside of the submitted work. T.T-K.H acknowledges grants or contracts from the Centers for Disease Control and Prevention, outside of the submitted work. Al.K acknowledges grants or contracts from EU Horizon 20, the Novo Nordisk Foundation, Innovation Fund Denmark, the Danish National Research Foundation, the Region of Southern Denmark, and AstraZeneca, royalties or licenses from Gyldendal, payment or honoraria from Norgine, Siemens, and Nordic Bioscience, patents from the Region of Southern Denmark and the University of Southern Denmark, advisory board participation for Norgine and Siemens, an unpaid role as Vice Secretary of the European Association for the Study of The Liver (EASL), and other support from Norgine, Siemens, Echosens, and Nordic Bioscience, outside of the submitted work. V.M acknowledges grants from 89bio, Akero Therapeutics Inc., Albireo Pharma Inc., Aligos Therapeutics, Alimentiv, AlloVir, Altimune Inc., AMRA Medical AB, Arrowhead Pharmaceuticals, ChemomAb Ltd., Covance Inc., Cymabay Therapeutics, E-Scopics, EA Pharma Co. LTD, Echosens, Eli Lilly & Company, ENYO Pharma SA, Galectin Therapeutics, Gilead Sciences Inc., GlaxoSmithKline PLC, HepQuant, High Tide Therapeutics, HistoIndex, Intercept Pharmaceuticals, Inventiva Pharma, Janssen, LabCorp, LG Chem Life Sciences, Merck & Company Inc., Northsea Therapeutics, Novartis Pharma AG, Novo Nordisk, Oncoustics, Pharmanest, Pliant Therapeutics, and Regeneron Pharmaceuticals to the Forum for Collaborative Research, outside of the submitted work. P.N.N acknowledges grants or contracts from Novo Nordisk, consulting fees from Novo Nordisk, Boehringer Ingelheim, Gilead, Intercept, Poxel Pharmaceuticals, BMS, Pfizer, Sun Pharma, Madrigal, and GSK, payment or honoraria from Novo Nordisk and AiCME, support for attending meetings and/or travel from Novo Nordisk, and advisory board participation for Novo Nordisk, Boehringer Ingelheim, Gilead, Intercept, Poxel Pharmaceuticals, BMS, Pfizer, Sun Pharma, Madrigal, and GSK, outside of the submitted work. M.E.R acknowledges consulting fees from Boehringer Ingelheim, Novo Nordisk, GSK, Intercept, Madrigal, and Cytodyn, outside of the submitted work. D.R. acknowledges remuneration to her institution from the European Association for the Study of The Liver (EASL) related to the current manuscript. M.S acknowledges grants to his institution from Inventiva, Zydus, and Merck, advisory board participation for Zydus, and unpaid leadership and membership roles for ALEH and the Global NASH Council, outside of the submitted work. C.W.S acknowledges payment or honoraria from Gilead Sciences and Abbott, support for attending meetings and/or travel from Gilead Sciences, and roles as Co-chair of the Society on Liver Disease of Africa and member of the International Advisory Board of *The Lancet Gastroenterology and Hepatology* journal, outside of the submitted work. E.A.T acknowledges consulting fees from Novo Nordisk, Boehringer, and Pfizer, payment or honoraria from Novo Nordisk and Dr Falk, board participation for Boehringer, Pfizer, and Novo Nordisk, and a role a member of the EASL Governing Board, outside of the submitted work. L.V acknowledges grants or contracts and support for attending meetings and/or travel from Gilead Sciences, consulting fees from Gilead, Pfizer, Astra Zeneca, Novo Nordisk, Intercept pharmaceuticals, Diatech Pharmacogenetics, IONIS, and Boehringer Ingelheim, payment or honoraria from MSD, Gilead, AlfaSigma, AbbVie, and Viatrix, patents from Takeda, and board participation for Intercept, Pfizer, Gilead, Novo Nordisk, and Boehringer Ingelheim, outside of the submitted work. M.V-R acknowledges consultancy fees from the European Association for the Study of The Liver (EASL) related to the current manuscript. S.Z-S has given presentations for and received support for attending meetings and/or travel from AbbVie, outside of the submitted work. J.M.S acknowledges research funding from Gilead Sciences, Boehringer Ingelheim, and Siemens Healthcare GmbH, consulting fees from Apollo Endosurgery, Albireo Pharma Inc, Bayer, BMS, Boehringer Ingelheim, Gilead Sciences, GSK, Intercept Pharmaceuticals, Ipsen, Inventiva Pharma, Madrigal, MSD, Northsea Therapeutics, Novartis, Novo Nordisk, Pfizer, Roche, Sanofi, and Siemens Healthcare GmbH, payment or honoraria from Boehringer Ingelheim, Echosens, MedPublico GmbH, Novo Nordisk, and Madrigal Pharmaceuticals, support for attending meetings and/or travel from Gilead Sciences, and stock from AGED diagnostics and Hepta Bio, outside of the submitted work. V.W-S.W acknowledges grants from Gilead Sciences to his institution, consulting fees from AbbVie, Boehringer Ingelheim, Echosens, Gilead Sciences, Intercept, Inventiva, Novo Nordisk, Pfizer, Sagimet Biosciences, and TARGET PharmaSolutions, payment and honoraria from Abbott,

AbbVie, Gilead Sciences, Novo Nordisk, and Unilab, support for attending meetings and/or travel from AbbVie and Gilead Sciences, and unpaid stock as co-founder of Illuminatio Medical Technology Limited, outside of the submitted work. Z.M.Y acknowledges consulting fees from Gilead, Intercept, Siemens, Novo Nordisk, Madrigal, Merck, Quest, and Bristol Myers Squibb, outside of the submitted work. J.P.A, P.C, S.A.A, M.A, G.C-N, D.I.W, A.K, C.J.K, and S.K.S have no conflicts of interest to disclose.

Please refer to the accompanying ICMJE disclosure forms for further details.

### Authors' contributions

This study was led by a core group of 40 co-authors. J.V.L led the core group and provided regular updates by email. Twenty-six core group members and 11 co-authors participated in a three-day in-person meeting hosted by Wilton Park, UK, in October 2022, which informed the development of the research priorities included in the Delphi study. Seven co-chairs (A.M.A, J.P.A, P.C, M.N, J.M.S, V.W-S.W, and Z.M.Y) led the drafting of seven evidence notes, including key priorities and challenges, ahead of the Wilton Park meeting and were supported by core group members (R.B, T.B, H.C-P, K.C, N.D, A.D, T.T-K.H, A.I.K, V.M, P.N.N, M.E.R, M.S, E.T, and S.Z-S). The evidence notes were reviewed by J.V.L and H.E.M and informed the drafting of the research priorities statements and actions. D.R and J.V.L led the methodology. J.V.L, H.E.M, P.N.B, C.J.K, D.R, D.I.W, and M.V-R reviewed comments submitted as part of the two survey rounds. J.V.L, H.E.M, and M.V-R reviewed all comments sent directly by email. All panel members provided two rounds of comments through Qualtrics XM. H.E.M, M.V-R, and J.V.L wrote the first draft of the manuscript, which was reviewed by the core group. Those fulfilling authorship criteria are named.

### Data availability statement

De-identified source data for all analyses will be made available by contacting the corresponding author (Jeffrey.Lazarus@sph.cuny.edu), with appropriate ethical approval and for fair use.

### Acknowledgments

The formation of the Healthy Livers, Healthy Lives global coalition builds on three years of work led initially by the EASL International Liver Foundation (EILF) and since 2021 by EASL. Collaboration has been at the centre of this work. Over 500 individuals and organisations spanning over 100 countries have engaged in these efforts, with multiple disciplines and sectors represented, including affected populations. As part of this process, in October 2022, delegates, including representatives from EASL, the American Association for the Study of Liver Diseases (AASLD), the Latin American Association for the Study of the Liver (*Asociación Latinoamericana para el Estudio del Hígado [ALEH]*) (representatives of the Asian Pacific Association for the Study of the Liver [APASL] were unable to attend), the Society on Liver Disease in Africa (SOLDA), the European Association for the Study of Diabetes (EASD), the European Association for the Study of Obesity (EASO), the European Society of Primary Care Gastroenterology (ESPCG), the Global NASH Council (GNC), United European Gastroenterology (UEG), the World Obesity Federation (WOF), and the World Organization of Family Doctors (WONCA), gathered for three days of discussion at Wilton Park, a UK-based forum for strategic dialogue.

We are grateful to the following experts for their contributions to the Delphi process: Manal F. Abdelmalek, Abdelmounem E. Abdo, Matthew J. Armstrong, Carol L. Brosgart, José Luis Calleja, Michael R. Charlton, Yogesh Chawla, Abhijit Chowdhury, David E. Cohen, Alessandro Demaio, Javier Diaz-Ferrer, Anna Mae Diehl, Alexander French, Cheryl Grainger, Nadege T. Gunn, Bela Hunyady, Robert James, Jia-Horng Kao, Rohit Kohli, Hye Won Lee, Cosmas Rinaldi A. Lesmana, Panu K. Luukkonen, Dina Mansour, Patrick Marcellin, Rui T. Marinho, Luca Miele, Tahiri Mohammed, Munkhjargal Ayurzana, Jude Oben, Missiani Ochwoto, Juan Paredes Méndez, Rachel Pryke, Puneet Puri, Marcus Ranney, Stuart K. Roberts, John F. Ryan, Arun J. Sanyal, Raymond Sayegh, Piotr Socha, Joan B. Soriano, Juan José Suárez, Ki-Chul Sung, Tawesak Tanwandee, Juan Turnes, Saskia van Mil, Kym Watt, Sarah H. Wild, Stavra A. Xanthakos, and Amany Zekry.

### The Healthy Livers, Healthy Lives Collaborators

Jeffrey V. Lazarus<sup>1,2,3</sup>, Henry E. Mark<sup>4,5</sup>, Alina M. Allen<sup>6</sup>, Juan Pablo Arab<sup>7,8,9</sup>, Patrizia Carrieri<sup>10</sup>, Mazen Noureddin<sup>11</sup>, William Alazawi<sup>12</sup>, Naim Alkhouri<sup>13</sup>, Saleh A. Alqahtani<sup>14</sup>, Marco Arrese<sup>9</sup>, Ramon Bataller<sup>15</sup>, Thomas Berg<sup>16</sup>, Paul N. Brennan<sup>17</sup>, Patrizia Burra<sup>18</sup>, Graciela E. Castro-Narro<sup>19,20,21</sup>, Helena Cortez-Pinto<sup>22</sup>, Kenneth Cusi<sup>23</sup>, Nikos Dedes<sup>24</sup>, Ajay Duseja<sup>25</sup>, Sven M. Francke<sup>26,27</sup>, Hannes Hagström<sup>28</sup>, Terry T-K. Huang<sup>3,29</sup>, Dana Ivancovsky Wajcman<sup>1</sup>, Achim

Kautz<sup>30</sup>, Christopher J. Kopka<sup>31</sup>, Aleksander Krag<sup>32</sup>, Veronica Miller<sup>33</sup>, Philip N. Newsome<sup>34</sup>, Mary E. Rinella<sup>35</sup>, Diana Romero<sup>36</sup>, Shiv Kumar Sarin<sup>37</sup>, Marcelo Silva<sup>38</sup>, C. Wendy Spearman<sup>39</sup>, Emmanuel A. Tsochatzidis<sup>40</sup>, Luca Valenti<sup>41,42</sup>, Marcela Villota-Rivas<sup>1</sup>, Shira Zelber-Sag<sup>43,44</sup>, Jörn M. Schattenberg<sup>45</sup>, Vincent Wai-Sun Wong<sup>46</sup>, Zobair M. Younossi<sup>47</sup>, Fredrick Aberg<sup>48</sup>, Leon A. Adams<sup>49</sup>, Khalid Al-Naamani<sup>50</sup>, Reda M. Albadawy<sup>51</sup>, Zinaida Alexa<sup>52</sup>, Michael Allison<sup>53</sup>, Faisal Abdullatif Alnaser<sup>54</sup>, Khalid Alswat<sup>55</sup>, Mario R. Alvares-da-Silva<sup>56</sup>, Domenico Alvaro<sup>57</sup>, Michele Alves-Bezerra<sup>58</sup>, Raul J. Andrade<sup>59</sup>, Quentin M. Anstee<sup>60</sup>, Yaw Asante Awuku<sup>61</sup>, Oidov Baatarkhuu<sup>62</sup>, Gyorgy Baffy<sup>63</sup>, Shokhista R. Bakieva<sup>64</sup>, Meena B. Bansal<sup>65</sup>, Robert Barouki<sup>66</sup>, Rachel L. Batterham<sup>67</sup>, Cynthia Behling<sup>68</sup>, Renata Belfort-DeAguiar<sup>69</sup>, Annalisa Berzigotti<sup>70</sup>, Michael Betel<sup>71</sup>, Cristiana Bianco<sup>72</sup>, Emanuele Bosi<sup>73</sup>, Jerome Boursier<sup>74</sup>, Elizabeth M. Brunt<sup>75</sup>, Elisabetta Bugianesi<sup>76</sup>, Christopher J. Byrne<sup>77</sup>, Maria Cecilia Cabrera Cabrejos<sup>78</sup>, Stephen Caldwell<sup>79</sup>, Rotonya Carr<sup>80</sup>, Marlen Ivón Castellanos Fernández<sup>81</sup>, Laurent Castera<sup>82</sup>, Maria Gabriela Castillo-López<sup>83</sup>, Cyrielle Caussy<sup>84</sup>, Eira Cerda-Reyes<sup>85</sup>, Antonio Ceriello<sup>86</sup>, Wah- Kheong Chan<sup>87</sup>, Yoosoo Chang<sup>88</sup>, Phunchai Charatcharoenwitthaya<sup>89</sup>, Norberto Chavez-Tapia<sup>90</sup>, Raymond T. Chung<sup>91</sup>, Massimo Colombo<sup>92</sup>, Kirsten J. Coppell<sup>93</sup>, Helma P. Cotrim<sup>94</sup>, Antonio Craxi<sup>95</sup>, Javier Crespo<sup>96</sup>, Anuradha Dassanayake<sup>97</sup>, Nicholas O. Davidson<sup>98</sup>, Robert J. de Knegt<sup>99</sup>, Victor de Ledinghen<sup>100</sup>, Münevver Demir<sup>101</sup>, Hailemichael Desalegn<sup>102</sup>, Moises Diago<sup>103</sup>, John F. Dillon<sup>77</sup>, Bruce Dimmig<sup>104</sup>, M. Ashworth Dirac<sup>105</sup>, Melissa Dirchwolf<sup>106</sup>, Jean-François Dufour<sup>107</sup>, Karel Dvorak<sup>108</sup>, Mattias Ekstedt<sup>109</sup>, Mohamed El-Kassas<sup>110</sup>, Osama M. Elsanousi<sup>111</sup>, Ahmed M. Elshar-kawy<sup>34</sup>, Reda M. Elwakil<sup>112</sup>, Wayne Eskridge<sup>113</sup>, Mohammed Eslam<sup>114</sup>, Gamal Esmat<sup>115</sup>, Jian- Gao Fan<sup>116</sup>, Maria Lucia Ferraz<sup>117</sup>, Robert Flisiak<sup>118</sup>, Davide Fortin<sup>10</sup>, Yasser Fouad<sup>119</sup>, Scott L. Friedman<sup>65</sup>, Michael Fuchs<sup>120</sup>, Adrian Gadano<sup>121</sup>, Amalia Gastaldelli<sup>122</sup>, Anja Geerts<sup>123</sup>, Andreas Geier<sup>124</sup>, Jacob George<sup>114</sup>, Lynn H. Gerber<sup>125</sup>, Hasmik L. Ghazinyan<sup>126</sup>, Liana Gheorghie<sup>127</sup>, Denise Giangola Kile<sup>128</sup>, Marcos Giral<sup>129</sup>, George Goh Boon Bee<sup>130</sup>, Nicolas Goossens<sup>131</sup>, Isabel Graupera<sup>132</sup>, Henning Grønbaek<sup>133</sup>, Saeed Hamid<sup>134</sup>, Vanessa Hebditch<sup>135</sup>, Zachary Henry<sup>79</sup>, Ingrid J. Hickman<sup>136</sup>, L. Ansley Hobbs<sup>3</sup>, Samantha L. Hocking<sup>137</sup>, Wolf Peter Hofmann<sup>138</sup>, Ramazan Idilman<sup>139</sup>, Paula Iruzubieta<sup>96</sup>, Scott Isaacs<sup>140</sup>, Vasily A. Isakov<sup>141</sup>, Mona H. Ismail<sup>142</sup>, Mohammad H. Jamal<sup>143</sup>, Helen Jarvis<sup>144</sup>, Peter Jepsen<sup>133</sup>, François R. Jormayvaz<sup>145</sup>, Sudhamshu K.C.<sup>146</sup>, Satoru Kakizaki<sup>147</sup>, Saul Karpen<sup>148</sup>, Takumi Kawaguchi<sup>149</sup>, Shelley E. Keating<sup>150</sup>, Yousef Khader<sup>151</sup>, Seung Up Kim<sup>152</sup>, Won Kim<sup>153</sup>, David E. Kleiner<sup>154</sup>, Ger Koek<sup>155</sup>, Narcisse Patrice Joseph Komar<sup>156</sup>, Loretta A. Kondili<sup>157</sup>, Bart G. Koot<sup>158</sup>, Marko Korenjak<sup>159</sup>, Eleni Kotsiliti<sup>160</sup>, Yiannoula Koulla<sup>161</sup>, Carina Kugelmas<sup>162</sup>, Marcelo Kugelmas<sup>163</sup>, Asma Labidi<sup>164</sup>, Naomi F. Lange<sup>165</sup>, Joel E. Lavine<sup>166</sup>, Mariana Lazo<sup>167</sup>, Nathalie Leite<sup>168</sup>, Han-Chieh Lin<sup>169</sup>, Udrnam Lkhagva<sup>62</sup>, Michelle T. Long<sup>170</sup>, Patricio Lopez-Jaramillo<sup>171</sup>, Adelina Lozano<sup>172</sup>, Maria Paula Macedo<sup>173</sup>, Reza Malekzadeh<sup>174</sup>, Giulio Marchesini<sup>175</sup>, Sebastian Marciano<sup>121</sup>, Kim Martinez<sup>176</sup>, Sophia E. Martínez Vázquez<sup>20</sup>, Lyudmila Mateva<sup>177</sup>, José M. Mato<sup>178</sup>, Charles N. Mbendi<sup>179</sup>, Alexis Gorden McCarty<sup>180</sup>, Jeff McIntyre<sup>181</sup>, Martin McKee<sup>182</sup>, Juan M. Mendive<sup>183</sup>, Ivana Mikolasevic<sup>184</sup>, Pamela S. Miller<sup>185</sup>, Tamara Milovanovic<sup>186</sup>, Terri Milton<sup>187</sup>, Rosalba Moreno-Alcantar<sup>188</sup>, Timothy R. Morgan<sup>189</sup>, Ayesha A. Motala<sup>190</sup>, Jean Muris<sup>191</sup>, Carla Musso<sup>192</sup>, Edna J. Nava-González<sup>193</sup>, Francesco Negro<sup>194</sup>, Alexander V. Nerse-sov<sup>195</sup>, Brent A. Neuschwander-Tetri<sup>196</sup>, Dafina Nikolova<sup>197</sup>, Suzanne Norris<sup>198</sup>, Katja Novak<sup>199</sup>, Ponsiano Ocama<sup>200</sup>, Janus P. Ong<sup>201</sup>, Arlinking Ong-Go<sup>202</sup>, Charles Onyekwere<sup>203</sup>, P. Martin Padilla-Machaca<sup>204</sup>, Raluca Pais<sup>205</sup>, Calvin Q. Pan<sup>206</sup>, Arturo Panduro<sup>207</sup>, Manas K. Panigrahi<sup>208</sup>, Georgios Papatheodoridis<sup>209</sup>, Imran Paruk<sup>190</sup>, Keyur Patel<sup>210</sup>, Carlos Penha-Goncalves<sup>211</sup>, Norma M. Pérez<sup>212</sup>, Juanita Pérez-Escobar<sup>213</sup>, Juan M. Pericás<sup>214</sup>, Gianluca Perseghin<sup>215</sup>, Mário Guimarães Pessoa<sup>216</sup>, Salvatore Petta<sup>217</sup>, Claudia Pinto Marques Souza de Oliveira<sup>218</sup>, Dorairaj Prabhakaran<sup>219</sup>, Nikolaos Prysopoulos<sup>220</sup>, Attoosa Rabiee<sup>221</sup>, Alnoor Ramji<sup>222</sup>, Vlad Ratziu<sup>223</sup>, Natarajan Ravendran<sup>224</sup>, Katrina Ray<sup>225</sup>, Michael Roden<sup>226</sup>, Stefano Romeo<sup>227</sup>, Manuel Romero-Gómez<sup>228</sup>, Yaron Rotman<sup>229</sup>, Samir Rouabhia<sup>230</sup>, Ian A. Rowe<sup>231</sup>, Shakhlo Sadirova<sup>232</sup>, Maryam Salem Alkhatry<sup>233</sup>, Riina Salupere<sup>234</sup>, Sanjaya K. Satapathy<sup>235</sup>, Jeffrey B. Schwimmer<sup>236</sup>, Giada Sebastiani<sup>237</sup>, Lynn Seim<sup>238</sup>, Yosuke Seki<sup>239</sup>, Abdel Karim Serme<sup>240</sup>, David Shapiro<sup>241</sup>, Lali Sharvadze<sup>242</sup>, Jonathan E. Shaw<sup>243</sup>, Isaac Thom Shawa<sup>244</sup>, Thirivikrama Shenoy<sup>245</sup>, Oren Shibolet<sup>246</sup>, Yusuke Shimakawa<sup>247</sup>, Jay H. Shubrook<sup>248</sup>, Shivaram Prasad Singh<sup>249</sup>, Edford Sinkala<sup>250</sup>, Lubomir Skladany<sup>251</sup>, Igor Skrypnik<sup>252</sup>, Myeong Jun Song<sup>253</sup>, Silvia Sookoian<sup>254</sup>, Kannan Sridharan<sup>255</sup>, Norbert Stefan<sup>256</sup>, Jonathan G. Stine<sup>257</sup>, Nikos Stratakis<sup>1</sup>, Dhastagir Sultan Sheriff<sup>258</sup>, Shikha S. Sundaram<sup>259</sup>, Gianluca Svegliati-Baroni<sup>260</sup>, Mark G. Swain<sup>261</sup>, Frank Tacke<sup>101</sup>, Shahrad Taheri<sup>262</sup>, Soek-Siam Tan<sup>263</sup>, Elliot B. Tapper<sup>264</sup>, Giovanni Targher<sup>265</sup>, Eugen Tcaciuc<sup>266</sup>, Maja Thiele<sup>32</sup>, Dina Tinia-cos<sup>267</sup>, Ieva Tolmane<sup>268</sup>, Aldo Torre<sup>269</sup>, Esther A. Torres<sup>270</sup>, Sombat Treeprasert-suk<sup>271</sup>, Michael Trenell<sup>272</sup>, Svetlana Turcan<sup>266</sup>, Adela Turcanu<sup>266</sup>, Jonas Valantinas<sup>273</sup>, Laurens A. van Kleef<sup>99</sup>, Jose Antonio Velarde Ruiz Velasco<sup>274</sup>, Mette Vesterhus<sup>275</sup>, Eduardo Vilar-Gomez<sup>276</sup>, Imam Waked<sup>277</sup>, Julia



- Wattacheril<sup>278</sup>, Heiner Wedemeyer<sup>279</sup>, Fonda Wilkins<sup>280</sup>, José Willems<sup>281</sup>, Robert J. Wong<sup>282</sup>, Yusuf Yilmaz<sup>283</sup>, Hannele Yki-Järvinen<sup>284</sup>, Ming-Lung Yu<sup>285</sup>, Volkan Yumuk<sup>286</sup>, Müjdat Zeybel<sup>287</sup>, Kenneth I. Zheng<sup>288</sup>, Ming-Hua Zheng<sup>288</sup>,  
\*Contributed equally.
- <sup>1</sup>Barcelona Institute for Global Health (ISGlobal), Hospital Clínic, University of Barcelona, Barcelona, Spain
- <sup>2</sup>Faculty of Medicine and Health Sciences, University of Barcelona, Barcelona, Spain
- <sup>3</sup>CUNY Graduate School of Public Health and Health Policy (CUNY SPH), New York, NY, USA
- <sup>4</sup>European Association for the Study of the Liver (EASL), Geneva, Switzerland
- <sup>5</sup>Independent consultant, Nottingham, UK
- <sup>6</sup>Division of Gastroenterology and Hepatology, Department of Medicine, Mayo Clinic, Rochester, MN, USA
- <sup>7</sup>Division of Gastroenterology, Department of Medicine, Schulich School of Medicine, Western University & London Health Sciences Centre, London, Ontario, Canada
- <sup>8</sup>Department of Epidemiology and Biostatistics, Schulich School of Medicine, Western University, London, Ontario, Canada
- <sup>9</sup>Department of Gastroenterology, School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile
- <sup>10</sup>Aix Marseille Univ, Inserm, IRD, SESSTIM, Sciences Economiques & Sociales de la Santé & Traitement de l'Information Médicale, ISSPAM, Marseille, France
- <sup>11</sup>Houston Methodist Hospital, Houston Research Institute, Houston, TX, USA
- <sup>12</sup>Barts Liver Centre, Blizard Institute, Queen Mary University of London, London, UK
- <sup>13</sup>Fatty Liver Program, Arizona Liver Health, Phoenix, AZ, USA
- <sup>14</sup>King Faisal Specialist Hospital & Research Centre, Riyadh, Saudi Arabia
- <sup>15</sup>Liver Unit, Hospital Clínic, University of Barcelona, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain
- <sup>16</sup>Division of Hepatology, Department of Medicine II, Leipzig University Medical Center, Leipzig, Germany
- <sup>17</sup>Division of Hepatology, University of Dundee, Dundee, Scotland, UK
- <sup>18</sup>Multivisceral Transplant Unit-Gastroenterology, Department of Surgery, Oncology and Gastroenterology at the Padua University Hospital, Padua, Italy
- <sup>19</sup>Department of Hepatology and Transplant, Hospital Médica Sur, Mexico City, Mexico
- <sup>20</sup>Department of Gastroenterology, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico
- <sup>21</sup>Asociación Latinoamericana para el Estudio del Hígado (ALEH), Santiago, Chile
- <sup>22</sup>Clinica Universitária de Gastrenterologia, Laboratório de Nutrição, Faculdade de Medicina, Universidade de Lisboa, Lisbon, Portugal
- <sup>23</sup>Division of Endocrinology, Diabetes & Metabolism, Department of Medicine, University of Florida, Gainesville, FL, USA
- <sup>24</sup>Greek Patients Association, Athens, Greece
- <sup>25</sup>Department of Hepatology, Postgraduate Institute of Medical Education and Research, Chandigarh, India
- <sup>26</sup>Department of Gastroenterology and Hepatology, Antwerp University Hospital, Edegem, Belgium
- <sup>27</sup>InflaMed Centre of Excellence, Laboratory for Experimental Medicine and Paediatrics, Translational Sciences in Inflammation and Immunology, Faculty of Medicine and Health Sciences, University of Antwerp, Wilrijk, Belgium
- <sup>28</sup>Department of Medicine, Huddinge, Karolinska Institutet, Stockholm, Sweden
- <sup>29</sup>CUNY Center for Systems and Community Design and NYU-CUNY Prevention Research Center, New York, NY, USA
- <sup>30</sup>Kautz 5 uGÜ, Köln, Germany
- <sup>31</sup>Independent researcher, Ponte de Lima, Portugal
- <sup>32</sup>Department of Gastroenterology and Hepatology, Odense University Hospital, Odense, Denmark
- <sup>33</sup>University California Berkeley School of Public Health, Berkeley, CA, USA
- <sup>34</sup>National Institute for Health Research Birmingham Biomedical Research Centre at University Hospitals Birmingham NHS Foundation Trust and the University of Birmingham, Birmingham, UK
- <sup>35</sup>Department of Medicine, University of Chicago, Chicago, IL, USA
- <sup>36</sup>Department of Community Health and Social Sciences, CUNY Graduate School of Public Health and Health Policy, New York, NY, USA
- <sup>37</sup>Department of Hepatology, Institute of Liver and Biliary Sciences, New Delhi, India
- <sup>38</sup>Hepatology and Clinical Research Units, Hospital Universitario Austral, Buenos Aires, Argentina
- <sup>39</sup>Division of Hepatology, Department of Medicine, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa
- <sup>40</sup>UCL Institute for Liver and Digestive Health, Royal Free Hospital, London, UK
- <sup>41</sup>Precision Medicine, Biological Resource Center, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy
- <sup>42</sup>Department of Pathophysiology and Transplantation, University of Milan, Milan, Italy
- <sup>43</sup>School of Public Health, Faculty of Social Welfare and Health Sciences, University of Haifa, Haifa, Israel
- <sup>44</sup>Department of Gastroenterology, Tel Aviv Medical Centre, Tel Aviv, Israel
- <sup>45</sup>Metabolic Liver Research Program, I. Department of Medicine, University Medical Centre Mainz, Mainz, Germany
- <sup>46</sup>The Chinese University of Hong Kong, Hong Kong, China
- <sup>47</sup>Center for Liver Disease, Inova, Falls Church, VA, USA
- <sup>48</sup>Transplantation and Liver Surgery, Helsinki University Hospital, Helsinki, Finland
- <sup>49</sup>Medical School, The University of Western Australia, Perth, Western Australia, Australia
- <sup>50</sup>Department of Internal Medicine, Division of Gastroenterology and Hepatology, Armed Forces Hospital, Muscat, Oman
- <sup>51</sup>Gastroenterology, Hepatology & Infectious Diseases Department, Benha University, Benha, Egypt
- <sup>52</sup>Republican Clinical Hospital "Timofei Mosneaga", Chişinău, Republic of Moldova
- <sup>53</sup>Liver Unit, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK
- <sup>54</sup>Department of Primary Care & Public Health, Faculty of Medicine, Imperial College, London, UK
- <sup>55</sup>Liver Disease Research Center, Department of Medicine, College of Medicine, King Saud University, Riyadh, Saudi Arabia
- <sup>56</sup>School of Medicine, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil
- <sup>57</sup>Department of Translational and Precision Medicine, Sapienza University of Rome, Rome, Italy
- <sup>58</sup>Department of Biomedicine, Biotechnology and Public Health, University of Cadiz, Cadiz, Spain
- <sup>59</sup>Servicio de Aparato Digestivo, Hospital Universitario Virgen de la Victoria, Instituto de Investigación Biomédica de Málaga y Plataforma en Nanomedicina-IBIMA Plataforma BIONAND, Universidad de Málaga, Málaga, Spain
- <sup>60</sup>Translational & Clinical Research Institute, Faculty of Medical Science, Newcastle University, Newcastle upon Tyne, UK
- <sup>61</sup>University of Health and Allied Science, Ho, Ghana
- <sup>62</sup>Mongolian National University of Medical Sciences, Ulaanbaatar, Mongolia
- <sup>63</sup>VA Boston Healthcare System and Harvard Medical School, Boston, MA, USA
- <sup>64</sup>Scientific Department, The Research Institute of Virology of the MoH of the Republic of Uzbekistan, Tashkent, Uzbekistan
- <sup>65</sup>Icahn School of Medicine at Mount Sinai, New York, NY, USA
- <sup>66</sup>Université Paris Cité, Inserm T<sup>3</sup>S, Paris, France
- <sup>67</sup>Centre for Obesity Research, Department of Medicine, University College London, London, UK
- <sup>68</sup>Pacific Rim Pathology Group, San Diego, CA, USA
- <sup>69</sup>Internal Medicine Department, Yale University, New Haven, CT, USA
- <sup>70</sup>Department of Visceral Surgery and Medicine, Bern University Hospital, University of Bern, Bern, Switzerland
- <sup>71</sup>Fatty Liver Alliance, Toronto, Ontario, Canada
- <sup>72</sup>Precision Medicine Lab, Biological Resource Center, and Transfusion Medicine, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy
- <sup>73</sup>IRCCS Ospedale San Raffaele, Milan, Italy
- <sup>74</sup>Angers University & Angers University Hospital, Angers, France
- <sup>75</sup>Dept. of Pathology and Immunology, Washington University School of Medicine, St. Louis, MO, USA
- <sup>76</sup>Dept. of Medical Sciences, University of Torino, Torino, Italy
- <sup>77</sup>School of Medicine, University of Dundee, Dundee, Scotland, UK
- <sup>78</sup>Liver Unit, Hospital G. Almenara, Universidad Mayor de San Marcos, Lima, Peru
- <sup>79</sup>Department of Medicine, University of Virginia, Charlottesville, VA, USA
- <sup>80</sup>Division of Gastroenterology, Department of Medicine, University of Washington, Seattle, WA, USA
- <sup>81</sup>Institute of Gastroenterology, La Havana, Cuba
- <sup>82</sup>Université Paris Cité, Department of Hepatology, Hospital Beaujon, AP-HP, Clichy, Paris, France
- <sup>83</sup>Departamento Unidad Metabólica, Hospital Universitario Fundación Favaloro, Buenos Aires, Argentina
- <sup>84</sup>Endocrinology Diabetes Nutrition Hospices Civils de Lyon, Lyon, France



- <sup>85</sup>Central Military Hospital, Asociación Latinoamericana para el Estudio del Hígado, Mexico City, Mexico
- <sup>86</sup>IRCCS MultiMedica, Milan, Italy
- <sup>87</sup>Gastroenterology and Hepatology Unit, Department of Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia
- <sup>88</sup>Center for Cohort Studies, Total Healthcare Center, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, Seoul, South Korea
- <sup>89</sup>Faculty of Medicine Siriraj Hospital, Bangkok, Thailand
- <sup>90</sup>Médica Sur Clinic & Foundation, Mexico City, Mexico
- <sup>91</sup>Liver Center, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA
- <sup>92</sup>European Association for the Study of the Liver (EASL) International Liver Foundation, Geneva, Switzerland
- <sup>93</sup>Department of Medicine, University of Otago, Wellington, New Zealand
- <sup>94</sup>School of Medicine, Federal University of Bahia, Salvador, Brazil
- <sup>95</sup>School of Medicine, University of Palermo, Palermo, Italy
- <sup>96</sup>Gastroenterology and Hepatology Department, Clinical and Translational Research in Digestive Diseases, Valdecilla Research Institute (DIVAL), Marqués de Valdecilla University Hospital, Santander, Spain
- <sup>97</sup>Faculty of Medicine, University of Kelaniya, Kelaniya, Sri Lanka
- <sup>98</sup>Washington University School of Medicine, St. Louis, MO 63110, USA
- <sup>99</sup>Erasmus MC University Medical Center, Rotterdam, the Netherlands
- <sup>100</sup>CHU Bordeaux, Bordeaux, France
- <sup>101</sup>Department of Hepatology & Gastroenterology, Charité - Universitätsmedizin Berlin, Berlin, Germany
- <sup>102</sup>St. Paul's Hospital Millennium Medical College, Addis Ababa, Ethiopia
- <sup>103</sup>Digestive Diseases Department, Hospital General Universitario de Valencia, Valencia, Spain
- <sup>104</sup>Banner Liver Support Group, Phoenix, AZ, USA
- <sup>105</sup>Department of Family Medicine, University of Washington, Seattle, WA, USA
- <sup>106</sup>Liver Unit, Hospital Privado de Rosario, Rosario, Argentina
- <sup>107</sup>Centre des Maladies Digestives Lausanne, Lausanne, Switzerland
- <sup>108</sup>Fourth Department of Internal Medicine, First Faculty of Medicine, Charles University, Prague, Czech Republic
- <sup>109</sup>Department of Health, Medicine and Caring Sciences, Linköping University, Linköping, Sweden
- <sup>110</sup>Endemic Medicine Department, Faculty of Medicine, Helwan University, Cairo, Egypt
- <sup>111</sup>Department of Surgery, Faculty of Medicine, National Ribat University, Khar-toum, Sudan
- <sup>112</sup>Tropical Medicine Department, Ain Shams University, Cairo, Egypt
- <sup>113</sup>Fatty Liver Foundation, Boise, ID, USA
- <sup>114</sup>Storr Liver Centre, Westmead Hospital, Westmead Institute for Medical Research, University of Sydney, Sydney, New South Wales, Australia
- <sup>115</sup>Endemic Medicine Department, Cairo University, Cairo, Egypt
- <sup>116</sup>Department of Gastroenterology, Xinhua Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai Key Lab of Pediatric Gastroenterology and Nutrition, Shanghai, China
- <sup>117</sup>Federal University of Sao Paulo, São Paulo, Brazil
- <sup>118</sup>Department of Infectious Diseases and Hepatology, Medical University of Bialystok, Bialystok, Poland
- <sup>119</sup>Faculty of Medicine, Minia University, Minya, Egypt
- <sup>120</sup>Central Virginia VA Health Care System and Virginia Commonwealth University, Richmond, VA, USA
- <sup>121</sup>Liver Unit, Hospital Italiano de Buenos Aires, Buenos Aires, Argentina
- <sup>122</sup>Institute of Clinical Physiology, National Research Council, Pisa, Italy
- <sup>123</sup>Department of Gastroenterology & Hepatology, University Hospital Ghent, Ghent, Belgium
- <sup>124</sup>Division of Hepatology, University Hospital Wuerzburg, Wuerzburg, Germany
- <sup>125</sup>Beatty Liver and Obesity Center, Inova Health System, Falls Church, VA, USA
- <sup>126</sup>Nikoméd Medical Centre, Yerevan, Armenia
- <sup>127</sup>Carol Davila University of Medicine and Pharmacy, Bucharest, Romania
- <sup>128</sup>Independent participant, Myrtle Beach, SC, USA
- <sup>129</sup>Departamento de Gastroenterología, Hospital de Clínicas, Universidad Nacional de Asunción, San Lorenzo, Paraguay
- <sup>130</sup>Department of Gastroenterology and Hepatology, Singapore General Hospital, Singapore
- <sup>131</sup>Division of Gastroenterology and Hepatology, Geneva University Hospital, Geneva, Switzerland
- <sup>132</sup>Liver Unit, Hospital Clínic, FCRB-IDIBAPS, University of Barcelona, Barcelona, Spain
- <sup>133</sup>Department of Hepatology and Gastroenterology, Aarhus University Hospital, Aarhus, Denmark
- <sup>134</sup>Department of Medicine, Aga Khan University, Karachi, Pakistan
- <sup>135</sup>British Liver Trust, Winchester, UK
- <sup>136</sup>Department of Nutrition and Dietetics, Princess Alexandra Hospital, Brisbane, Queensland, Australia
- <sup>137</sup>Central Clinical School, Faculty of Medicine and Health, University of Sydney, Sydney, New South Wales, Australia
- <sup>138</sup>Medical Care Center for Gastroenterology Bayerischer Platz, Berlin, Germany
- <sup>139</sup>Department of Gastroenterology, Ankara University School of Medicine, Ankara, Turkey
- <sup>140</sup>Department of Medicine, Emory University School of Medicine, Atlanta, GA, USA
- <sup>141</sup>Department of Gastroenterology & Hepatology, Federal Research Centre of Nutrition, Biotechnology & Food Safety, Moscow, Russia
- <sup>142</sup>King Fahd Hospital of the University, Al-Khobar, and College of Medicine, Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia
- <sup>143</sup>Department of Transplantation and Surgery, Kuwait University, Kuwait City, Kuwait
- <sup>144</sup>Newcastle University, Newcastle, UK
- <sup>145</sup>Geneva University Hospital, Geneva, Switzerland
- <sup>146</sup>National Academy of Medical Sciences, Kathmandu, Nepal
- <sup>147</sup>Department of Clinical Research, National Hospital Organization Takasaki General Medical Center, Takasaki, Japan
- <sup>148</sup>Emory University School of Medicine, Atlanta, GA, USA
- <sup>149</sup>Division of Gastroenterology, Department of Medicine, Kurume University School of Medicine, Kurume, Japan
- <sup>150</sup>School of Human Movement and Nutrition Sciences, The University of Queensland, Brisbane, Queensland, Australia
- <sup>151</sup>Department of Public Health, Jordan University of Science and Technology, Irbid, Jordan
- <sup>152</sup>Department of Internal Medicine, Yonsei University College of Medicine, Seoul, Republic of Korea
- <sup>153</sup>Department of Internal Medicine, Seoul National University College of Medicine, Seoul Metropolitan Government Boramae Medical Center, Seoul, Republic of Korea
- <sup>154</sup>Laboratory of Pathology, Center for Cancer Research, National Cancer Institute, NIH, Bethesda, MD, USA
- <sup>155</sup>Maastricht University Medical Center, Maastricht, the Netherlands
- <sup>156</sup>Institut Pasteur de Bangui, Bangui, Central African Republic
- <sup>157</sup>Center for Global Health, Istituto Superiore Di Sanità (ISS), UniCamillus-Saint Camillus International University of Health Sciences, Rome, Italy
- <sup>158</sup>Department of Pediatric Gastroenterology, Emma's Children Hospital, Amsterdam University Medical Center, Amsterdam, the Netherlands
- <sup>159</sup>European Liver Patients' Association, Brussels, Belgium
- <sup>160</sup>Nature Reviews Gastroenterology & Hepatology, Berlin, Germany
- <sup>161</sup>Cyprus Liver Patients Association, Nicosia, Cyprus
- <sup>162</sup>Department of Pediatrics, Denver Health Medical Center, Denver, CO, USA
- <sup>163</sup>South Denver Gastroenterology, Englewood, CO, USA
- <sup>164</sup>Gastroenterology "A" Department, Rabta University Hospital, Faculty of Medicine of Tunis, University of Tunis El Manar, Tunis, Tunisia
- <sup>165</sup>Department of Visceral Surgery and Medicine, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland
- <sup>166</sup>Department of Pediatrics, Columbia University, New York, NY, USA
- <sup>167</sup>Urban Health Collaborative, Dornsife School of Public Health, Drexel University, Philadelphia, PA, USA
- <sup>168</sup>Department of Internal Medicine, University Hospital Clementino Fraga Filho, School of Medicine, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil
- <sup>169</sup>Division of Gastroenterology and Hepatology, Taipei Veterans General Hospital, Taipei, Taiwan
- <sup>170</sup>Novo Nordisk A/S, Vandtårnsvej 108-110, 2860 Søborg, Denmark
- <sup>171</sup>Masira Research Institute, Medical School, Universidad de Santander (JDES), Bucaramanga, Colombia
- <sup>172</sup>Cayetano Heredia Peruvian University, Lima, Peru
- <sup>173</sup>NOVA4Health, NOVA Medical School, Faculdade de Ciências Médicas, NMS, FCM, Universidade NOVA de Lisboa, Lisbon, Portugal
- <sup>174</sup>Digestive Disease Research Institute, Tehran University of Medical Sciences, Tehran, Iran
- <sup>175</sup>Alma Mater-University of Bologna, Bologna, Italy
- <sup>176</sup>GLI, Lakewood, NJ, USA
- <sup>177</sup>University Hospital "St Ivan Rilski", Medical University of Sofia, Sofia, Bulgaria
- <sup>178</sup>CIC bioGUNE, Technology Park of Bizkaia, Derio, Spain
- <sup>179</sup>Service of Hepatology and Gastroenterology, Department of Internal Medicine, University Clinics of Kinshasa, Kinshasa, Democratic Republic of the Congo
- <sup>180</sup>Mid-Atlantic Permanente Medical Group, Rockville, MD, USA

- <sup>181</sup>Liver Health Programs, Global Liver Institute, Washington, DC, USA
- <sup>182</sup>London School of Hygiene & Tropical Medicine, London, UK
- <sup>183</sup>La Mina Primary Health Care Academic Centre, Catalan Health Institute, University of Barcelona, Barcelona, Spain
- <sup>184</sup>Department of Gastroenterology, UHC Rijeka, Rijeka, Croatia
- <sup>185</sup>Independent participant, Powell, OH, USA
- <sup>186</sup>School of Medicine, University of Belgrade, University Clinical Center of Serbia, Belgrade, Serbia
- <sup>187</sup>Independent consultant, Houston, TX, USA
- <sup>188</sup>Gastroenterology Department, HE CMN SXXI, IMSS, Mexico City, Mexico
- <sup>189</sup>Medical Service, VA Long Beach Healthcare System, Long Beach, CA, USA
- <sup>190</sup>University of KwaZulu-Natal, Durban, South Africa
- <sup>191</sup>Dept. General Practice, Care and Public Health Research Institute (CAPHRI), Maastricht University, Maastricht, the Netherlands
- <sup>192</sup>Diabetes Metabolic Department, Hospital Universitario Fundación Favaloro, Buenos Aires, Argentina
- <sup>193</sup>Facultad de Salud Pública y Nutrición, Universidad Autónoma de Nuevo León, Monterrey, Mexico
- <sup>194</sup>University of Geneva, Geneva, Switzerland
- <sup>195</sup>Department of Gastroenterology, SD Asfendiyarov Kazakh National Medical University, Almaty, Kazakhstan
- <sup>196</sup>Saint Louis University, St. Louis, MO, USA
- <sup>197</sup>University Clinic for Gastroenterohepatology, University Ss. Cyril and Methodius, Skopje, Macedonia
- <sup>198</sup>Department of Hepatology, St James's Hospital, Dublin, Ireland
- <sup>199</sup>Dept. of Gastroenterology and Hepatology, University Medical Center Ljubljana, Ljubljana, Slovenia
- <sup>200</sup>Makerere University College of Health Sciences, Kampala, Uganda
- <sup>201</sup>University of the Philippines Manila, Manila, Philippines
- <sup>202</sup>Section of Gastroenterology and Hepatology, University of Santo Tomas Faculty of Medicine and Surgery, Manila, Philippines
- <sup>203</sup>Department of Internal Medicine, Lagos State University College of Medicine Ikeja, Lagos, Nigeria
- <sup>204</sup>Liver Unit, Guillermo Almenara National Hospital, National University of San Marcos, Lima, Peru
- <sup>205</sup>Sorbonne Université, Assistance Publique-Hôpitaux de Paris, Hôpital Pitié-Salpêtrière, Institute of Cardiometabolism and Nutrition, Centre de Recherche Saint Antoine, INSERM UMRS\_938 Paris, France
- <sup>206</sup>Division of Gastroenterology and Hepatology, NYU Langone Health, NYU Grossman School of Medicine, New York, NY, USA
- <sup>207</sup>Genomic Medicine in Hepatology, Hospital Civil de Guadalajara/CUCS, UdeG, Guadalajara, Mexico
- <sup>208</sup>All India Institute of Medical Sciences, Bhubaneswar, India
- <sup>209</sup>Medical School of National and Kapodistrian University of Athens, Athens, Greece
- <sup>210</sup>University Health Network, Toronto, Ontario, Canada
- <sup>211</sup>Instituto Gulbenkian de Ciência, Oeiras, Portugal
- <sup>212</sup>Gastroenterología-Hepatólogía-Trasplante Hepático, Hospital General de la Plaza de la Salud, Santo Domingo, Dominican Republic
- <sup>213</sup>Gastroenterology Department, Hospital Juárez de México, Mexico City, Mexico
- <sup>214</sup>Liver Unit, Vall d'Hebron University Hospital, Vall d'Hebron Institute for Research, Centros de Investigación Biomédica en Red Enfermedades Hepáticas y Digestivas (CIBERehd), Barcelona, Spain
- <sup>215</sup>Department of Medicine and Surgery, Università degli Studi di Milano-Bicocca, Milan, Italy
- <sup>216</sup>Division of Gastroenterology and Hepatology, University of São Paulo School of Medicine, São Paulo, Brazil
- <sup>217</sup>Section of Gastroenterology and Hepatology, PROMISE, University of Palermo, Palermo, Italy
- <sup>218</sup>Department of Gastroenterology, University of São Paulo School of Medicine, São Paulo, Brazil
- <sup>219</sup>Centre for Chronic Disease Control, New Delhi, India
- <sup>220</sup>Rutgers New Jersey Medical School, Newark, NJ, USA
- <sup>221</sup>Washington DC VA Medical Center, Washington DC, USA
- <sup>222</sup>University of British Columbia, Vancouver, British Columbia, Canada
- <sup>223</sup>Sorbonne Université, Paris, France
- <sup>224</sup>Department of Hepatology, Johns Hopkins School of Medicine, Baltimore, MD, USA
- <sup>225</sup>Nature Reviews Gastroenterology & Hepatology, London, UK
- <sup>226</sup>Department of Endocrinology and Diabetology, Medical Faculty, Heinrich Heine University, University Hospital, Düsseldorf, Germany
- <sup>227</sup>Department of Molecular and Clinical Medicine, Gothenburg University, Cardiology Department, Sahlgrenska University Hospital, Gothenburg, Sweden
- <sup>228</sup>Digestive Diseases Department and CIBERehd, Virgen del Rocio University Hospital, Institute of Biomedicine of Seville (HUVR/CSIC/US), University of Seville, Seville, Spain
- <sup>229</sup>Liver & Energy Metabolism Section, Liver Diseases Branch, National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), National Institutes of Health (NIH), Bethesda, MD, USA
- <sup>230</sup>Internal Medicine Department, Touhami Benflis University Hospital Centre, Batna, Algeria
- <sup>231</sup>University of Leeds, Leeds, UK
- <sup>232</sup>The Research Institute of Virology of the MoH of the Republic of Uzbekistan, Tashkent, Uzbekistan
- <sup>233</sup>Ibrahim bin Hamad Obaidullah Hospital, Emirates Health Services, RAK, UAE
- <sup>234</sup>Tartu University Hospital, University of Tartu, Tartu, Estonia
- <sup>235</sup>North Shore University Hospital, Zucker School of Medicine at Hofstra/Northwell Health, Hempstead, NY, USA
- <sup>236</sup>Division of Gastroenterology, Hepatology, and Nutrition, Department of Pediatrics, University of California San Diego School of Medicine, La Jolla, CA, USA
- <sup>237</sup>Division of Gastroenterology and Hepatology, Department of Medicine, McGill University Health Centre, Montreal, Quebec, Canada
- <sup>238</sup>American Liver Foundation, West Orange, NJ, USA
- <sup>239</sup>Weight Loss and Metabolic Surgery Center, Yotsuya Medical Cube, Tokyo, Japan
- <sup>240</sup>University Joseph KI-ZERBO, Ouagadougou, Burkina Faso
- <sup>241</sup>Integrated Quality Resources, San Diego, CA, USA
- <sup>242</sup>Infectious Diseases, AIDS and Clinical Immunology Research Center, Tbilisi, Georgia
- <sup>243</sup>Baker Heart and Diabetes Institute, Melbourne, Victoria, Australia
- <sup>244</sup>University of Derby, Derby, UK
- <sup>245</sup>Sree Gokulam Medical College and Research Foundation, Venjar-ammoodu, India
- <sup>246</sup>Department of Gastroenterology & Hepatology, Tel Aviv Medical Center & Tel Aviv University, Tel Aviv, Israel
- <sup>247</sup>Institut Pasteur, Université Paris Cité, Unité d'Épidémiologie des Maladies Émergentes, Paris, France
- <sup>248</sup>Touro University California, Vallejo, CA, USA
- <sup>249</sup>Kalinga Gastroenterology Foundation, Cuttack, India
- <sup>250</sup>The University of Zambia, School of Medicine, Department of Internal Medicine, Lusaka, Zambia
- <sup>251</sup>HEGITO Liver & Transplant Unit, Dept. Internal Medicine of the Slovak Medical University, F.D. Roosevelt Teaching Hospital, Banská Bystrica, Slovakia
- <sup>252</sup>Internal Medicine №1 Department, Poltava State Medical University, Poltava, Ukraine
- <sup>253</sup>Department of Internal Medicine, College of Medicine, The Catholic University of Korea, Seoul, Republic of Korea
- <sup>254</sup>Clinical and Molecular Hepatology, Centro de Altos Estudios en Ciencias Humanas y de la Salud (CAECIHS), Universidad Abierta Interamericana, Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Buenos Aires, Argentina
- <sup>255</sup>Arabian Gulf University, Manama, Bahrain
- <sup>256</sup>University Hospital of Tübingen, Tübingen, Germany
- <sup>257</sup>Penn State Health Milton S. Hershey Medical Center, Hershey, PA, USA
- <sup>258</sup>Anna Medical College, Montagne Blanche, Mauritius
- <sup>259</sup>Digestive Health Institute, Children's Hospital Colorado, University of Colorado School of Medicine, Aurora, CO, USA
- <sup>260</sup>Liver Injury and Transplant Unit, Polytechnic University of Marche, Ancona, Italy
- <sup>261</sup>Department of Medicine, University of Calgary, Calgary, Alberta, Canada
- <sup>262</sup>Hamad Medical Corporation, Doha, Qatar
- <sup>263</sup>Department of Hepatology, Selayang Hospital, Batu Caves, Malaysia
- <sup>264</sup>University of Michigan, Ann Arbor, MI, USA
- <sup>265</sup>Section of Diabetes and Endocrinology, University of Verona, Verona, Italy
- <sup>266</sup>Discipline of Gastroenterology, Nicolae Testemitanu State University of Medicine and Pharmacy, Chişinău, Republic of Moldova
- <sup>267</sup>Dept. of Pathology, Aretaieion Hospital, Medical School, National and Kapodistrian University of Athens, Athens, Greece
- <sup>268</sup>Riga East University Hospital, University of Latvia, Riga, Latvia
- <sup>269</sup>Metabolic Unit, Instituto Nacional de Ciencias Médicas y Nutrición "Salvador Zubirán", Mexico City, Mexico
- <sup>270</sup>Department of Medicine, University of Puerto Rico School of Medicine, San Juan, Puerto Rico

- <sup>271</sup>Chulalongkorn University, Bangkok, Thailand
- <sup>272</sup>Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK
- <sup>273</sup>Vilnius University Hospital Santaros Clinics, Vilnius, Lithuania
- <sup>274</sup>Hospital Civil de Guadalajara Fray Antonio Alcalde, Guadalajara, Mexico
- <sup>275</sup>Dept. of Clinical Science, University of Bergen, Dept. of Medicine, Haraldsplass Deaconess Hospital, Bergen, Norway
- <sup>276</sup>Indiana University School of Medicine, Indianapolis, IN, USA
- <sup>277</sup>National Liver Institute, Shebeen El-Kom, Egypt
- <sup>278</sup>Department of Medicine, Division of Digestive and Liver Diseases, Columbia University Irving Medical Center, Center for Liver Disease and Transplantation, New York, NY, USA
- <sup>279</sup>Department of Gastroenterology, Hepatology, Infectious Diseases and Endocrinology, Hannover Medical School, Hannover, Germany
- <sup>280</sup>Lexington Medical Center, West Columbia, SC, USA
- <sup>281</sup>Dutch Liver Patients Association, Hoogland, the Netherlands
- <sup>282</sup>Division of Gastroenterology and Hepatology, Stanford University School of Medicine, Veterans Affairs Palo Alto Healthcare System, Palo Alto, CA, USA
- <sup>283</sup>Department of Gastroenterology, School of Medicine, Recep Tayyip Erdoğan University, Rize, Turkey
- <sup>284</sup>University of Helsinki and Minerva Foundation Institute for Medical Research, Helsinki, Finland
- <sup>285</sup>Hepatobiliary Division, Department of Internal Medicine, Kaohsiung Medical University Hospital, Kaohsiung Medical University, School of Medicine, College of Medicine, National Sun Yet-sen University, Kaohsiung City, Taiwan
- <sup>286</sup>Istanbul University-Cerrahpaşa, Cerrahpaşa Medical Faculty, Division of Endocrinology, Metabolism and Diabetes, Istanbul, Turkey
- <sup>287</sup>Department of Gastroenterology and Hepatology, Koç University, Istanbul, Turkey
- <sup>288</sup>MAFLD Research Center, Department of Hepatology, the First Affiliated Hospital of Wenzhou Medical University, Wenzhou, China

## Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jhep.2023.04.035>.

## References

*Author names in bold designate shared co-first authorship.*

- [1] Younossi ZM, Golabi P, Paik JM, Henry A, Van Dongen C, Henry L. The global epidemiology of nonalcoholic fatty liver disease (NAFLD) and nonalcoholic steatohepatitis (NASH): a systematic review. *Hepatology* 2023;77(4):1135–1347.
- [2] Riazhi K, Azhari H, Charette JH, Underwood FE, King JA, Afshar EE, et al. The prevalence and incidence of NAFLD worldwide: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol* 2022;7(9):851–861.
- [3] Sweeny KF, Lee CK. Nonalcoholic Fatty Liver Disease in Children. *Gastroenterol Hepatol (N Y)* 2021;17(12):579–587.
- [4] Araújo AR, Rosso N, Bedogni G, Tiribelli C, Bellentani S. Global epidemiology of non-alcoholic fatty liver disease/non-alcoholic steatohepatitis: What we need in the future. *Liver Int* 2018;38(Suppl 1):47–51.
- [5] Kanwal F, Kramer JR, Mapakshi S, Natarajan Y, Chayanupatkul M, Richardson PA, et al. Risk of Hepatocellular Cancer in Patients With Non-Alcoholic Fatty Liver Disease. *Gastroenterology* 2018;155(6):1828–1837.e2.
- [6] McSweeney L, Breckons M, Fattakhova G, Oluboyede Y, Vale L, Ternent L, et al. Health-related quality of life and patient-reported outcome measures in NASH-related cirrhosis. *JHEP Rep* 2020;2(3):100099.
- [7] Sayiner M, Stepanova M, Pham H, Noor B, Walters M, Younossi ZM. Assessment of health utilities and quality of life in patients with non-alcoholic fatty liver disease. *BMJ Open Gastroenterol* 2016;3(1):e000106.
- [8] Younossi Z, Aggarwal P, Shrestha I, Fernandes J, Johansen P, Augusto M, et al. The burden of non-alcoholic steatohepatitis: A systematic review of health-related quality of life and patient-reported outcomes. *JHEP Rep* 2022;4(9):100525.
- [9] Global Burden of Disease Cancer Collaboration. Global, Regional, and National Cancer Incidence, Mortality, Years of Life Lost, Years Lived With Disability, and Disability-Adjusted Life-Years for 29 Cancer Groups, 1990 to 2017: A Systematic Analysis for the Global Burden of Disease Study. *JAMA Oncol* 2019;5(12):1749–1768.
- [10] Schattenberg JM, Lazarus JV, Newsome PN, Serfaty L, Aghemo A, Augustin S, et al. Disease burden and economic impact of diagnosed non-alcoholic steatohepatitis in five European countries in 2018: A cost-of-illness analysis. *Liver Int* 2021;41(6):1227–1242.
- [11] Younossi ZM, Blissett D, Blissett R, Henry L, Stepanova M, Younossi Y, et al. The economic and clinical burden of nonalcoholic fatty liver disease in the United States and Europe. *Hepatology* 2016;64(5):1577–1586.
- [12] Ludwig J, Viggiano TR, McGill DB, Oh BJ. Nonalcoholic steatohepatitis: Mayo Clinic experiences with a hitherto unnamed disease. *Mayo Clin Proc* 1980;55(7):434–438.
- [13] Anstee QM, Targher G, Day CP. Progression of NAFLD to diabetes mellitus, cardiovascular disease or cirrhosis. *Nat Rev Gastroenterol Hepatol* 2013;10(6):330–344.
- [14] Li AA, Kim D, Ahmed A. Association of sarcopenia and NAFLD: an overview. *Clin Liver Dis (Hoboken)* 2020;16(2):73–76.
- [15] Targher G, Tilg H, Byrne CD. Non-alcoholic fatty liver disease: a multisystem disease requiring a multidisciplinary and holistic approach. *Lancet Gastroenterol Hepatol* 2021;6(7):578–588.
- [16] Lazarus JV, Anstee QM, Hagström H, Cusi K, Cortez-Pinto H, Mark HE, et al. Defining comprehensive models of care for NAFLD. *Nat Rev Gastroenterol Hepatol* 2021;18(10):717–729.
- [17] Hussain A, Patel PJ, Rhodes F, Srivastava A, Patch D, Rosenberg W. Decompensated cirrhosis is the commonest presentation for NAFLD patients undergoing liver transplant assessment. *Clin Med (Lond)* 2020;20(3):313–318.
- [18] Newton JL. Systemic symptoms in non-alcoholic fatty liver disease. *Dig Dis* 2010;28(1):214–219.
- [19] Castera L. Non-invasive tests for liver fibrosis in NAFLD: creating pathways between primary healthcare and liver clinics. *Liver Int* 2020;40(Suppl 1):77–81.
- [20] Boursier J, Hagström H, Ekstedt M, Moreau C, Bonacci M, Cure S, et al. Non-invasive tests accurately stratify patients with NAFLD based on their risk of liver-related events. *J Hepatol* 2022;76(5):1013–1020.
- [21] Loomba R, Huang DQ, Sanyal AJ, Anstee QM, Trauner M, Lawitz EJ, et al. Liver stiffness thresholds to predict disease progression and clinical outcomes in bridging fibrosis and cirrhosis. *Gut* 2023;72(3):581–589.
- [22] Chalmers J, Wilkes E, Harris R, Kent L, Kinra S, Aithal GP, et al. The Development and Implementation of a Commissioned Pathway for the Identification and Stratification of Liver Disease in the Community. *Frontline Gastroenterol* 2020;11(2):86–92.
- [23] Moolla A, Motohashi K, Marjot T, Shard A, Ainsworth M, Gray A, et al. A multidisciplinary approach to the management of NAFLD is associated with improvement in markers of liver and cardio-metabolic health. *Frontline Gastroenterol* 2019;10(4):337–346.
- [24] Neilson LJ, Macdougall L, Lee PS, Hardy T, Beaton D, Chandrapalan S, et al. Implementation of a care bundle improves the management of patients with non-alcoholic fatty liver disease. *Frontline Gastroenterol* 2021;12(7):578–585.
- [25] Srivastava A, Gailer R, Tanwar S, Trembling P, Parkes J, Rodger A, et al. Prospective evaluation of a primary care referral pathway for patients with non-alcoholic fatty liver disease. *J Hepatol* 2019;71(2):371–378.
- [26] Dufour JF, Anstee QM, Bugianesi E, Harrison S, Loomba R, Paradis V, et al. Current therapies and new developments in NASH. *Gut* 2022;71(10):2123–2134.
- [27] Harrison SA, Allen AM, Dubourg J, Noureddin M, Alkhouri N. Challenges and opportunities in NASH drug development. *Nat Med* 2023;29(3):562–573.
- [28] Duseja A, Singh SP, De A, Madan K, Rao PN, Shukla A, et al. Indian National Association for Study of the Liver (INASL) Guidance Paper on Nomenclature, Diagnosis and Treatment of Nonalcoholic Fatty Liver Disease (NAFLD). *J Clin Exp Hepatol* 2023;13(2):273–302.
- [29] Romero-Gómez M, Zelber-Sagi S, Trenell M. Treatment of NAFLD with diet, physical activity and exercise. *J Hepatol* 2017;67(4):829–846.
- [30] Viveiros K. The Role of Life Style Modifications in Comprehensive Non-Alcoholic Fatty Liver Disease Treatment. *Clin Liver Dis (Hoboken)* 2021;17(1):11–14.
- [31] Díaz LA, Fuentes-López E, Ayares G, Idaisoaga F, Arnold J, Márquez-Lomas A, et al. The establishment of public health policies and the burden of non-alcoholic fatty liver disease in the Americas. *Lancet Gastroenterol Hepatol* 2022;7(6):552–559.
- [32] Lazarus JV, Colombo M, Cortez-Pinto H, Huang TTK, Miller V, Ninburg M, et al. NAFLD – sounding the alarm on a silent epidemic. *Nat Rev Gastroenterol Hepatol* 2020;17(7):377–379.
- [33] Lazarus JV, Mark HE, Colombo M, Demais S, Dillon JF, George J, et al. A sustainable development goal framework to guide multisectoral action on NAFLD through a societal approach. *Aliment Pharmacol Ther* 2022;55(2):234–243.
- [34] Lazarus JV, Mark HE, Anstee QM, Arab JP, Batterham RL, Castera L, et al. Advancing the global public health agenda for NAFLD: a consensus statement. *Nat Rev Gastroenterol Hepatol* 2022;19(1):60–78.



- [35] Jafri A, Mathe N, Aglago EK, Konyole SO, Ouedraogo M, Audain K, et al. Food availability, accessibility and dietary practices during the COVID-19 pandemic: a multi-country survey. *Public Health Nutr* 2021;24(7):1798–1805.
- [36] Mattioli AV, Sciomer S, Cocchi C, Maffei S, Gallina S. Quarantine during COVID-19 outbreak: Changes in diet and physical activity increase the risk of cardiovascular disease. *Nutr Metab Cardiovasc Dis* 2020;30(9):1409–1417.
- [37] Grebely J, Bruneau J, Lazarus JV, Dalgard O, Bruggmann P, Treloar C, et al. Research priorities to achieve universal access to hepatitis C prevention, management and direct-acting antiviral treatment among people who inject drugs. *Int J Drug Pol* 2017;47:51–60.
- [38] Eyre H, Kahn R, Robertson RM. Preventing cancer, cardiovascular disease, and diabetes: a common agenda for the American Cancer Society, the American Diabetes Association, and the American Heart Association. *Diabetes Care* 2004;27(7):1812–1824.
- [39] McKinnon RA, Orleans CT, Kumanyika SK, Haire-Joshu D, Krebs-Smith SM, Finkelstein EA, et al. Considerations for an obesity policy research agenda. *Am J Prev Med* 2009;36(4):351–357.
- [40] Lazarus JV, Safreed-Harmon K, Kamarulzaman A, Anderson J, Leite RB, Behrens G, et al. Consensus statement on the role of health systems in advancing the long-term well-being of people living with HIV. *Nat Commun* 2021;12(1):4450.
- [41] Rubino F, Puhl RM, Cummings DE, Eckel RH, Ryan DH, Mechanick JI, et al. Joint international consensus statement for ending stigma of obesity. *Nat Med* 2020;26(4):485–497.
- [42] Karlsen TH, Sheron N, Zelber-Sagi S, Carrieri P, Dusheiko G, Bugianesi E, et al. The EASL-Lancet Liver Commission: protecting the next generation of Europeans against liver disease complications and premature mortality. *Lancet* 2022;399(10319):61–116.
- [43] Iruzubieta P, Bataller R, Arias-Loste MT, Arrese M, Calleja JL, Castro-Narro G, et al. Research Priorities for Precision Medicine in NAFLD. *Clin Liver Dis* 2023;27(2):535–551.
- [44] Ekstedt M, Nasr P, Kechagias S. Natural History of NAFLD/NASH. *Curr Hepatol Rep* 2017;16(4):391–397.
- [45] Hagström H, Nasr P, Ekstedt M, Hammar U, Stål P, Hultcrantz R, et al. Fibrosis stage but not NASH predicts mortality and time to development of severe liver disease in biopsy-proven NAFLD. *J Hepatol* 2017;67(6):1265–1273.
- [46] Sanyal AJ, Van Natta ML, Clark J, Neuschwander-Tetri BA, Diehl A, Dasarthy S, et al. Prospective Study of Outcomes in Adults with Nonalcoholic Fatty Liver Disease. *N Engl J Med* 2021;385(17):1559–1569.
- [47] Younossi ZM, Anstee QM, Wong VWS, Trauner M, Lawitz EJ, Harrison SA, et al. The Association of Histologic and Noninvasive Tests With Adverse Clinical and Patient-Reported Outcomes in Patients With Advanced Fibrosis Due to Nonalcoholic Steatohepatitis. *Gastroenterology* 2021;160(5):1608–1619.e13.
- [48] Anstee QM, Castera L, Loomba R. Impact of non-invasive biomarkers on hepatology practice: Past, present and future. *J Hepatol* 2022;76(6):1362–1378.
- [49] Allen AM, Lazarus JV, Younossi ZM. Healthcare and socioeconomic costs of NAFLD: A global framework to navigate the uncertainties. *J Hepatol* 2023. S0168-8278(23)00079-X.
- [50] Hagström H, Nasr P, Ekstedt M, Hammar U, Widman L, Stål P, et al. Health Care Costs of Patients With Biopsy-Confirmed Nonalcoholic Fatty Liver Disease Are Nearly Twice Those of Matched Controls. *Clin Gastroenterol Hepatol* 2020;18(7):1592–1599.e8.
- [51] Talens M, Tumas N, Lazarus JV, Benach J, Pericàs JM. What Do We Know About Inequalities in NAFLD Distribution and Outcomes? A Scoping Review. *J Clin Med* 2021;10(21):5019.
- [52] Di Cesare M, Khang Y-H, Asaria P, Blakely T, Cowan MJ, Farzadfar F, et al. Inequalities in non-communicable diseases and effective responses. *Lancet* 2013;381(9866):585–597.
- [53] Eslam M, Wong GL, Hashem AM, Chan HL, Nielsen MJ, Leeming DJ, et al. A Sequential Algorithm Combining ADAPT and Liver Stiffness Can Stage Metabolic-Associated Fatty Liver Disease in Hospital-Based and Primary Care Patients. *Am J Gastroenterol* 2021;116(5):984–993.
- [54] Majumdar A, Campos S, Gurusamy K, Pinzani M, Tsochatzidis EA. Defining the Minimum Acceptable Diagnostic Accuracy of Noninvasive Fibrosis Testing in Cirrhosis: A Decision Analytic Modeling Study. *Hepatology* 2020;71(2):627–642.
- [55] Vali Y, Lee J, Boursier J, Petta S, Wonders K, Tiniakos D, et al. Biomarkers for staging fibrosis and non-alcoholic steatohepatitis in non-alcoholic fatty liver disease (the LITMUS project): a comparative diagnostic accuracy study. *Lancet Gastroenterol Hepatol* 2023. S2468-1253(23)00017-1.
- [56] Castera L, Friedrich-Rust M, Loomba R. Noninvasive Assessment of Liver Disease in Patients With Nonalcoholic Fatty Liver Disease. *Gastroenterology* 2019;156(5):1264–1281.e4.
- [57] Gidener T, Dierkhising RA, Mara KC, Therneau TM, Venkatesh SK, Ehman RL, et al. Change in serial liver stiffness measurement by magnetic resonance elastography and outcomes in NAFLD. *Hepatology* 2023;77(1):268–274.
- [58] Crossan C, Majumdar A, Srivastava A, Thorburn D, Rosenberg W, Pinzani M, et al. Referral pathways for patients with NAFLD based on non-invasive fibrosis tests: Diagnostic accuracy and cost analysis. *Liver Int* 2019;39(11):2052–2060.
- [59] Vilar-Gomez E, Lou Z, Kong N, Vuppalanchi R, Imperiale TF, Chalasani N. Cost Effectiveness of Different Strategies for Detecting Cirrhosis in Patients With Nonalcoholic Fatty Liver Disease Based on United States Health Care System. *Clin Gastroenterol Hepatol* 2020;18(10):2305–2314.e12.
- [60] Lazarus JV, Castera L, Mark HE, Allen AM, Adams LA, Anstee QM, et al. Real-world evidence on non-invasive tests and associated cut-offs used to assess fibrosis in routine clinical practice. *JHEP Rep* 2022;5(1):100596.
- [61] Glass LM, Hunt CM, Fuchs M, Su GL. Comorbidities and Nonalcoholic Fatty Liver Disease: The Chicken, the Egg, or Both? *Fed Pract* 2019;36(2):64–71.
- [62] Kumar S, Wong R, Newberry C, Yeung M, Peña JM, Sharaiha RZ. Multidisciplinary Clinic Models: A Paradigm of Care for Management of NAFLD. *Hepatology* 2021;74(6):3472–3478.
- [63] Mencin AA, Loomba R, Lavine JE. Caring for children with NAFLD and navigating their care into adulthood. *Nat Rev Gastroenterol Hepatol* 2015;12(11):617–628.
- [64] Mitrani R, Kohut T, Panganiban J, Carr RM. Transition of Care Model for Pediatric Patients With Nonalcoholic Fatty Liver Disease. *Clin Liver Dis (Hoboken)* 2021;18(1):30–36.
- [65] Nobili V, Svegliati-Baroni G, Alisi A, Miele L, Valenti L, Vajro P. A 360-degree overview of paediatric NAFLD: recent insights. *J Hepatol* 2013;58(6):1218–1229.
- [66] DeVore S, Kohli R, Lake K, Nicholas L, Dietrich K, Balistreri WF, et al. A multidisciplinary clinical program is effective in stabilizing BMI and reducing transaminase levels in pediatric patients with NAFLD. *J Pediatr Gastroenterol Nutr* 2013;57(1):119–123.
- [67] World Health Organization. WHO guideline: recommendations on digital interventions for health system strengthening. Geneva: WHO; 2019.
- [68] Lazarus JV, Villota-Rivas M, Jiménez-González C, Santos-Laso A, Iruzubieta P, Arias-Loste MT, et al. Physicians' Use of Digital Health Interventions in the Management of Nonalcoholic Fatty Liver Disease. *Clin Liver Dis* 2023;27(2):515–533.
- [69] Lim SL, Johal J, Ong KW, Han CY, Chan YH, Lee YM, et al. Lifestyle Intervention Enabled by Mobile Technology on Weight Loss in Patients With Nonalcoholic Fatty Liver Disease: Randomized Controlled Trial. *JMIR Mhealth Uhealth* 2020;8(4):e14802.
- [70] Sato M, Akamatsu M, Shima T, Ikegami T, Yanase M, Mikami S, et al. Impact of a Novel Digital Therapeutics System on Nonalcoholic Steatohepatitis: The NASH App Clinical Trial. *Am J Gastroenterol* 2023. <https://doi.org/10.14309/ajg.0000000000002143>.
- [71] Stine JG, Rivas G, Hummer B, Duarte-Rojo A, May CN, Geyer N, et al. Mobile health lifestyle intervention program leads to clinically significant loss of body weight in patients with NASH. *Hepatol Commun* 2023;7(4):e0052.
- [72] Chatterjee A, Prinz A, Gerdes M, Martinez S. Digital Interventions on Healthy Lifestyle Management: Systematic Review. *J Med Internet Res* 2021;23(11):e26931.
- [73] Zhang Y, Pratap A, Folarin AA, Sun S, Cummins N, Matcham F, et al. Long-term participant retention and engagement patterns in an app and wearable-based multinational remote digital depression study. *NPJ Digit Med* 2023;6(1):25.
- [74] Allen AM, Shah VH, Therneau TM, Venkatesh SK, Mounajjed T, Larson JJ, et al. Multiparametric Magnetic Resonance Elastography Improves the Detection of NASH Regression Following Bariatric Surgery. *Hepatol Commun* 2019;4(2):185–192.
- [75] Paudel S, Sharma N, Joshi A, Randall M. Development of a Shared Decision Making Model in a Community Mental Health Center. *Commun Ment Health J* 2018;54(1):1–6.
- [76] Tamhane S, Rodriguez-Gutierrez R, Hargraves I, Montori VM. Shared Decision-Making in Diabetes Care. *Curr Diab Rep* 2015;15(12):112.
- [77] Vilar-Gomez E, Calzadilla-Bertol L, Wong VWS, Castellanos M, Aller-de la Fuente R, Metwally M, et al. Fibrosis Severity as a Determinant of Cause-Specific Mortality in Patients With Advanced Nonalcoholic Fatty Liver Disease: A Multi-National Cohort Study. *Gastroenterology* 2018;155(2):443–457.e17.



- [78] Polyzos SA, Kechagias S, Tsochatzis EA. Review article: non-alcoholic fatty liver disease and cardiovascular diseases: associations and treatment considerations. *Aliment Pharmacol Ther* 2021;54(8):1013–1025.
- [79] Islam KB, Brandman D, Chu JN, Goldman ML, Fox RK. Primary Care Providers and Nonalcoholic Fatty Liver Disease: A Needs Assessment Survey. *Dig Dis Sci* 2023;68(2):434–438.
- [80] Vidal-Cevallos P, Ordóñez-Vázquez AL, Procopio-Mosso O, Cardoso-Arias R, Uribe M, Chávez-Tapia NC. Cross-sectional pilot study to assess primary healthcare workers' knowledge of nonalcoholic fatty liver disease in a marginalized community in Mexico. *Sci Rep* 2021;11(1):12100.
- [81] Wessels DH, Rosenberg Z. Awareness of non-alcoholic steatohepatitis and treatment guidelines: What are physicians telling us? *World J Hepatol* 2021;13(2):233–241.
- [82] ElSayed NA, Aleppo G, Aroda VR, Bannuru RR, Brown FM, Bruemmer D, et al. Introduction and Methodology: Standards of Care in Diabetes-2023. *Diabetes Care* 2023;46(Suppl 1):S1–S4.
- [83] Carrieri P, Mourad A, Marcellin F, Trylesinski A, Calleja JL, Protopopescu C, et al. Knowledge of liver fibrosis stage among adults with NAFLD/NASH improves adherence to lifestyle changes. *Liver Int* 2022;42(5):984–994.
- [84] Alemany-Pagès M, Moura-Ramos M, Araújo S, Macedo MP, Ribeiro RT, do Ó D, et al. Insights from qualitative research on NAFLD awareness with a cohort of T2DM patients: time to go public with insulin resistance? *BMC Public Health* 2020;20(1):1142.
- [85] Wieland AC, Mettler P, McDermott MT, Crane LA, Cicutto LC, Bambha KM. Low awareness of nonalcoholic fatty liver disease among patients at high metabolic risk. *J Clin Gastroenterol* 2015;49(1):e6–e10.
- [86] Assimakopoulos K, Karaivazoglou K, Tsermpini EE, Diamantopoulou G, Triantos C. Quality of life in patients with nonalcoholic fatty liver disease: A systematic review. *J Psychosom Res* 2018;112:73–80.
- [87] Golubeva JA, Sheptulina AF, Yafarova AA, Mamutova EM, Kiselev AR, Drapkina OM. Reduced Quality of Life in Patients With Non-Alcoholic Fatty Liver Disease May Be Associated With Depression and Fatigue. *Healthcare (Basel)* 2022;10(9):1699.
- [88] Shea S, Lionis C, Atkinson L, Kite C, Lagojda L, Chaggar SS, et al. Support Needs and Coping Strategies in Non-Alcoholic Fatty Liver Disease (NAFLD): A Multidisciplinary Approach to Potential Unmet Challenges Beyond Pharmacological Treatment. *Livers* 2023;3(1):1–20.
- [89] Remmers C, Hibbard J, Mosen DM, Wagenfield M, Hoye RE, Jones C. Is patient activation associated with future health outcomes and healthcare utilization among patients with diabetes? *J Ambul Care Manage* 2009;32(4):320–327.
- [90] Weingart SN, Zhu J, Chiappetta L, Stuver SO, Schneider EC, Epstein AM, et al. Hospitalized patients' participation and its impact on quality of care and patient safety. *Int J Qual Health Care* 2011;23(3):269–277.
- [91] Gershkowitz BD, Hillert CJ, Crotty BH. Digital Coaching Strategies to Facilitate Behavioral Change in Type 2 Diabetes: A Systematic Review. *J Clin Endocrinol Metab* 2021;106(4):e1513–e1520.
- [92] Milani RV, Lavie CJ, Bober RM, Milani AR, Ventura HO. Improving Hypertension Control and Patient Engagement Using Digital Tools. *Am J Med* 2017;130(1):14–20.
- [93] Hallsworth K, McPherson S, Anstee QM, Flynn D, Haigh L, Avery L. Digital Intervention With Lifestyle Coach Support to Target Dietary and Physical Activity Behaviors of Adults with Nonalcoholic Fatty Liver Disease: Systematic Development Process of VITALISE Using Intervention Mapping. *J Med Internet Res* 2021;23(1):e20491.
- [94] Mazzotti A, Caletti MT, Brodosi L, Di Domizio S, Forchielli ML, Petta S, et al. An internet-based approach for lifestyle changes in patients with NAFLD: Two-year effects on weight loss and surrogate markers. *J Hepatol* 2018;69(5):1155–1163.
- [95] Francque SM, Marchesini G, Kautz A, Walmsley M, Dörner R, Lazarus JV, et al. Non-alcoholic fatty liver disease: A patient guideline. *JHEP Rep* 2021;3(5):100322.
- [96] Lazarus JV, Mark HE, Villota-Rivas M, Palayew A, Carrieri P, Colombo M, et al. The global NAFLD policy review and preparedness index: Are countries ready to address this silent public health challenge? *J Hepatol* 2022;76(4):771–780.
- [97] World Health Organization. Global action plan for the prevention and control of NCDs 2013–2020. Geneva: WHO; 2013.
- [98] World Health Organization. Political declaration of the third high-level meeting of the General Assembly on the prevention and control of non-communicable diseases, and mental health. Geneva: WHO; 2023.
- [99] Gilmore AB, Fabbri A, Baum F, Bertscher A, Bondy K, Chang H-J, et al. Defining and conceptualising the commercial determinants of health. *Lancet* 2023;401(10383):1194–1213.

## Supplemental information

### **A global research priority agenda to advance public health responses to fatty liver disease**

**Jeffrey V. Lazarus, Henry E. Mark, Alina M. Allen, Juan Pablo Arab, Patrizia Carrieri, Mazen Nouredin, William Alazawi, Naim Alkhouri, Saleh A. Alqahtani, Marco Arrese, Ramon Bataller, Thomas Berg, Paul N. Brennan, Patrizia Burra, Graciela E. Castro-Narro, Helena Cortez-Pinto, Kenneth Cusi, Nikos Dedes, Ajay Duseja, Sven M. Francque, Hannes Hagström, Terry T.-K. Huang, Dana Ivancovsky Wajcman, Achim Kautz, Christopher J. Kopka, Aleksander Krag, Veronica Miller, Philip N. Newsome, Mary E. Rinella, Diana Romero, Shiv Kumar Sarin, Marcelo Silva, C. Wendy Spearman, Emmanuel A. Tsochatzis, Luca Valenti, Marcela Villota-Rivas, Shira Zelber-Sagi, Jörn M. Schattenberg, Vincent Wai-Sun Wong, Zobair M. Younossi, on behalf of the, and Healthy Livers, Healthy Lives Collaborators**

## Supplementary materials

# A global research priority agenda to advance public health responses to fatty liver disease

Jeffrey V. Lazarus, Henry E. Mark, Alina M. Allen, Juan Pablo Arab, Patrizia Carrieri, Mazen Nouredin, William Alazawi, Naim Alkhouri, Saleh A. Alqahtani, Marco Arrese, Ramon Bataller, Thomas Berg, Paul N. Brennan, Patrizia Burra, Graciela E. Castro-Narro, Helena Cortez-Pinto, Kenneth Cusi, Nikos Dedes, Ajay Duseja, Sven M. Francque, Hannes Hagström, Terry T-K. Huang, Dana Ivancovsky Wajcman, Achim Kautz, Christopher J. Kopka, Aleksander Krag, Veronica Miller, Philip N. Newsome, Mary E. Rinella, Diana Romero, Shiv Kumar Sarin, Marcelo Silva, C. Wendy Spearman, Emmanuel A. Tsochatzis, Luca Valenti, Marcela Villota-Rivas, Shira Zelber-Sagi, Jörn M. Schattenberg, Vincent Wai-Sun Wong, Zobair M. Younossi

The list above is of core authors. For the full authorship list, please refer to the manuscript.

## Table of Contents

Table S1. Core group members (n=40).....	2
Table S2. Wilton Park meeting delegates. ....	5
Table S3. Research priorities that achieved less than 80% agreement. ....	9

**Table S1. Core group members (n=40).**

<b>Name</b>	<b>Primary affiliation</b>
William Alazawi	Barts Liver Centre, Blizard Institute, Queen Mary University of London, London, UK
Naim Alkhouri	Fatty Liver Program, Arizona Liver Health, Phoenix, AZ, USA
Alina M. Allen	Division of Gastroenterology and Hepatology, Department of Medicine, Mayo Clinic, Rochester, MN, USA
Saleh A. Alqahtani	King Faisal Specialist Hospital & Research Centre, Riyadh, Saudi Arabia
Juan Pablo Arab	Division of Gastroenterology, Department of Medicine, Schulich School of Medicine, Western University & London Health Sciences Centre, London, Ontario, Canada
Marco Arrese	Department of Gastroenterology, School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile
Ramon Bataller	Liver Unit, Hospital Clínic, University of Barcelona, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain
Thomas Berg	Division of Hepatology, Department of Medicine II, Leipzig University Medical Center, Leipzig, Germany
Paul N. Brennan	Division of Hepatology, University of Dundee, Dundee, Scotland, UK
Patrizia Burra	Multivisceral Transplant Unit-Gastroenterology, Department of Surgery, Oncology and Gastroenterology at the Padua University Hospital, Padua, Italy
Patrizia Carrieri	Aix Marseille Univ, Inserm, IRD, SESSTIM, Sciences Economiques & Sociales de la Santé & Traitement de l'Information Médicale, ISSPAM, Marseille, France
Graciela E. Castro-Narro	Department of Hepatology and Transplant, Hospital Médica Sur, Mexico City, Mexico
Helena Cortez-Pinto (Non-voting)	Clinica Universitária de Gastrenterologia, Laboratório de Nutrição, Faculdade de Medicina, Universidade de Lisboa, Lisbon, Portugal
Kenneth Cusi	Division of Endocrinology, Diabetes & Metabolism, Department of Medicine, University of Florida, Gainesville, FL, USA
Nikos Dedes	Greek Patients Association, Athens, Greece
Ajay Duseja	Department of Hepatology, Postgraduate Institute of Medical Education and Research, Chandigarh, India
Sven M. Francque (Non-voting)	Department of Gastroenterology and Hepatology, Antwerp University Hospital, Edegem, Belgium



Hannes Hagström	Department of Medicine, Huddinge, Karolinska Institutet, Stockholm, Sweden
Terry T-K. Huang	CUNY Graduate School of Public Health and Health Policy (CUNY SPH), New York, NY, USA
Dana Ivancovsky Wajcman	Barcelona Institute for Global Health (ISGlobal), Hospital Clínic, University of Barcelona, Barcelona, Spain
Achim Kautz (Non-voting)	Kautz 5 gUG, Köln, Germany
Christopher J. Kopka	Independent researcher, Ponte de Lima, Portugal
Aleksander Krag	Department of Gastroenterology and Hepatology, Odense University Hospital, Odense, Denmark
Jeffrey V. Lazarus (Chair)	Barcelona Institute for Global Health (ISGlobal), Hospital Clínic, University of Barcelona, Barcelona, Spain
Henry E. Mark	European Association for the Study of the Liver (EASL), Geneva, Switzerland
Veronica Miller	University California Berkeley School of Public Health, Berkeley, CA, USA
Philip N. Newsome	National Institute for Health Research Birmingham Biomedical Research Centre at University Hospitals Birmingham NHS Foundation Trust and the University of Birmingham, Birmingham, UK
Mazen Nouredin	Houston Methodist Hospital, Houston Research Institute, Houston, TX, USA
Mary E. Rinella	Department of Medicine, University of Chicago, Chicago, IL, USA
Diana Romero (Non-voting)	Department of Community Health and Social Sciences, CUNY Graduate School of Public Health and Health Policy, New York, NY, USA
Shiv Kumar Sarin	Department of Hepatology, Institute of Liver and Biliary Sciences, New Delhi, India
Jörn M. Schattenberg	Metabolic Liver Research Program, I. Department of Medicine, University Medical Centre Mainz, Mainz, Germany
Marcelo Silva	Hepatology and Clinical Research Units, Hospital Universitario Austral, Buenos Aires, Argentina
C. Wendy Spearman	Division of Hepatology, Department of Medicine, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa
Emmanuel A. Tsochatzis	UCL Institute for Liver and Digestive Health, Royal Free Hospital, London, UK

Luca Valenti	Precision Medicine, Biological Resource Center, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy
Marcela Villota-Rivas (Non-voting)	Barcelona Institute for Global Health (ISGlobal), Hospital Clínic, University of Barcelona, Barcelona, Spain
Vincent Wai-Sun Wong	The Chinese University of Hong Kong, Hong Kong, China
Zobair M. Younossi	Center for Liver Disease, Inova, Falls Church, VA, USA
Shira Zelber-Sagi	School of Public Health, Faculty of Social Welfare and Health Sciences, University of Haifa, Haifa, Israel

**Table S2. Wilton Park meeting delegates.**

<b>First Name</b>	<b>Last Name</b>	<b>Position</b>	<b>Institution</b>	<b>Country</b>
William	Alazawi	Professor of Hepatology	Queen Mary University of London	UK
Naim	Alkhouri	Chief Medical Officer and Director of the Fatty Liver Program	Arizona Liver Health	USA
Alina M.	Allen	Director, NAFLD Clinic	Mayo Clinic	USA
Saleh A.	Alqahtani	Chairman, Board of Directors and Executive Director	King Faisal Specialist Hospital & Research Centre	Saudi Arabia
Juan Pablo	Arab	Associate Professor of Medicine	Western University	Canada
Marco	Arrese	Professor of Medicine	Pontificia Universidad Católica de Chile	Chile
Thomas	Berg	Secretary General of the European Association for the Study of the Liver (EASL); Head of Division of Hepatology	Leipzig University Medical Center	Germany
Jacqueline	Bowman-Busato	Policy Lead	European Association for the Study of Obesity	Belgium
Paul N.	Brennan	Clinical Lecturer and Specialist Registrar in Hepatology	University of Dundee	UK
Patrizia	Burra	Head of Multivisceral Transplant Unit	Padua University Hospital	Italy
Massimo	Colombo	Chair of the International Liver Foundation	EASL	Italy

Kenneth	Cusi	Professor of Medicine	University of Florida	USA
Nikos	Dedes	Chair	Greek Patients Association	Greece
Davide	Fortin	Post Doc	Aix Marseille University	France
Sven M.	Francque	Professor	Antwerp University Hospital	Belgium
Alexander	French	Head of Education and Capacity Building	World Obesity Federation	UK
Amalia	Gastaldelli	Research Director	Institute of Clinical Physiology	Italy
Hannes	Hagström	Consultant hepatologist	Karolinska Institutet	Sweden
Saeed	Hamid	Professor of Medicine, Director Clinical Trials Unit	Aga Khan University	Pakistan
Vanessa	Hebditch	Director of Communications and Policy	British Liver Trust	UK
Achim	Kautz	Chief Executive Officer	Kautz 5 gUG	Germany
Loreta A.	Kondili	Hepatologist Clinical Researcher	Instituto Superiore di Sanità	Italy
Eleni	Kotsiliti	Associate Editor	Nature Reviews Gastroenterology & Hepatology	Germany
Aleksander	Krag	Professor, Head of Center	Odense University Hospital	Denmark
Jeffrey V.	Lazarus	Head of Health Systems Research Group and Associate Research Professor	Barcelona Institute for Global Health (ISGlobal)	Spain
Maria Paula	Macedo	Professor	NOVA Medical School	Portugal

Henry E.	Mark	Consultant Programme Director	Wilton Park	UK
Juan M.	Mendive	Family Physician	Catalan Health Institute	Spain
Yoanna	Nedelcheva	Policy, Advocacy and Public Health Manager	EASL	Switzerland
Mazen	Noureddin	Director	Houston Research Institute	USA
Rachel	Pryke	General practitioner	Bewdley Medical Centre	UK
Mary E.	Rinella	Professor of Medicine	University of Chicago	USA
Jörn M.	Schattenberg	Director, Metabolic Liver Research Program	University Medical Centre Mainz	Germany
Marcelo	Silva	Hepatology Consultant	Hospital Universitario Austral	Argentina
C. Wendy	Spearman	Head of the Division of Hepatology, Department of Medicine, Faculty of Health Sciences	University of Cape Town	South Africa
Emmanuel A.	Tsochatzis	Professor of Hepatology	UCL Institute for Liver and Digestive Health	UK
Luca	Valenti	Professor of Internal Medicine	Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico	Italy
Zobair M.	Younossi	President, Inova Medicine	Inova Health System	USA
Volkan	Yumuk	Professor of Medicine	Istanbul University-Cerrahpaşa	Turkey



Shira	Zelber-Sagi	Head of School of Public Health, Faculty of Social Welfare and Health Sciences	University of Haifa	Israel
-------	-------------	---	---------------------	--------

**Table S3. Research priorities that achieved less than 80% agreement.**

<b>Statement</b>		<b>A (%)</b>	<b>SA (%)</b>
<b>Section 1: The human and economic burden</b>			
1.5	Report all data disaggregated by sex, race, ethnicity, age, socioeconomic status, education level, and other variables related to inequities.	74.2	23.3
<b>Section 2: Defining and implementing models of care</b>			
2.5	Explore how novel digital technologies (e.g., artificial intelligence, data-based analytics, digital health applications and therapeutics) can be utilised within healthcare settings.	73.2	21.3
2.6	Assess how digital health (e.g., applications, interventions, therapeutics) can support patients to achieve lifestyle behavioural change.	69.8	26.7
2.7	Further explore the use of artificial intelligence to improve diagnostics for fatty liver disease.	63.9	31.2
<b>Section 3: Treatment and care</b>			
3.3	Evaluate patient-centred decision making in relation to fatty liver disease treatment and care outcomes.	79.5	16.3
<b>Section 4: Education and awareness</b>			
4.3	Study the effectiveness of strategies to impact fatty liver disease knowledge, attitudes, beliefs, and practices (KABPs), prioritising KABPs among healthcare professionals and high-risk groups.	72.8	25.4
<b>Section 5: Patient and community perspectives</b>			

5.1	Conduct research to understand the needs and experiences of fatty liver disease patients and at-risk communities (e.g., perspectives around prevention, treatment, and care, including mental health).	78.0	19.2
5.3	Assess if published patient guidelines result in patients having an improved understanding of fatty liver disease and taking a more active role in their disease management.	71.9	25.7
5.4	Evaluate the efficacy of patient-led self-care programmes in improving fatty liver disease outcomes.	74.6	22.0
5.5	Explore the potential of new technologies (e.g., digital health applications and therapeutics, mobile interventions) to foster patient engagement in treatment and care.	75.5	21.3
5.6	Evaluate the effect of interventions to reduce liver disease stigma among patients, the public, and healthcare providers.	74.7	21.2
<b>Section 6: Leadership and policies for the fatty liver disease public health agenda</b>			
6.1	Conduct periodic studies of national and sub-national policies and guidelines for the prevention and management of fatty liver disease, to identify trends and gaps, and assess their implementation.	79.4	18.8
6.3	Monitor, study, and report mentions of fatty liver disease within patient groups and professional societies outside of the field of hepatology (e.g., at events, in publications).	66.8	28.0
A, agree; SA, somewhat agree.			