

PAPER • OPEN ACCESS

## Quantum transfer of interacting qubits

To cite this article: Tony J G Apollaro *et al* 2022 *New J. Phys.* **24** 083025

View the [article online](#) for updates and enhancements.

You may also like

- [Effective Hamiltonians for interacting superconducting qubits: local basis reduction and the Schrieffer–Wolff transformation](#)  
Gioele Consani and Paul A Warburton
- [Exact Entanglement Dynamics in Three Interacting Qubits](#)  
Wen-Bin He, , Xi-Wen Guan et al.
- [Entanglement and the speed of evolution of two interacting qubits](#)  
C Zander, A Borras, A R Plastino et al.



## PAPER

## Quantum transfer of interacting qubits

## OPEN ACCESS

RECEIVED  
9 May 2022REVISED  
20 July 2022ACCEPTED FOR PUBLICATION  
4 August 2022PUBLISHED  
23 August 2022

Original content from  
this work may be used  
under the terms of the  
[Creative Commons  
Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/).  
Any further distribution  
of this work must  
maintain attribution to  
the author(s) and the  
title of the work, journal  
citation and DOI.

Tony J G Apollaro<sup>1,\*</sup> , Salvatore Lorenzo<sup>2</sup> , Francesco Plastina<sup>3,4</sup>,  
Mirko Consiglio<sup>1</sup>  and Karol Życzkowski<sup>5,6</sup><sup>1</sup> Department of Physics, University of Malta, Msida MSD 2080, Malta<sup>2</sup> Università degli Studi di Palermo, Dipartimento di Fisica e Chimica - Emilio Segrè, via Archirafi 36, I-90123 Palermo, Italy<sup>3</sup> Dipartimento di Fisica, Università della Calabria, 87036 Arcavacata di Rende (CS), Italy<sup>4</sup> INFN, Gruppo Collegato di Cosenza, Italy<sup>5</sup> Institute of Theoretical Physics, Jagiellonian University, ul. Łojasiewicza 11, 30-348 Kraków, Poland<sup>6</sup> Center for Theoretical Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668 Warszawa, Poland

\* Author to whom any correspondence should be addressed.

E-mail: [tony.apollaro@um.edu.mt](mailto:tony.apollaro@um.edu.mt)**Keywords:** high-dimensional quantum state transfer, quantum dynamical maps, quantum spin chains, quantum information processing, open quantum systems**Abstract**

The transfer of quantum information between different locations is key to many quantum information processing tasks. Whereas, the transfer of a single qubit state has been extensively investigated, the transfer of a many-body system configuration has insofar remained elusive. We address the problem of transferring the state of  $n$  interacting qubits. Both the exponentially increasing Hilbert space dimension, and the presence of interactions significantly scale-up the complexity of achieving high-fidelity transfer. By employing tools from random matrix theory and using the formalism of quantum dynamical maps, we derive a general expression for the average and the variance of the fidelity of an arbitrary quantum state transfer protocol for  $n$  interacting qubits. Finally, by adopting a weak-coupling scheme in a spin chain, we obtain the explicit conditions for high-fidelity transfer of three and four interacting qubits.

**1. Introduction**

Quantum information processing (QIP) is shaping the 21st century technology by means of the advantage it provides, with respect to its classical counterpart, in fields ranging from computation to cryptography [1]. A basic building block of several QIP protocols is the transfer of quantum information between different locations. In particular, a great variety of different protocols have been devised to achieve the high fidelity transfer of the quantum state of a single qubit (one-QST). They can be classified into three broad classes: protocols employing flying qubits [2], teleportation-based ones [3], and those employing spin- $\frac{1}{2}$  chains as quantum data bus [4]. In this paper, we will focus on the last approach. Since the seminal paper of Bose [4], numerous one-QST protocols have been proposed, and implemented, in systems ranging from evanescently coupled optical waveguides [5, 6] and cavity-coupled atoms and ions [7], to transmon qubits [8], nitrogen vacancy centers [9] and Rydberg atoms [10], just to name a few.

As a direct generalization of the original set-up, the quantum state transfer of  $n$ -qubits ( $n$ -QST) has received some attention in the last few years, with a focus on the transfer of entangled qubit states. In the majority of cases, however, the state to be sent is that of a set of non-interacting qubits [11–14]. On the other hand, the transfer of the quantum state of *interacting*  $n$ -body systems has been insofar barely addressed. In fact, for many QIP protocols, fast and efficient non-interacting  $n$ -QST would already constitute an important achievement: in distributed quantum computing architectures [15], for instance, the  $n$ -qubit output state of a computation has to be distributed among different quantum processors. In quantum secret sharing protocols an entangled  $n$ -qubit state is shared among several users [16], while a fully-fledged quantum internet [17] requires nodes capable of exchanging, ideally, arbitrary many-body states. However, the quantum state transfer of interacting qubits ( $n$ -iQST) could be far more beneficial both for QIP protocols (where, e.g., qubit-decoupling operations prior to transfer could be avoided), and, in

perspective, to also achieve the transfer of the full physical configuration of complex systems, where complexity is embodied not only by the dimensionality of the system's Hilbert space [18] but also by the interactions among its constituents [19]. The transfer of an interacting system's state may also constitute a significant advantage in those experiments with many-body systems, where state preparation is not easily feasible in the same set-up in which the rest of the experiment is performed. In these cases, one could, e.g., prepare the state embodying the properties under investigation in a different location and send it later, via a quantum channel, in order to load it into the main set-up.

Although being closely related,  $n$ -QST and  $n$ -iQST represent two different aspects of quantum transfer, with the latter being a non-trivial generalisation of the former. In the non-interacting case, indeed, the  $n$ -qubit state could in principle be transferred sequentially, with qubits sent one by one. On the other hand, if we aim at transferring the quantum state while the system is interacting, sequential transfer is not feasible as it has to be accomplished simultaneously for the entire system. Similarly to  $n$ -QST, for  $n$ -iQST, the recipient is required to have a physical system able to store the received quantum information. In our case, it is sufficient to work with a qubit system with the same interaction scheme of the one that is sent. Therefore, the quality of a  $n$ -iQST can be assessed with the same figure of merits as those used for  $n$ -QST; namely, the fidelity.

The  $n$ -QST problem has been posed shortly after one-QST via spin- $\frac{1}{2}$  chain was proposed. Following the protocol for perfect state transfer, which entails fully-engineered couplings of the quantum chain, it has been shown that mirror-periodic Hamiltonians allow for mirror-inversion of an arbitrary quantum state with respect to the center of the chain [20, 21]. However, in order to apply the same idea to achieve  $n$ -iQST, one would require a fully-engineered quantum channel, implying, in general, a modification of the couplings among the qubits embodying the sender's system. On the other hand, uniformly-coupled chains allow for high-quality  $n$ -QST for specific lengths related to prime number theory [22], and extensive research has been devoted to investigate the transfer of few-qubit entangled states over such spin chains [12–14, 23–26]. However, a general approach to the  $n$ -QST problem has not been put forward yet, and, moreover, specific  $n$ -QST protocols may not be applicable to the  $n$ -iQST case due to the dynamical evolution of the sender's qubits.

Because of the exponential increase of the Hilbert space dimension  $d = 2^n$  of an  $n$ -qubit state, QST protocols based on local operations and classical communication (LOCC) perform poorly, displaying a maximal fidelity scaling as  $\frac{2}{d+1}$  [27]. It is, therefore, of the utmost importance to identify the conditions on a quantum dynamical map that allows it to surpass the LOCC limit of  $n$ -QST, and to single out physical models that realise these maps.

In this paper, we address the  $n$ -iQST problem via a novel approach, by combining quantum dynamical maps [18] and random matrix theory [28], in order to derive the average fidelity, and its variance, of an arbitrary  $n$ -iQST protocol. Our approach will be similar to that employed in reference [29] to determine the average fidelity for the QST of a qudit. We find that the  $n$ -iQST average fidelity can be decomposed into two contributions. A first, classical one achieving the LOCC limit, resulting only from map elements connecting the diagonal elements of the sender and receiver density matrices, and a second, quantum contribution embodied by the map elements connecting off-diagonal density matrix elements of the sender and the receiver. Finally, we apply our information-theoretical formalism to the  $n$ -iQST via a spin- $\frac{1}{2}$  chain obtaining the conditions for high-quality iQST via a uniformly coupled system for  $n = 3$  and  $n = 4$  spins, utilising a generalization of the weak-coupling protocol already employed for one- and two-QST [30–33].

The paper is organised as follows: in section 2 we set the stage and derive expressions for the average fidelity and for the variance of fidelity of an arbitrary  $n$ -iQST protocol in terms of dynamical map elements. After having discussed in section 3 the special case of sequential transmission, which can be employed for  $n$ -QST only, in section 4, we derive the dynamical map elements for a quantum channel modeled by an  $U(1)$ -symmetric spin- $\frac{1}{2}$  Hamiltonian. We employ these results in section 5, where we show that, for a quantum channel modeled by an  $XX$  Hamiltonian, efficient three- and four-iQST is achievable via the weak-coupling protocol; in section 6 we draw our conclusions. Finally, in the appendix we report the explicit expressions for the elements of the dynamical maps used in the paper.

## 2. Average fidelity and variance

As stated in the introduction, to assess the performance of an  $n$ -iQST protocol, the same quantum-information theoretical tools used for  $n$ -QST can be utilised. An important figure of merit for the efficiency of a QST protocol is the average fidelity  $\langle F \rangle$ , defined as the fidelity averaged over all pure input states with respect to the unitarily invariant measure,

$$\langle F \rangle = \frac{1}{\Omega} \int_{\Omega} d\Omega F(|\Psi\rangle, \rho(t)), \quad (1)$$

with  $\Omega$  denoting the space of pure states and

$$F(|\Psi\rangle, \rho(t)) = \langle \Psi | \rho | \Psi \rangle \quad (2)$$

is the Uhlmann–Jozsa fidelity [34]. The evaluation of  $\langle F \rangle$  requires a parametrization of the pure state vector space in order to carry out integration. While this can be easily done for systems having low Hilbert space dimension (as it is the case for one [4] and two qubits [33]), for an arbitrary pure state in  $d$ -dimensions, one needs to integrate over  $2(d-1)$  real parameters. Alternative methods have been devised, involving invariant integration over the  $SU(d)$  group [35, 36], or algebraic approaches using products of Pauli matrices [37]. However, in much of the existing literature about high-dimensional systems, the fidelity between quantum states has been used in a purely information-theoretical perspective, see, e.g., reference [38] and references therein, and very little is known about the dynamics of the average fidelity in technologically relevant scenarios, such as the interacting many-body quantum state transfer protocols we are interested in.

Here, we develop an approach to the  $n$  interacting qubits QST combining the formalism of quantum dynamical maps and invariant  $SU(d)$  group integration, in order to obtain the average fidelity of an arbitrary  $n$ -qubit QST protocol. To this end, consider the map  $\Lambda : \mathcal{S} \rightarrow \mathcal{R}$  that sends the input state of the sender into the output state of the receiver [39]

$$\hat{\rho}^R = \Lambda[\hat{\rho}^S]. \quad (3)$$

Taking an initial pure state for the sender  $\hat{\rho}^S = |s\rangle\langle s|$ , and expressing the map  $\Lambda$  in its Kraus decomposition  $\Lambda[\hat{\rho}] = \sum_k \hat{E}_k \hat{\rho} \hat{E}_k^\dagger$ , with  $\sum_k \hat{E}_k^\dagger \hat{E}_k = \mathbf{I}$ , the output fidelity reads

$$F = \sum_k \langle s | \hat{E}_k | s \rangle \langle s | \hat{E}_k^\dagger | s \rangle = \sum_k |\langle s | \hat{E}_k | s \rangle|^2. \quad (4)$$

The average fidelity over all possible input states can be obtained by integrating equation (4) with respect to the Haar measure on the unitary group  $U_{\mathcal{S}}$  acting on  $\mathcal{S}$  [40]:

$$\langle F \rangle = \frac{1}{\Omega} \int_{\Omega} d\Omega \sum_k |\langle s | \hat{E}_k | s \rangle|^2 = \sum_k \int dU_{\mathcal{S}} \left| \langle \tilde{s} | \hat{U}_{\mathcal{S}}^\dagger \hat{E}_k \hat{U}_{\mathcal{S}} | \tilde{s} \rangle \right|^2, \quad (5)$$

where we write the input state  $|s\rangle$  as the unitarily transformed reference state  $|\tilde{s}\rangle$ .

Performing integration we arrive at

$$\begin{aligned} \langle F \rangle &= \sum_k \frac{1}{d_{\mathcal{S}}(d_{\mathcal{S}}+1)} \left( \text{Tr}\{\hat{E}_k^\dagger \hat{E}_k\} + \text{Tr}\{\hat{E}_k^\dagger\} \text{Tr}\{\hat{E}_k\} \right) \\ &= \frac{1}{(d_{\mathcal{S}}+1)} + \frac{1}{d_{\mathcal{S}}(d_{\mathcal{S}}+1)} \sum_k |\text{Tr}\{\hat{E}_k\}|^2, \end{aligned} \quad (6)$$

where  $d_{\mathcal{S}}$  stands for the dimension of the Hilbert space  $\mathcal{S}$ . Note that the same expression can also be used to evaluate the average gate fidelity of a quantum channel [41, 42]. Expressing the input state  $|s\rangle$  in some complete orthonormal basis of the sender Hilbert space  $\{|0\rangle_{\mathcal{S}}, \dots, |d_{\mathcal{S}}-1\rangle_{\mathcal{S}}\}$ ,

$$\hat{\rho}^R = \Lambda[\hat{\rho}^S] = \sum_k \sum_{n,m=0}^{d_{\mathcal{S}}-1} a_n a_m^* \hat{E}_k |n\rangle_{\mathcal{S}} \langle m| \hat{E}_k^\dagger, \quad (7)$$

and choosing a basis also for the receiver Hilbert space  $\mathcal{R}$ ,  $\{|0\rangle_{\mathcal{R}}, \dots, |d_{\mathcal{R}}-1\rangle_{\mathcal{R}}\}$ , we obtain

$$\begin{aligned} \hat{\rho}^R &= \Lambda[\hat{\rho}^S] = \sum_k \sum_{n,m=0}^{d_{\mathcal{S}}-1} \sum_{i,j=0}^{d_{\mathcal{R}}-1} a_n a_m^* \langle i | \hat{E}_k | n \rangle_{\mathcal{S}} \langle m | \hat{E}_k^\dagger | j \rangle_{\mathcal{R}} |i\rangle_{\mathcal{R}} \langle j| \\ &= \sum_{n,m=0}^{d_{\mathcal{S}}-1} \sum_{i,j=0}^{d_{\mathcal{R}}-1} A_{ij}^{nm} a_n a_m^* |i\rangle_{\mathcal{R}} \langle j|, \end{aligned} \quad (8)$$

and, hence, define the elements of the map  $\Lambda$

$$A_{ij}^{nm} = \sum_k \langle i | \hat{E}_k | n \rangle_{\mathcal{S}} \langle m | \hat{E}_k^\dagger | j \rangle_{\mathcal{R}}. \quad (9)$$

The dynamical map in equation (8) is usually represented as a  $ij \times nm$  matrix acting on the input (sender) density matrix  $\rho^S$  expressed as a column vector and giving the output (receiver) density matrix,  $\vec{\rho}^R = A(t)\vec{\rho}^S$ . In the following, we will consider only the case where the Hilbert spaces  $\mathcal{S}$  and  $\mathcal{R}$  have the same dimensions, i.e.  $d_S = d_R = d$ , as our aim is to apply the present formalism to the  $n$ -iQST protocol.

The fidelity between  $\vec{\rho}^R$  and an arbitrary pure input state

$$|\Psi\rangle = \sum_{p=0}^{d-1} a_p |p\rangle_S, \tag{10}$$

is given in terms of the dynamical map elements in equation (9), by

$$F(|\Psi\rangle, \rho) = \langle \Psi | \rho | \Psi \rangle = \sum_{pqijnm=0}^{d-1} a_p^* a_q a_n a_m^* A_{ij}^{nm} \delta_{pi} \delta_{jq} = \sum_{ijnm=0}^{d-1} a_i^* a_j a_n a_m^* A_{ij}^{nm}, \tag{11}$$

where all of the  $a$ 's refer to the initial state of equation (10), while the  $\delta$ 's arise from choosing the same basis for  $\rho^S$  and  $\rho^R$ . Although from equation (11) the full probability distribution function (PDF) for the fidelity can be derived for a given map, we will focus, in the following, only on the first and second moments of the distribution for arbitrary maps. Related results for the PDF between arbitrary quantum states can be found in reference [43].

In order to evaluate the average fidelity, we use the results of reference [36], where it is shown that the only non-zero averages are given by

$$\langle |a_i|^2 \rangle = \frac{1}{d}, \quad \langle |a_i|^4 \rangle = \frac{2}{d(d+1)}, \quad \langle |a_i|^2 |a_j|^2 \rangle_{i \neq j} = \frac{1}{d(d+1)}, \tag{12}$$

which, in our case, yield

$$\langle F \rangle = \frac{1}{d(d+1)} \left( 2 \sum_{i=0}^{d-1} A_{ii}^{ii} + \sum_{i \neq j=0}^{d-1} A_{ij}^{ij} + 2 \Re \left\{ \sum_{i>j=0}^{d-1} A_{ij}^{ij} \right\} \right). \tag{13}$$

Similarly to the results of references [41, 42, 44–46], Equation (13) provides quite a simple expression for the average fidelity in terms of the quantum dynamical map elements (notice, indeed, that only  $\frac{d}{2}(3d-1)$ , out of  $d^4$ , of the quantum map's elements really matter). It will be exploited many times in the following, in order to assess the performance of  $d$ -dimensional QST in various physically relevant cases.


It is straightforward to show that equation (13) encompasses both the trivial case  $\Phi(t) = \mathbf{I}$ , i.e.,  $A_{ij}^{nm} = \delta_{in} \delta_{jm}$ , yielding  $\langle F \rangle = 1$ , and the LOCC-limit. The latter, in particular, is obtained from the first two contributions to  $\langle F \rangle$ , which come from those elements of the map  $A$  connecting all of the diagonal elements of the density matrices of  $S$  and  $R$ . The third term, instead, is due to off-diagonal map elements, connecting input to output coherences. It has, thus, a purely quantum origin, and it disappears for a classical map. The LOCC limit is obtained by setting  $A_{ij}^{nm} = \delta_{ij} \delta_{nm}$ , yielding  $\langle F \rangle = \frac{2}{d+1}$ . This is the maximum possible value achievable from the first two terms only, as one can infer by using the following constraints on the map elements, which are obtained from the fact that  $A$  represents a CPTP map in the chosen basis:

$$\begin{aligned} \sum_i A_{ii}^{nm} &= \sum_i \sum_k \mathcal{R} \langle i | \hat{E}_k | n \rangle_S \langle m | \hat{E}_k^\dagger | i \rangle_{\mathcal{R}} \\ &= \sum_i \sum_k \mathcal{S} \langle m | \hat{E}_k^\dagger | i \rangle_{\mathcal{R}} \langle i | \hat{E}_k | n \rangle_S \\ &= \mathcal{S} \langle m | \sum_k \hat{E}_k^\dagger \hat{E}_k | n \rangle_S \\ &= \delta_{nm}, \end{aligned} \tag{14}$$

$$\sum_i A_{ii}^{nm} = 1 \rightarrow 0 \leq A_{ii}^{nn} \leq 1, \quad \sum_i A_{ii}^{mm} = d, \tag{15}$$

$$A_{ij}^{nm} = (A_{mn}^{ji})^*, \quad A_{ij}^{ij} = \left( A_{ji}^{ii} \right)^*. \tag{16}$$

Thus, the last quantum term is crucial for a good performance of the channel, and it becomes more and more relevant by increasing the dimension  $d$ , as it may amount to a value up to  $1 - \frac{2}{d-1} = \frac{d-1}{d+1}$ . As a final comment to the expression above for  $\langle F \rangle$  in equation (13), we notice that map's elements connecting different off-diagonal elements of  $\rho^R$  and  $\rho^S$  do not play any role, as they are averaged out by the integral over unitaries.



### 2.1. Variance of the fidelity

Up to now, we have focused on the average fidelity. However, this does not give a complete characterization of the performance of a quantum state transfer protocol, and thus, we shall now turn our attention to the variance of the fidelity distribution, which provides a description of the dispersion of the values of  $F$  for different input states. To obtain the variance, which is defined in the usual way as

$$(\Delta F)^2 = \langle F^2 \rangle - \langle F \rangle^2, \quad (17)$$

we need to evaluate the second moment of the distribution of the possible values of the fidelity taken over all pure input states, equation (11), which is obtained by averaging the following expression

$$F^2(|\Psi\rangle, \rho) = \sum_{ijnmpqrs=0}^{d-1} a_i^* a_j a_n a_m^* a_p^* a_q a_r a_s^* A_{ij}^{nm} A_{pq}^{rs}. \quad (18)$$

Using, again, the results of reference [47], we get

$$\begin{aligned} \langle F^2 \rangle &= \frac{1}{d(d+1)(d+2)(d+3)} \\ &\times \sum_{i,m,p,s=0}^{d-1} \left( (A_{ii}^{mm} + A_{im}^{im})(A_{pp}^{ss} + A_{ps}^{ps}) + (A_{ii}^{pm} + A_{ip}^{im})(A_{pm}^{ss} + A_{ps}^{ms}) \right. \\ &+ (A_{ii}^{sm} + A_{is}^{im})(A_{pm}^{ps} + A_{pp}^{ms}) + (A_{im}^{pm} + A_{ip}^{mm})(A_{pi}^{ss} + A_{ps}^{is}) \\ &\left. + (A_{im}^{sm} + A_{is}^{mm})(A_{pi}^{ps} + A_{pp}^{is}) + (A_{ip}^{sm} + A_{is}^{pm})(A_{pi}^{ms} + A_{pm}^{is}) \right). \quad (19) \end{aligned}$$


As an easy check for this expression, if the map is the unit map, i.e.,  $A_{ij}^{nm} = \delta_{in}\delta_{jm}$ , then  $\langle F^2 \rangle = 1$ . In the next sections, we will explicitly evaluate the second moment and the variance for the case of an  $n$ -qubit channel, both in the absence and in the presence of interactions. Let us stress that the formalism outlined in this section is not limited to unitary evolutions of the quantum channel as the formulation in terms of Krauss operators allow us to include open system dynamics scenarios.

### 3. $n$ -QST over independent channels

Before turning our attention to the transfer of  $n$  interacting qubits via a single quantum channel, embodied by a spin- $\frac{1}{2}$  chain, let us analyse the case of  $n$  non-interacting qubits transferred ‘in parallel’ across  $n$ -independent channels. The topology of the independent channels can be arbitrary, and here we model them as  $U(1)$ -symmetric spin- $\frac{1}{2}$  networks as depicted in figure 1. For two- and three-dimensional spin networks allowing for perfect QST see, e.g., reference [48].

Besides being interesting in itself, this case will serve as a benchmark to compare the performance of other, more involved, transmission set-ups.

With this approach, we need to transfer one qubit per channel; thus, we can make use of the well known result that, for an initially fully polarized channel + receiver system, one-QST is completely determined by a



single parameter, the transition amplitude for one-spin excitation to be transferred from sender to receiver [4]. Such a transition amplitude, that can be taken to be real (see also next section), and that we call here  $f$ , can be manipulated in various ways, using any of the control schemes reported in the literature. Whatever approach one employs to maximize  $f$ , once its value is set, the dynamical map representing the single qubit transmission has the amplitude damping form, and it is reported in equation (A.1) of appendix A.

Under the assumption that the dynamical maps  $A$  of the various parallel transmission channels are all the same, the  $n$ -QST average fidelity of equation (13) becomes

$$\langle F_n \rangle = \frac{1}{d+1} + \frac{1}{d(d+1)} |1+f|^{2n}. \quad (20)$$

In order to overcome the classical  $n$ -QST LOCC limit, the single-particle amplitude  $f$  has to exceed the one-QST LOCC limit,  $f > \sqrt{2} - 1$ . This is independent of the number of qubits  $n$  and results in a polynomially decaying fidelity for decreasing  $f$ . In figure 2 we report the  $n$ -QST average fidelity as a function of the transition amplitude  $f$  and show that for a large number of qubits, e.g.,  $n \simeq 20$ , high-quality QST is achieved only for almost-unit single-particle transition amplitude.

From equation (20), we notice that the average fidelity  $\langle F_n \rangle \neq \prod_{i=1}^n \langle F_1 \rangle$ . In fact, the product of  $\langle F_1 \rangle$  gives the average fidelity of the QST only for fully factorized states, i.e., if  $|\Psi\rangle_n = \bigotimes_{i=1}^n |\psi\rangle_i$ . As, in general, for independent processes, the average of the product of a set of random variables is equal to the product of their averages, we conclude that, when  $n > 2$ , entanglement gives rise to a breakdown of independence in the parallel transfer processes, and it does so in such a way as to reduce the fidelity.



In order to quantify the effect of entanglement on the  $n$ -QST with the set-up reported in figure 1, we introduce, as a figure of merit, the ratio  $R$  between the fidelity of the  $n$ -QST of the subset of product states and of the full set of states.  $R$  can then be expressed both as a function of the transition amplitude  $f$  and as a function of the fidelity.

In the first case, we obtain

$$R(f) = \frac{\langle F_1 \rangle^n}{\langle F_n \rangle} = \frac{3^{-n}(2^n + 1)(f(f+2) + 3)^n}{(f+1)^{2n} + 2^n} \geq 1, \quad (21)$$

with equality holding for  $f = 0, 1$ . The maximum of  $R(f)$  occurs precisely at the amplitude value  $f_{\text{LOCC}} = \sqrt{2} - 1$  that saturates the LOCC-limit, yielding  $R(f_{\text{LOCC}}) = \frac{1}{2} \left( \left( \frac{2}{3} \right)^n + \left( \frac{4}{3} \right)^n - 2 \right)$ . Equation (21) is displayed in the left panel of figure 3, which shows that, at fixed transition amplitude  $f$ , product states enjoy a higher  $n$ -QST fidelity than entangled states. To further sustain the claim that product states are better transferred than entangled ones by independent channels, it is instructive to report also the ratio,  $R(F)$ , between the  $n$ -QST fidelity averaged over product states only, and over the full set of states at a given value,  $F$ , of the average fidelity. From equation (20), we obtain that  $\langle F_n \rangle = F$  when the transition amplitude attains the value  $f_F = \sqrt{2} \left( (2^n + 1) \left( F - \frac{1}{2^{n+1}} \right) \right)^{\frac{1}{2n}} - 1$ . Using this value of the transition amplitude,  $f_F$ , into the  $n$ -QST average fidelity over the subset of product states  $\langle F_1 \rangle^n$ , we get the ratio  $R$  to be

$$R(F) = \frac{\left( 1 + (2^n F + F - 1)^{\frac{1}{n}} \right)^n}{3^n F} \xrightarrow{n \rightarrow \infty} F^{-\frac{1}{3}}. \quad (22)$$



In the right panel of figure 3, we plot the ratio in equation (22) and show that, for  $n > 1$ , at a fixed fidelity of the full set of states, the subset of product states enjoys a higher  $n$ -QST average fidelity.

Let us now turn our attention to the variance evaluated for the case of independent channels. The variance for an  $n$ -qubit arbitrary state is given by equation (17), whereas the variance restricted to product states input follows the law  $(\Delta F)^2 = \langle F_1^2 \rangle^n - \langle F_1 \rangle^{2n}$ , where  $F_1$  is the one-QST fidelity. From the left panel in figure 4 we see that the variance for product states is greater for high-average fidelity values. Intuitively, this can be explained by noticing that the set of separable pure states is of zero measure within the set of pure states. Hence, there are less states in the neighborhood of the sender state giving the targeted fidelity, resulting in a flatter PDF of the fidelity, and an increased variance with respect to the variance obtained for the full set of pure states.

To conclude this section, we report the coefficient of variation  $CV$  as a figure of merit of the relative variability for the fidelity

$$CV = \frac{\sigma}{\langle F_n \rangle} = \sqrt{\frac{\langle F_n^2 \rangle}{\langle F_n \rangle^2} - 1}, \quad (23)$$

where  $\sigma = \Delta F$  is the standard deviation. For the case of independent channels we are dealing with here,  $CV$  is reported in the right panel of figure 4 as a function of  $\langle F_n \rangle$ . From the plot, we can conclude that, as perhaps expected, the relative dispersion becomes smaller as the average fidelity increases. Moreover, at fixed fidelity,  $CV$  tends to zero with increasing  $n$ , meaning that the average fidelity  $\langle F_n \rangle$  is a self-averaging quantity, for  $n \gg 1$  and  $\langle F_n \rangle > \frac{1}{d}$ .





**Figure 5.** Setup of an  $n$  interacting qubit state transfer protocol ( $n$ -iQST). The sender and receiver blocks are weakly coupled by  $J_0$  at both edges of a wire. Each part consists of a 1D lattice described by the Hamiltonian in equation (33) with (i) nearest neighbor couplings only, (ii)  $\Delta = 0$ , (iii) uniform parameters (that is,  $J_i = J$  and  $h_i = h \forall i$ ) and (iv)  $J_0 \ll J$ .  $n_s$  ( $= n_r$ ) indicates the number of sender (receiver) qubits, respectively in red and blue,  $n_w$  the number of qubits in the wire, shown in green, while  $N = n_s + n_r + n_w$  is the total number of qubits in the system described by the Hamiltonian in equation (33).

#### 4. $n$ -iQST via spin chains

In this section, we discuss quantum transfer for  $n$  interacting qubits via spin chains, where the sender's, the receiver's as well as the wire's qubit dynamics are described by the same Hamiltonian. Whereas the latter condition is not a requirement for the validity of the formalism outlined in section 2, our aim here is to showcase an efficient  $n$ -iQST protocol, and we will leave to future works the investigation of efficient  $n$ -iQST protocols where the Hamiltonians of the sender, the receiver and the wire differ. A much investigated QST protocol models the channel as an open boundary linear spin- $\frac{1}{2}$  chain [49], where the sender and the receiver are located at opposite edges, see figure 5 for a pictorial representation of the protocol applied to  $n$ -qubit QST via weak-coupling. This protocol has been shown to allow for both one- and two-qubit high-fidelity QST under a variety of different dynamical parameters [30, 50–53]. For  $n > 2$ , however, only a few results have been obtained up to now.

Here, after revisiting the one- and two-QST using our formalism, we will move to  $n > 2$  iQST, and propose a protocol for high-quality three and four-iQST. In particular, we will analyze the set-up where the whole sender + wire + receiver system has  $U(1)$  symmetry, i.e., it preserves the total magnetization along the  $z$ -axis, and we assume both our channel (i.e., the wire), and the receiver qubits to be initially fully polarized. This class of  $U(1)$ -symmetric Hamiltonians encompasses a large number of models, including, e.g., the following general Heisenberg-type Hamiltonian

$$H = \sum_n \sum_r (J_{n,r} (\hat{\sigma}_n^x \hat{\sigma}_{n+r}^x + \hat{\sigma}_n^y \hat{\sigma}_{n+r}^y) + \Delta_{n,r} \hat{\sigma}_n^z \hat{\sigma}_{n+r}^z + h_n \hat{\sigma}_n^z), \quad (24)$$

where the sum runs over lattice sites  $n$  and interaction range  $r$ , and where we allowed for an anisotropic exchange coupling between spins along the  $z$ -axis (so that, in general,  $J_{n,r} \neq \Delta_{n,r}$ ), and for an external transverse field  $h_n$ .

An arbitrary initial pure state, having up to  $n_s$  excitations, i.e., spin-up states  $|1\rangle$ , located in the sender block, with the rest of the system being in  $|\mathbf{0}\rangle \equiv |00\dots 00\rangle$ , can be written in the computational basis (with  $|n\rangle \equiv |0_1, 0_2, \dots, 1_n \dots 0\rangle$ ), as

$$\begin{aligned} |\Psi(0)\rangle &= a_0 |\mathbf{0}\rangle + \sum_{n \in \{n_s\}} a_n |n\rangle + \sum_{n < m \in \{n_s\}} a_{nm} |nm\rangle \\ &+ \sum_{n < m < p \in \{n_s\}} a_{nmp} |nmp\rangle + \sum_{n < m < p < q \in \{n_s\}} a_{nmpq} |nmpq\rangle + \dots \end{aligned} \quad (25)$$

According to the dynamics generated by the Hamiltonian in equation (24), such an initial state evolves into [32]

$$\begin{aligned} |\Psi(t)\rangle &= a_0 |\mathbf{0}\rangle + \sum_{m=1}^N \left( \sum_{n \in \{n_s\}} a_n f_n^m \right) |m\rangle + \sum_{p < q=1}^N \left( \sum_{n < m \in \{n_s\}} a_{nm} f_{nm}^{pq} \right) |pq\rangle \\ &+ \sum_{r < s < t=1}^N \left( \sum_{n < m < p \in \{n_s\}} a_{nmp} f_{nmp}^{rst} \right) |rst\rangle + \sum_{r < s < t < u=1}^N \left( \sum_{n < m < p < q \in \{n_s\}} a_{nmpq} f_{nmpq}^{rstu} \right) |rstu\rangle + \dots, \end{aligned} \quad (26)$$

where  $f_{s_1 s_2 \dots s_n}^{r_1 r_2 \dots r_n} = \langle r_1 r_2 \dots r_n | e^{-i\hat{H}t} | s_1 s_2 \dots s_n \rangle$  is the transition amplitude for  $n$ -excitation states from sites  $s_1 s_2 \dots s_n$  to  $r_1 r_2 \dots r_n$ .

The state of the receiver's qubits at sites  $\{n_r\}$  is generally given by a density matrix, which can be obtained, via a lengthy but straightforward calculation, by tracing out all but the receiver's qubits. Finally, the dynamical map  $A$  is derived from the latter by comparison with equation (8). Below, we give a brief overview of this procedure for the cases of  $n = 1$  and  $n = 2$ , and then move to the three- and four-iQST.

### 4.1. One-QST

For the QST of one qubit, the average fidelity of equation (13) reads

$$\langle F_1 \rangle = \frac{1}{6} (2(A_{00}^{00} + A_{11}^{11}) + A_{00}^{11} + A_{11}^{00} + 2 \Re\{A_{10}^{10}\}). \tag{27}$$

The non-zero map elements entering this expression (see appendix A) are

$$A_{00}^{00} = 1, \quad A_{11}^{11} = |f_1^N|^2, \quad A_{10}^{10} = (f_1^N)^*, \quad A_{00}^{11} = 1 - |f_1^N|^2, \tag{28}$$

where  $f_1^N = \langle N | e^{-i\hat{H}t} | 1 \rangle$  is the transition amplitude of one-spin excitation to travel from the sender location 1 to the receiver location  $N$ . With these map elements, equation (27) becomes

$$\langle F_1 \rangle = \frac{1}{2} + \frac{|f_1^N|^2}{6} + \frac{\Re\{f_1^N\}}{3} = \frac{1}{2} + \frac{|f_1^N|^2}{6} + \frac{|f_1^N| \cos \phi}{3}, \tag{29}$$

which coincides with the result in reference [4], and has been used also before, in section 3, when analysing transmission along parallel channels. In the previous expression,  $\phi$  is the argument of the complex number  $f_1^N$ . In order to maximise  $\langle F_1 \rangle$ , one chooses to perform a rotation on the receiver (or to apply a magnetic field to the quantum channel) such that  $\cos \phi = 1$ , so that only the modulus of the transition amplitude matters.

### 4.2. Two-QST

The transfer of a (possibly entangled) two-qubit state along a spin chain has been analyzed, e.g. in reference [32]. There, it was shown that the receiver’s state and the fidelity can be written in terms of one- and two-spin excitation transfer amplitudes. The map allowing to obtain the output (receiver) state from the input (sender) one, is reported in equation (B.1) in appendix B. Using these results, a lengthy but straightforward calculation gives for the average fidelity of equation (13)

$$\begin{aligned} \langle F_2 \rangle = & \frac{1}{4} + \frac{1}{20} (|f_1^N|^2 + |f_2^{N-1}|^2 + |f_{12}^{NN-1}|^2) \\ & + \frac{1}{10} \Re\{f_1^N + f_2^{N-1} + f_{12}^{NN-1} + f_2^{N-1} (f_1^N)^* + f_{12}^{NN-1} (f_1^N)^* + f_{12}^{NN-1} (f_2^{N-1})^*\}, \end{aligned} \tag{30}$$

which simplifies the expression already obtained in reference [33], and has a straightforward physical interpretation: to achieve unit fidelity, all of the excitations initially located on sender sites  $i$  need to reach their mirror-symmetric sites  $N + 1 - i$  located in the receiver’s block, at the same time, with unit transition amplitude. Notice that a similar map has been used in reference [54] to investigate the non-Markovian dynamics of two qubits coupled to spin environments.

Let us stress here that, at variance with the one-QST in section 4.1, we cannot, in general, turn the expression contained in the last term into a sum of transition amplitude moduli with a common phase, as the different transition amplitudes will have, in general, different arguments. The same will be true for the average fidelity of every  $n > 1$ -QST protocol.

### 4.3. n-iQST

The  $n$ -iQST average fidelity  $\langle F_n \rangle$  can be derived, for a quantum map given by  $U(1)$ -symmetric Hamiltonians where the receiver and the channel are initially fully polarized, by a procedure similar to that outlined in section 4.2. The  $n$ -qubit dynamical map (which we do not report for the sake of brevity) is derived in terms of  $p \rightarrow q$  qubit transition amplitudes, with  $p, q \in [1, n]$ . A straightforward calculation yields

$$\langle F_n \rangle = \frac{1}{d} + \frac{1}{d(d+1)} \sum_S |f_S^S|^2 + \frac{2}{d(d+1)} \Re \left\{ \frac{1}{2} \sum_S f_S^S \left( 1 + \sum_{S' \neq S} f_{S'}^{S'} \right)^* \right\}. \tag{31}$$

We recognise equation (31) to have the same structure as  $\langle F_2 \rangle$  in equation (30), with the classical contribution given by the sum of all transition probabilities between the mirror-symmetric partitions of the sender and receiver blocks, and the quantum part given by the product of all of the  $n$ -qubit transition amplitudes times the (complex conjugate) of all of the  $m$ -qubit transition amplitudes between the  $S$  and  $R$ , with  $m < n$ . Before analysing this expression for the particular cases of  $n = 3$  and  $n = 4$ , some observations are in order: the first term in equation (31) corresponds to a random guess  $\langle F_n \rangle = \frac{1}{d}$ ; on the other hand, the summation in the second term runs over all possible transitions  $i \in S \rightarrow i' \in R$  with  $i = i'$ , achieving a total number of  $\sum_{r=1}^n \binom{n}{r} = 2^n - 1$ . The sum of the first two terms, then, gives the LOCC limit  $\langle F_n \rangle = \frac{2}{d+1}$  if all

transition probabilities are equal to unity. As mentioned before, the third terms contains the product of all the transition amplitudes that build up the coherence in the output state, and for this reason we consider this to be a purely quantum contribution.

Due to mirror-symmetry, we can employ a smarter notation in which  $S = \{(i), (ij), (ijk), (ijkl), \dots\}$  are the labels of the sender sites (for the different  $n$  values) and in which we label the receiver site  $r = N + 1 - s$  by the same numbering,  $s$ , as its mirror-symmetric counterpart on the sender block. With this convention, equation (31) can be rewritten in a more compact form as

$$\langle F_n \rangle = \frac{1}{d+1} + \frac{1}{d(d+1)} \left| 1 + \sum_S f_S^S \right|^2. \quad (32)$$

In the following we will evaluate the  $n$ -iQST for three and four qubits for an  $XX$  Hamiltonian. The fidelity and the variance is reported with respect to two possible scenarios: (a.) the sender qubits are let to evolve for a time  $t^*$ , yielding the state  $|\psi_s(t^*)\rangle$ , when the sender and the receiver block, still made out of interacting qubits, are coupled to the wire and (b.) one loads an arbitrary state onto the sender qubits (on a time-scale shorter than the typical evolution of the full Hamiltonian) while they are connected to the wire + receiver system. In scenario (a.) the fidelity is evaluated with respect to the state at the time  $t^*$ , i.e., when the sender (receiver) qubits are connected to the wire; in the scenario (b.) the fidelity is reported with respect to the state that is loaded onto the sender qubits.

## 5. Efficient three- and four-iQST in the $XX$ spin- $\frac{1}{2}$ model with weak links

Here, we propose a high-quality three- and four-iQST protocol by means of an integrable  $U(1)$ -symmetric Hamiltonian (see equation (24)) based on a weak-coupling protocol between the sender (receiver) blocks and the wire. We consider a 1D spin- $\frac{1}{2}$  chain with isotropic interactions in the  $XY$  plane

$$\hat{H} = \frac{1}{4} \sum_i^N J_i (\hat{\sigma}_i^x \hat{\sigma}_{i+1}^x + \hat{\sigma}_i^y \hat{\sigma}_{i+1}^y) + \frac{h_i}{2} \hat{\sigma}_i^z, \quad (33)$$

where  $\hat{\sigma}_i^\alpha$  ( $\alpha = x, y, z$ ) is the Pauli operator sitting on site  $i$ , and we assume open boundary conditions  $\hat{\sigma}_{N+1}^\alpha = 0$ . In the following, we will also assume that the couplings  $J_i$  are all uniform, except for the couplings  $J_i = J_0$  between the sender (receiver) block and the wire (see figure 5). This is the so-called weak-coupling scheme which has been already successfully investigated for one- and two-qubit QST [30, 31, 33, 55]. We will also set the coupling within the sender (receiver) block and within the wire as our time and energy unit  $J_i = J = 1$ . Note that these assumptions are unnecessary for the diagonalisation of the model we are going to outline.

Using the Jordan–Wigner transformation, equation (33) is mapped to a spinless quadratic fermion model [56],

$$\hat{H} = \sum_i^N \frac{J_i}{2} (\hat{c}_i^\dagger \hat{c}_{i+1} + \text{h.c.}) + \sum_i^N h_i \hat{c}_i^\dagger \hat{c}_i - \sum_i^N \frac{h_i}{2}, \quad (34)$$

where, hereafter, the energy is rescaled by the constant term.

The  $U(1)$  symmetry of the model implies that the number operator,  $\hat{N} = \sum_{i=1}^N \hat{c}_i^\dagger \hat{c}_i$  is a conserved quantity. This allows the dynamics to be addressed in excitation-number invariant subspaces. Moreover, due to the quadratic, i.e., non-interacting, nature of the Hamiltonian, the  $n$ -qubit dynamics can be expressed in terms of the single-particle transition amplitudes. In the single-particle sector, equation (34) is diagonalised as

$$\hat{H} = \sum_{k=1}^N \omega_k |\phi_k\rangle \langle \phi_k| \equiv \sum_{k=1}^N \omega_k \hat{c}_k^\dagger \hat{c}_k, \quad (35)$$

where  $\{\omega_k, |\phi_k\rangle\}$ , with  $|\phi_k\rangle = \hat{c}_k^\dagger |0\rangle$ , are the eigenvalues and the eigenvectors of the tridiagonal matrix,  $A \equiv \langle i | \hat{H} | j \rangle = \frac{J}{2} (\delta_{i,j+1} + \delta_{i,j-1}) + h_i \delta_{i,j}$ , describing the single-particle dynamics in the direct space basis,  $|i\rangle \equiv \hat{c}_i^\dagger |0\rangle$ .

Finally, the single-particle transition amplitude from site  $i$  to site  $j$  reads

$$f_i^j(t) = \langle j | e^{-it\hat{H}} | i \rangle = \sum_{k=1}^N e^{-i\omega_k t} \langle j | \phi_k \rangle \langle \phi_k | i \rangle = \sum_{k=1}^N e^{-i\omega_k t} \phi_{jk} \phi_{ki}, \quad (36)$$

and builds up the transition amplitude Hermitian matrix

$$\mathcal{F}(t) = \begin{pmatrix} f_1^1(t) & f_1^2(t) & \dots & f_1^N(t) \\ f_2^1(t) & f_2^2(t) & \dots & f_2^N(t) \\ \vdots & \vdots & \ddots & \vdots \\ f_N^1(t) & f_N^2(t) & \dots & f_N^N(t) \end{pmatrix}. \quad (37)$$

The transition amplitude for the transfer of  $n_s$  excitations, residing on the sender sites  $\{n_s\} = \{s_1, s_2, \dots, s_{n_s}\}$ , to the receiver sites  $r$ , residing on the receiver sites  $\{n_r\} = \{r_1, r_2, \dots, r_{n_r}\}$ , is the minor  $\mathcal{F}(t)_{\{n_s\}}^{\{n_r\}}$  of  $\mathcal{F}(t)$ , i.e., the determinant of the matrix where only the  $\{n_s\}$  rows and  $\{n_r\}$  columns of  $\mathcal{F}(t)$  are considered. The minor  $\{n_r\} = \{r_1, r_2, \dots, r_{n_r}\}$  is the quantity of interest entering equation (32) where it is represented by  $f_S^{\{n_r\}}$ .

Let us also recall that, whereas the  $n$ -iQST formalism developed in sections 2 and 4 is valid, respectively, for arbitrary Hamiltonians and for  $U(1)$ -symmetric ones, the results outlined in this section are valid only for quadratic,  $U(1)$ -symmetric Hamiltonians. Relaxing either, or both, of these constraints significantly increases the complexity of the problem already for one-QST [57]. E.g., including in the Hamiltonian of equation (33) the ZZ-interaction, does not allow the  $n$ -body transition amplitude to be expressed as a determinant of single-particle transition amplitudes as in equation (36), due to the interacting nature of the XXZ Hamiltonian. As a consequence, the scattering events of excitations will likely degrade the  $n$ -iQST efficiency of the channel. Similarly, relaxing the  $U(1)$ -symmetry allows the dynamics to access different particle-number subspaces than those given by the initial conditions, which will similarly introduce a detrimental effect on the  $n$ -iQST efficiency.

### 5.1. Three-iQST via weak links

From the previous discussion, we derive that, in order to have  $\langle F_3 \rangle = 1$ , each of the transition amplitudes  $f_i^j$  belonging to distinct sets of equation (32), i.e., those belonging to a different row and column of equation (37), need to have unit modulus and the same phase.


Without loss of generality, we choose  $i - j$  (or, equivalently,  $i + j$ ) to be even, so that  $f_i^j$  is purely real (imaginary) by choosing the transition amplitudes between sites  $(1, N - 2)$ ,  $(2, N - 1)$ ,  $(3, N)$  for  $N$  odd (even). From reference [58], we know that for a length of the wire  $n_w = 4l + 1$  and  $n_w = 4l + 3$  there are, respectively, one and three resonant single-particle levels with the sender (receiver) block. We will refer to these two cases as non-resonant and resonant three-iQST respectively, because, in the former case, the QST time is ruled by the non-resonant energy level splitting and in the latter by the resonant energy level splitting. For the non-resonant case, i.e., for length of the wire  $n_w = 4l + 1$ , each single-particle transition amplitude in equation (36) can be approximated, up to 2nd-order, by

$$f_i^j(t) = \sum_{k=1}^7 e^{-i\omega_k t} \phi_{jk} \phi_{ki}, \quad (38)$$

where only the quasi-degenerate energy levels enter the sum, and occupy, in the increasing ordered energy spectrum  $\omega_k < \omega_{k'}$  for  $k < k'$ , the following positions: the four 2nd-order perturbed energy levels are at  $\{\lceil \frac{N-5}{4} \rceil, \lceil \frac{N-5}{4} \rceil + 1, \lceil \frac{N-5}{4} \rceil + n_w, \lceil \frac{N-5}{4} \rceil + n_w + 1\}$ , where  $\lceil x \rceil$  is the ceiling function, and the three 1st-order perturbed energy levels are at  $\{\frac{N+1}{2} - 1, \frac{N+1}{2}, \frac{N+1}{2} + 1\}$ . In figure 6, an instance of the single-particle energy levels is given for the non-resonant case with  $n_w = 9$ . Exploiting the parity relations for the eigenvectors of mirror-symmetric matrices [59],  $\phi_{k, N+1-i} = (-1)^k \phi_{k, N+1-i}$ , and elementary trigonometric identities, it is easy to show that the longest time-scale is governed by the 2nd-order perturbative energy splitting. As a consequence, the envelope of the  $n$ -qubit QST average fidelity is given by  $\langle F \rangle_{\text{env}} \simeq \left| \sin^2 \left( \frac{\delta\omega}{2} t \right) \right|^2$ , where  $\delta\omega = \omega_{\lceil \frac{N-5}{4} \rceil + 1} - \omega_{\lceil \frac{N-5}{4} \rceil}$ . Within the transfer time  $\tau = \frac{\pi}{\delta\omega}$ , oscillations on a timescale of order of  $J$  occur because of the internal dynamics of the receiver block. Nevertheless, the fidelity reaches its maximum value of  $\langle F \rangle = 1 - O(J_0^2)$  multiple times, giving the receiver the opportunity to read-out the state within a time-window of the order of  $J$ . In figure 6 we show an instance of the aforementioned timescales.

### 5.2. Four-iQST via weak links

Here we turn our attention to the QST of four interacting qubits, again over the channel depicted in figure 5, and with the same Hamiltonian as in equation (33), where, now, the weak-coupling condition entails  $J_i = J_0(\delta_{i,4} + \delta_{i, N-4})$  and we set  $J_i = 1$  otherwise. However, it has been shown in reference [58] that,



for such a uniform coupling scheme, either all, or none, of the four single-particle energy levels are in resonance with the wire's energy level. As a consequence, because of the incommensurability of the frequencies entering equation (38) (where now the summations extends over 12 or eight frequencies for the all-resonant and the non-resonant cases, respectively), the average fidelity of the four-iQST will not approach unity. In order to introduce a time-scale separation, as done in the three-iQST case, a minimal-engineering solution can be achieved by acting on the intra-sender (-receiver) couplings  $J_i$  such that two (symmetric) energy levels are in resonance with the wire's energies and two are left out-of-resonance. The values for  $J_s$  can be readily found by setting



$$J_s = \frac{\cos \frac{k\pi}{n_w+1}}{\cos \frac{s\pi}{5}}, \quad (39)$$

where  $s \in (1, 2)$  represents the sth-energy level of the sender put in resonance with the kth-energy level of the wire for that value of  $J_s$ . In figure 7 a schematic representation is given for  $n_w = 10$ ,  $k = 2$ , and  $s = 1$ . With such a protocol, we achieve a separation of time-scales in the single-particle transition amplitudes, and achieve, once again, a transfer time of the order of magnitude of the 2nd-order energy perturbation correction, represented in figure 7 by the sinusoidal envelop of  $\bar{F}_4$ . Without loss of generality, we report in figure 7 the single-particle transition amplitudes  $f_1^1 = f_4^4 \simeq f_2^2 = f_3^3 > 0.99$ , resulting in  $\langle F_4 \rangle \simeq 0.98$  at the optimal transfer time.

### 5.3. Comparison between $n$ -QST and $n$ -iQST

In sections 3 and 5, we showed that a single interacting channel is able to efficiently transfer the quantum state of  $n$  spins, with  $n$  independent (parallel) transmissions if they are non-interacting, and in a single transfer instance, if we are in the presence of interactions between the sender's qubits, up to  $n = 4$ . Here, we address the question: which of these two type of transfer performs more reliably at a fixed value of given average fidelity? We do this by using the variance in equation (19) as a figure of merit.


As can be seen from figure 8, at fixed average fidelity  $\langle F \rangle$ , the variance  $\Delta F^2$  is always greater for the case of  $n$ -iQST with a single channel, than for  $n$ -QST with  $n$  independent channels. This can be readily



explained by noticing that the presence of interactions allows the receivers to explore a higher portion of the Hilbert space during the evolution, with respect to the case of independent channels. E.g., in the case of independent channels, the entanglement of the receivers cannot exceed that of the senders, whereas this is not the case for the single channel scenario.

#### 5.4. Effect of imperfections and disorder

For experimentally feasible implementations of the proposed  $n$ -iQST protocol, one needs to take into account that both imperfections in tuning the weak coupling  $J_0$  to specific values and the presence of



disorder in the wire may affect detrimentally the efficiency of the transfer. Both type of deviations from ideal protocols have been largely investigated for one-QST, see, e.g., references [30, 60–64] where both imperfections and the disorder is mainly considered to be static, i.e., values deviating from the ideal (weak)-coupling and random distributions of spin–spin couplings and magnetic fields in the wire, respectively. We perform a similar analysis for the  $n$ -iQST average fidelity by allowing the sender (receiver) block to be coupled to the wire via values of  $J_0$  different from those reported in figures 6 and 7 and we consider the case of disorder, independently for  $h$  and  $J$ , in the wire with a uniform distribution in the interval  $[-\Delta h, \Delta h]$  and  $[J - \Delta J, J + \Delta h]$ , respectively.

Figure 9 shows that the three-iQST fidelity is more resilient in the presence of spin–spin coupling disorder than in the presence of magnetic field disorder. However, in both cases, an average fidelity higher than the LOCC limit is obtained also in the presence of strong disorder of order  $\mathcal{O}(1)$ , whereas, for ultrastrong disorder the random guess scenario is realised. Clearly, being the weak-coupling protocol based on the resonance of a sender (receiver) single-particle energy level with one of the wire, disorder starts to significantly affect the fidelity only once the energy level of the wire is brought out of resonance, which occurs earlier for magnetic disorder.

Similarly, for values of  $J_0$  different than those used in sections 5.1 and 5.2, the quality of the three- and four-iQST remains well above the LOCC limit also for values of  $J_0$  up to 0.1, whereas, for values approaching unit and above, the random guess scenario is approached, as can be seen in the middle and right panel of figure 9.

## 6. Concluding remarks

Quantum state transfer of an  $n$ -qubit system is a key protocol in many QIP tasks. Whereas single qubit quantum state transfer has been intensively investigated, and also experimentally realised on a variety of different experimental platforms, arbitrary  $n$ -qubit quantum state transfer is still a goal to be achieved. The quantum transfer of a many-body interacting system is a formidable task, made difficult both by the exponentially-increasing dimensionality of the Hilbert space, and by the complexity due to the particle interactions. In this paper, we have provided a new approach to the  $n$  interacting qubit QST via dynamical maps. By considering the receiver block as an open quantum system coupled to an environment, embodied both by the sender block and the quantum channel, we have derived a general expression for the average fidelity of  $n$ -iQST in terms of quantum dynamical maps elements. We have also analyzed the dispersion of the values of the fidelity, when evaluated on all possible pure input states, that we expressed by the fidelity variance, for which we provided a general expression as well. Then, we specialised to short-range transmission obtained by coupling senders and receivers to a linear spin chain. In the case of  $U(1)$ -symmetric dynamics, we expressed the average fidelity in terms of transition amplitudes in the occupation-number invariant subspaces. We investigated in detail the  $U(1)$ -symmetric  $XX$  spin- $\frac{1}{2}$  Hamiltonian, and, by exploiting its non-interacting nature in the fermionic representation, we were able to express the  $n$ -qubit iQST average fidelity only in terms of single-particle transition amplitudes.

Our formalism also encompasses the non-interacting scenario, where we have shown that independent channels achieve a higher  $n$ -QST fidelity, at a fixed single-particle transition amplitude, for product states than for entangled states, although the variance of the former is greater than that of the latter, at a fixed value of the average fidelity. Interestingly, we obtained that the average fidelity is a self-averaging quantity, as quantified by the vanishing coefficient of variation for  $n \gg 1$  at high-values of the average fidelity.

Finally, we have proposed a protocol for the high-fidelity transfer of the quantum state of both three and four interacting qubits, arranged in a linear chain with uniform couplings, via a weak-coupling scheme to a non-engineered  $XX$  spin- $\frac{1}{2}$  chain. Whereas up to  $n = 4$ , high-quality iQST can be achieved by means of a uniform channel, it appears that for  $n > 4$  our protocol has to be substantially modified because of the impossibility to introduce a time-scale separation in the single-particle transition amplitudes when the involved energy levels become too numerous.

Considering the importance of quantum state transfer of a many-qubit system in several QIP tasks, ranging from cryptography to quantum computation, it is crucial to establish a theoretical framework that can encompass all of the various possibilities. We believe that this can be provided by the quantum map approach supplemented by the investigation of the statistics of the fidelity, that we started to establish in this paper, and that we applied to the general  $n$ -qubit state transfer. We provided analytical expressions of the 1st and 2nd moment of the  $n$ -iQST PDF and random matrix theory may be used to derive all moments of the QST PDE. Knowledge of the full PDF is relevant not only from a quantum information theoretical perspective but also in scenarios where single-shot QST is implemented. We will leave this topic as future works as the PDF is highly dependent on the chosen QST protocol. Moreover, our approach explicitly includes interactions among the sender's particles, opening the way to investigate quantum transfer protocols of complex interacting systems.

## Acknowledgments

TJGA thanks Abolfazl Bayat for pointing him out to reference [29], where useful expressions for the  $d$ -level fidelity can be found. SL acknowledges support from MIUR through project PRIN Project 2017SRN-BRK QUSHIP. TJGA acknowledges funding through the IPAS+ (Internationalisation Partnership Awards Scheme +) ACROSS project by the MCST (The Malta Council for Science & Technology). KŻ acknowledges support by Foundation for Polish Science through TEAM-NET Project under Contract POIR.04.04.00-00-17C1/18-00. We also thanks the organizers of the Quantum Hiking Conference in 2019 where useful discussions concerning this work have taken place.

## Data availability statement

The data that support the findings of this study are available upon reasonable request from the authors.

## Appendix A. One-qubit map

The one-qubit density matrix map  $\hat{\rho}_N = \Phi(t)\hat{\rho}_1$  is given by

$$\begin{pmatrix} \rho_{00} \\ \rho_{01} \\ \rho_{10} \\ \rho_{11} \end{pmatrix}_N = \begin{pmatrix} 1 & 0 & 0 & 1 - |f_1^N|^2 \\ 0 & f_1^N & 0 & 0 \\ 0 & 0 & (f_1^N)^* & 0 \\ 0 & 0 & 0 & |f_1^N|^2 \end{pmatrix} \begin{pmatrix} \rho_{00} \\ \rho_{01} \\ \rho_{10} \\ \rho_{11} \end{pmatrix}_1. \quad (\text{A.1})$$

## Appendix B. Two-qubit map

From reference [32], we derive the following two-qubit map's elements  $\hat{\rho}_{N-1,N} = \Phi(t)\hat{\rho}_{1,2}$ ,



$$\begin{aligned}
 A_{00}^{00} &= 1, & A_{00}^{11} &= 1 - |f_1^{N-1}|^2 - |f_1^N|^2, & A_{00}^{22} &= 1 - |f_2^{N-1}|^2 - |f_2^N|^2, \\
 A_{00}^{33} &= 1 - |f_{12}^{mN-1}|^2 - |f_{12}^{mN}|^2 - |f_{12}^{N-1N}|^2, \\
 A_{00}^{12} &= -f_1^{N-1} (f_2^{N-1})^* - f_1^N (f_2^N)^*, & A_{00}^{21} &= -f_2^{N-1} (f_1^{N-1})^* - f_2^N (f_1^N)^*, \\
 A_{01}^{01} &= (f_1^N)^*, & A_{01}^{02} &= (f_2^N)^*, & A_{01}^{13} &= f_1^m (f_{12}^{mN})^*, & A_{01}^{23} &= f_2^m (f_{12}^{mN})^*, \\
 A_{02}^{01} &= (f_1^{N-1})^*, & A_{02}^{02} &= (f_2^{N-1})^*, & A_{02}^{13} &= f_1^m (f_{12}^{mN-1})^*, & A_{02}^{23} &= f_2^m (f_{12}^{mN-1})^* \\
 A_{03}^{03} &= (f_{12}^{N-1N})^*, \\
 A_{11}^{11} &= |f_1^N|^2, & A_{11}^{12} &= f_1^N (f_2^N)^*, & A_{11}^{21} &= f_2^N (f_1^N)^*, & A_{11}^{22} &= |f_2^N|^2, & A_{11}^{33} &= |f_{12}^{mN}|^2, \\
 A_{12}^{11} &= f_1^N (f_1^{N-1})^*, & A_{12}^{12} &= f_1^N (f_2^{N-1})^*, & A_{12}^{21} &= f_2^N (f_1^{N-1})^*, & A_{12}^{22} &= f_2^N (f_2^{N-1})^*, \\
 A_{12}^{33} &= f_{12}^{mN} (f_{12}^{mN-1})^*, \\
 A_{13}^{13} &= f_1^N (f_{12}^{N-1N})^*, & A_{13}^{23} &= f_2^N (f_{12}^{N-1N})^*, \\
 A_{22}^{33} &= |f_{12}^{mN-1}|^2, & A_{22}^{11} &= |f_1^{N-1}|^2, & A_{22}^{22} &= |f_2^{N-1}|^2, & A_{22}^{12} &= f_1^{N-1} (f_2^{N-1})^*, \\
 A_{22}^{21} &= f_2^{N-1} (f_1^{N-1})^*, \\
 A_{23}^{13} &= f_1^{N-1} (f_{12}^{N-1N})^*, & A_{23}^{23} &= f_2^{N-1} (f_{12}^{N-1N})^*, & A_{33}^{33} &= |f_{12}^{N-1N}|^2,
 \end{aligned} \tag{B.1}$$

where  $m$  denotes the summation over all  $i \neq S, R$

$$\begin{pmatrix} \rho_{00} \\ \rho_{01} \\ \rho_{02} \\ \rho_{03} \\ \rho_{10} \\ \rho_{11} \\ \rho_{12} \\ \rho_{13} \\ \rho_{20} \\ \rho_{21} \\ \rho_{22} \\ \rho_{23} \\ \rho_{30} \\ \rho_{31} \\ \rho_{32} \\ \rho_{33} \end{pmatrix}_{N-1,N} = \begin{pmatrix} A_{00}^{00} & 0 & 0 & 0 & 0 & A_{00}^{11} & A_{00}^{12} & 0 & 0 & A_{00}^{21} & A_{00}^{22} & 0 & 0 & 0 & 0 & A_{00}^{33} \\ 0 & A_{01}^{01} & A_{01}^{02} & 0 & 0 & 0 & 0 & A_{01}^{13} & 0 & 0 & 0 & A_{01}^{23} & 0 & 0 & 0 & 0 \\ 0 & A_{02}^{01} & A_{02}^{02} & 0 & 0 & 0 & 0 & A_{02}^{13} & 0 & 0 & 0 & A_{02}^{23} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & A_{03}^{03} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & A_{10}^{10} & 0 & 0 & 0 & A_{10}^{20} & 0 & 0 & 0 & 0 & A_{10}^{31} & A_{10}^{32} & 0 \\ 0 & 0 & 0 & 0 & 0 & A_{11}^{11} & A_{11}^{12} & 0 & 0 & A_{11}^{21} & A_{11}^{22} & 0 & 0 & 0 & 0 & A_{11}^{33} \\ 0 & 0 & 0 & 0 & 0 & A_{12}^{11} & A_{12}^{12} & 0 & 0 & A_{12}^{21} & A_{12}^{22} & 0 & 0 & 0 & 0 & A_{12}^{33} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & A_{13}^{13} & 0 & 0 & 0 & A_{13}^{23} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & A_{20}^{10} & 0 & 0 & 0 & A_{20}^{20} & 0 & 0 & 0 & 0 & A_{20}^{31} & A_{20}^{32} & 0 \\ 0 & 0 & 0 & 0 & 0 & A_{21}^{11} & A_{21}^{12} & 0 & 0 & A_{21}^{21} & A_{21}^{22} & 0 & 0 & 0 & 0 & A_{21}^{33} \\ 0 & 0 & 0 & 0 & 0 & A_{22}^{11} & A_{22}^{12} & 0 & 0 & A_{22}^{21} & A_{22}^{22} & 0 & 0 & 0 & 0 & A_{22}^{33} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & A_{23}^{13} & 0 & 0 & 0 & A_{23}^{23} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & A_{30}^{30} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & A_{31}^{31} & A_{31}^{32} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & A_{32}^{31} & A_{32}^{32} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & A_{33}^{33} \end{pmatrix} \begin{pmatrix} \rho_{00} \\ \rho_{01} \\ \rho_{02} \\ \rho_{03} \\ \rho_{10} \\ \rho_{11} \\ \rho_{12} \\ \rho_{13} \\ \rho_{20} \\ \rho_{21} \\ \rho_{22} \\ \rho_{23} \\ \rho_{30} \\ \rho_{31} \\ \rho_{32} \\ \rho_{33} \end{pmatrix}_{1,2}$$

### ORCID iDs

Tony J G Apollaro  <https://orcid.org/0000-0002-9324-9336>

Salvatore Lorenzo  <https://orcid.org/0000-0002-0827-5549>

Mirko Consiglio  <https://orcid.org/0000-0001-8730-0206>

### References

- [1] Nielsen M A and Chuang I L 2010 *Quantum Computation and Quantum Information* (Cambridge: Cambridge University Press)
- [2] Northup T E and Blatt R 2014 Quantum information transfer using photons *Nat. Photon.* **8** 356–63
- [3] Bennett C H, Brassard G, Crépeau C, Jozsa R, Peres A and Wootters W K 1993 Teleporting an unknown quantum state via dual classical and Einstein–Podolsky–Rosen channels *Phys. Rev. Lett.* **70** 1895–9
- [4] Bose S 2003 Quantum communication through an unmodulated spin chain *Phys. Rev. Lett.* **91** 207901
- [5] Tamascelli D, Olivares S, Rossotti S, Osellame R and Paris M G A 2016 Quantum state transfer via Bloch oscillations *Sci. Rep.* **6** 26054
- [6] Chapman R J, Santandrea M, Huang Z, Corrielli G, Crespi A, Yung M-H, Osellame R and Peruzzo A 2016 Experimental perfect state transfer of an entangled photonic qubit *Nat. Commun.* **7** 11339
- [7] Vogell B, Vermersch B, Northup T E, Lanyon B P and Muschik C A 2017 Deterministic quantum state transfer between remote qubits in cavities *Quantum Sci. Technol.* **2** 045003
- [8] Li X et al 2018 Perfect quantum state transfer in a superconducting qubit chain with parametrically tunable couplings *Phys. Rev. Appl.* **10** 054009
- [9] Ali H, Basit A, Badshah F and Ge G-Q 2018 Quantum state transfer between nitrogen vacancy centers coupled to photonic crystal molecule in the off resonant regime *Physica E* **104** 261–7

- [10] Dlaska C, Vermersch B and Zoller P 2017 Robust quantum state transfer via topologically protected edge channels in dipolar arrays *Quantum Sci. Technol.* **2** 015001
- [11] Vieira R and Rigolin G 2019 Robust and efficient transport of two-qubit entanglement via disordered spin chains *Quantum Inf. Process.* **18** 135
- [12] Vieira R and Rigolin G 2020 Almost perfect transmission of multipartite entanglement through disordered and noisy spin chains *Phys. Lett. A* **384** 126536
- [13] Yousefjani R and Bayat A 2021 Parallel entangling gate operations and two-way quantum communication in spin chains *Quantum* **5** 460
- [14] Yousefjani R and Bayat A 2020 Simultaneous multiple-user quantum communication across a spin-chain channel *Phys. Rev. A* **102** 012418
- [15] Preskill J 2018 Quantum computing in the NISQ era and beyond *Quantum* **2** 79
- [16] Ouyang Y, Tan S H, Zhao L and Fitzsimons J F 2017 Computing on quantum shared secrets *Phys. Rev. A* **96** 052333
- [17] Wehner S, Elkouss D and Hanson R 2018 Quantum internet: a vision for the road ahead *Science* **362** eaam9288
- [18] Bengtsson I and Życzkowski K 2017 *Geometry of Quantum States* (Cambridge: Cambridge University Press)
- [19] Anderson P W 1972 More is different *Science* **177** 393–6
- [20] Albanese C, Christandl M, Datta N and Ekert A 2004 Mirror inversion of quantum states in linear registers *Phys. Rev. Lett.* **93** 230502
- [21] Kay A 2010 Perfect, efficient, state transfer and its application as a constructive tool *Int. J. Quantum Inf.* **8** 641–76
- [22] Sousa R and Omar Y 2014 Pretty good state transfer of entangled states through quantum spin chains *New J. Phys.* **16** 123003
- [23] Lorenz P and Stolze J 2014 Transferring entangled states through spin chains by boundary-state multiplets *Phys. Rev. A* **90** 044301
- [24] Apollaro T J G, Sanavio C, Chetcuti W J and Lorenzo S 2020 Multipartite entanglement transfer in spin chains *Phys. Lett. A* **384** 126306
- [25] Vieira R and Rigolin G 2018 Almost perfect transport of an entangled two-qubit state through a spin chain *Phys. Lett. A* **382** 2586–94
- [26] Almeida G M A, Souza A M C, de Moura F A B F and Lyra M L 2019 Robust entanglement transfer through a disordered qubit ladder *Phys. Lett. A* **383** 125847
- [27] Horodecki M, Horodecki P and Horodecki R 1999 General teleportation channel, singlet fraction, and quasidistillation *Phys. Rev. A* **60** 1888–98
- [28] Aubert S and Lam C S 2004 Invariant and group theoretical integrations over the  $U(n)$  group *J. Math. Phys.* **45** 3019–39
- [29] Bayat A and Karimipour V 2006 Transfer of  $d$ -level quantum states through spin chains by random swapping *Phys. Rev. A* **75** 022321
- [30] Wójcik A, Łuczak T, Kurzyński P, Grudka A, Gdala T and Bednarska M 2005 Unmodulated spin chains as universal quantum wires *Phys. Rev. A* **72** 034303
- [31] Lorenzo S, Apollaro T J G, Trombetti A, Paganelli S and Paganelli S 2017 Two-qubit quantum state transfer in spin chains and cold atoms with weak links *Int. J. Quantum Inf.* **15** 1750037
- [32] Apollaro T J G, Lorenzo S, Sindona A, Paganelli S, Giorgi G L and Plastina F 2015 Many-qubit quantum state transfer via spin chains *Phys. Scr.* **T165** 014036
- [33] Lorenzo S, Apollaro T J G, Paganelli S, Palma G M and Plastina F 2015 Transfer of arbitrary two-qubit states via a spin chain *Phys. Rev. A* **91** 042321
- [34] Jozsa R 1994 Fidelity for mixed quantum states *J. Mod. Opt.* **41** 2315–23
- [35] Bagan E, Baig M and Muñoz-Tapia R 2003 Minimal measurements of the gate fidelity of a qudit map *Phys. Rev. A* **67** 014303
- [36] Bayat A and Karimipour V 2007 Transfer of  $d$ -level quantum states through spin chains by random swapping *Phys. Rev. A* **75** 022321
- [37] Cabrera R and Baylis W E 2007 Average fidelity in  $n$ -qubit systems *Phys. Lett. A* **368** 25–8
- [38] Liang Y C, Yeh Y H, Mendonça P E M F, Teh R Y, Reid M D and Drummond P D 2018 Quantum fidelity measures for mixed states (arXiv:1810.08034)
- [39] Lorenzo S, Plastina F, Consiglio M and Apollaro T J G 2021 Quantum map approach to entanglement transfer and generation in spin chains.
- [40] Collins B and Śniady P 2006 Integration with respect to the Haar measure on unitary, orthogonal and symplectic group *Commun. Math. Phys.* **264** 773–95
- [41] Nielsen M A 2002 A simple formula for the average gate fidelity of a quantum dynamical operation *Phys. Lett. A* **303** 249–52
- [42] Johnston N and Kribs D W 2011 Quantum gate fidelity in terms of Choi matrices *J. Phys. A: Math. Theor.* **44** 495303
- [43] Życzkowski K and Sommers H J 2005 Average fidelity between random quantum states *Phys. Rev. A* **71** 032313
- [44] Pedersen L H, Møller N M and Mølmer K 2008 The distribution of quantum fidelities *Phys. Lett. A* **372** 7028–32
- [45] Pedersen L H, Møller N M and Mølmer K 2007 Fidelity of quantum operations *Phys. Lett. A* **367** 47–51
- [46] Mayer K and Knill E 2018 Quantum process fidelity bounds from sets of input states *Phys. Rev. A* **98** 052326
- [47] Mello P A 1990 Averages on the unitary group and applications to the problem of disordered conductors *J. Phys. A: Math. Gen.* **23** 4061–80
- [48] Karimipour V, Rad M S and Asoudeh M 2012 Perfect quantum state transfer in two- and three-dimensional structures *Phys. Rev. A* **85** 010302
- [49] Bose S 2007 Quantum communication through spin chain dynamics: an introductory overview *Contemp. Phys.* **48** 13–30
- [50] Banchi L, Apollaro T J G, Cuccoli A, Vaia R and Verrucchi P 2011 Long quantum channels for high-quality entanglement transfer *New J. Phys.* **13** 123006
- [51] Hermes S, Apollaro T J G, Paganelli S and Macri T 2020 Dimensionality-enhanced quantum state transfer in long-range-interacting spin systems *Phys. Rev. A* **101** 053607
- [52] Almeida G M A, Ciccarello F, Apollaro T J G and Souza A M C 2015 Quantum-state transfer in staggered coupled-cavity arrays *Phys. Rev. A* **93** 032310
- [53] Pavlis A K, Nikolopoulos G M and Lambropoulos P 2016 Evaluation of the performance of two state-transfer Hamiltonians in the presence of static disorder *Quantum Inf. Process.* **15** 2553–68
- [54] Apollaro T J G, Cuccoli A, Di Franco C, Paternostro M, Plastina F and Verrucchi P 2010 Manipulating and protecting entanglement by means of spin environments *New J. Phys.* **12** 083046
- [55] Apollaro T J G and Chetcuti W J 2020 Two-excitation routing via linear quantum channels *Entropy* **23** 51
- [56] Elliott L, Schultz T and Mattis D 1961 Two soluble models of an antiferromagnetic chain *Ann. Phys., NY* **16** 407–66

- [57] Bayat A and Bose S 2010 Information-transferring ability of the different phases of a finite XXZ spin chain *Phys. Rev. A* **81** 012304
- [58] Chetcuti W J, Sanavio C, Lorenzo S and Apollaro T J G 2020 Perturbative many-body transfer *New J. Phys.* **22** 033030
- [59] Banchi L and Vaia R 2013 Spectral problem for quasi-uniform nearest-neighbor chains *J. Math. Phys.* **54** 043501
- [60] Zwick A and Osenda O 2011 Quantum state transfer in a XX chain with impurities *J. Phys. A: Math. Theor.* **44** 105302
- [61] Lorenzo S, Apollaro T J G, Sindona A and Plastina F 2013 Quantum-state transfer via resonant tunneling through local-field-induced barriers *Phys. Rev. A* **87** 042313
- [62] Zwick A, Álvarez G A, Bensity G and Kurizki G 2014 Optimized dynamical control of state transfer through noisy spin chains *New J. Phys.* **16** 065021
- [63] Almeida G M A, de Moura F A B F and Lyra M L 2018 Quantum-state transfer through long-range correlated disordered channels *Phys. Lett. A* **382** 1335–40
- [64] Keele C and Kay A 2022 Combating the effects of disorder in quantum state transfer *Phys. Rev. A* **105** 032612