

[PRINT](#)[CLOSE WINDOW](#)

FISV 2008 - 10th Annual Congress Riva del Garda, 24-27 September 2008

<http://fisv2008.azuleon.org>

Cellular stress, apoptosis and autophagy

25/09/2008, 08:30 - [---]

(Organisers: [T. Pozzan](#), [M. Piacentini](#))

Abstract UAI: fc6c68ea0630150557

Stress induced and physiological apoptosis during early development of sea urchin *Paracentrotus lividus*

M. Agnello¹, S. Filosto¹, F. Trinchella², M.C. Roccheri¹

¹Dip. Biologia Cellulare e dello Sviluppo "A. Monroy", Univ. Palermo (Palermo); ²Univ. Federico II, Dip.di Scienze Biologiche, Sezione di Biologia Evolutiva e Comparata (Naples)

We investigate the toxic effect of cadmium on *P. lividus* embryo development. We already ascertained that 1mM and 100µM CdCl₂ induce apoptosis and found a competition between Cd and Ca uptake during early development. Using calcein-AM we show that intracellular amount of Ca in Cd-treated embryos is effectively lower than controls. We find cadmium to induce apoptosis through caspase-3 activation, while caspase-7 is the main effector of physiological apoptosis. Caspase-10 results active only in physiological apoptosis, since caspase-8 is predominantly involved in stress-induced apoptosis. Another stress inducer, the TPA, causes activation of all examined caspases. Immunocytochemical experiments with anti-cleaved-caspase-9 antibody, immunoblotting with anti- cytochrome-C antibody and fluorescent assays with JC-1 fluorochrome fail to find a mitochondrial involvement in apoptosis. Moreover, we show a detectable generation of reactive oxygen species, following exposure to Cd. We studied also caspases expressions during early physiological development and demonstrate that caspase-3 and caspase-10 are well-represented only after mesenchyme blastula stage.