

Berseem-annual ryegrass intercropping: effect of plant arrangement and seeding ratio on N₂ fixation and yield

STRINGI, L., GIAMBALVO, D., SCARPELLO, C., ATTARDO, C., FREANDA, A. S. AND TRAPANI, P.

Dipartimento di Agronomia Ambientale e Territoriale, Università di Palermo, Viale delle Scienze, 90128 Palermo, Italy

Abstract

Various agronomic factors can affect the productivity and the efficiency of legume-grass intercropping systems. This research was carried out in a Mediterranean semi-arid environment (37°30'N; 13°31'E; 178 m a.s.l.) with the aim to study on berseem clover- annual ryegrass mixture (*Trifolium alexandrinum* L. – *Lolium multiflorum* Lam subsp. *wersterwoldicum*) the effects of different plant arrangement (sowing of the two components in alternate rows or in the same row) and seeding ratios (100:0; 75:25; 50:50; 25:75; 0:100) on forage yield, nitrogen content and nitrogen fixation. The experimental design was a split-plot with four replications. The ¹⁵N isotope dilution technique was used to estimate nitrogen fixation by berseem clover. All plots were cut four times (first cut 85 DAS; rest period of four weeks). The DM yield of mixtures and pure stand clover were similar; annual ryegrass in pure stand produced the lowest yields. No significant differences for DM yield were observed due to the plant arrangement and seeding ratio in the mixture. Intercropping berseem had always a significant higher percentage of Ndfa than monocropped berseem. The %Ndfa of mixed berseem was not influenced by plant arrangement but gradually decreased when proportion of berseem in the mixture increased.

Keywords: berseem clover, annual ryegrass, mixture, plant arrangement, seeding ratio, nitrogen fixation

Introduction

Intercropping is the growing of two (or more) crops together on the same area of ground and often produces an advantage in terms of more yield and less variation in yield than comparable areas of sole crops (Willey, 1979; Bulson *et al.*, 1997). The most commonly accepted reason explaining why it is possible to obtain better yields with crop mixtures is that the component crops can use more efficiently the limiting resources differing in their growth requirements. Systems which combine grass and legume crops are highly regarded because of their potential for replacing fertilizer N with symbiotically fixed atmospheric N₂ and the economic and environmental benefits which may result from this (Stern, 1993). However, in the literature there are few data available on biological nitrogen fixation by annual forage legumes in Mediterranean environments. As well as few data is available on the contribution of biological nitrogen fixed by annual legume to the nitrogen nutrition of a non-legume grown in mixed stand and on the effects of intercropping with grass on the quantity of N₂ fixed by forage legume component. Moreover the productivity and efficiency of grass-legume intercrop system are affected by various agronomic variables such as component crop density, plant spacing and arrangement, time of sowing of component crop, N fertilization (Ofori and Stern, 1987).

The aim of this research was to evaluate on berseem clover – annual ryegrass mixture the effects of different plant arrangement and seeding ratios on forage yield, nitrogen content and nitrogen fixation in a typical rain-fed Mediterranean environment.

Materials and Methods

The research was carried out during 2002/03 in a hilly area of Sicily (37°30'N; 13°31'E; 178 m a.s.l.) on a *Vertic haploxerepts* soil and with wheat as previous crop. The topsoil (0–40 cm) had the following characteristics: 38% clay, 25% silt, 36% sand, 8.4 pH (in water), 1.27% organic matter, 0.85‰ total nitrogen.

The experiment was set up in a complete randomized block design with four replications. Treatments were: berseem (*Trifolium alexandrinum* L. cv Lilibeo) in pure stand (B); annual ryegrass (*Lolium multiflorum* Lam. subsp. *wersterwoldicum* cv Elunaria) in pure stand (R); and their mixture with different plant arrangement [sowing of the two components in alternate rows (BR_{AR}) or in the same row (BR_{SR})] and seeding ratios (B₂₅-R₇₅; B₅₀-R₅₀; B₇₅-R₂₅). Pure stands were seeded at 50 kg seeds ha⁻¹. Mixed and pure stands were sown by hand on 16 December 2002. No herbicide treatments were used; all plots were kept free of weeds by hand-weeding.

The ¹⁵N isotope dilution technique was used to estimate nitrogen fixation by clover. ¹⁵N fertilizer [(NH₄)₂SO₄ with an isotopic composition of 10 atom% ¹⁵N] was uniformly applied in a liquid form, following the application procedure described by Høgh-Jensen and Schjøerring (1994). ¹⁵N fertilizer (8 kg N ha⁻¹) was applied to a 2.88 m² microplot in the middle of each plot at the emergence of the crops and again two days after the 1st and 2nd utilizations. The rest of the plot outside the ¹⁵N-labelled area received a top-dressing of ammonium sulphate in an amount equivalent to the microplots. After each application of labelled fertilizer an irrigation of all plots was used in order to avoid the retention of ¹⁵N fertilizer on the plant leaves.

All plots were cut simultaneously at 85d after sowing and defoliation frequency of 28d was used (4 times). All plots were harvested by hand at 5 cm stubble height and total fresh weight was determined. At each harvest, a sample of plant material, taken in the centre of microplots, was hand-disaggregated into its botanical components (B, R), dried at 60 °C for 36 h, weighed and ground to a fine powder (sieved using a 0.1 mm mesh size) in a fast running mill and analysed by the Iso-Analytical laboratory (EA-IRMS technique) for ¹⁵N enrichment and total N content.

Data on ¹⁵N enrichment of biomass were used to calculate the percentage of clover nitrogen derived from symbiotic N₂ fixation (%Ndfa) according to Fried and Middleboe (1977).

The GLM procedure of the Statistical Analysis System (SAS) was used to analyze the variance. Least significant difference (LSD at P≤0.05) was used to make comparisons among treatments.

Results and Discussion

During growing season total rainfall was close to the long-term mean (532 mm vs 550 mm) and distribution was generally favourable for crop growth.

Dry matter yield (kg ha⁻¹) for annual ryegrass and berseem clover in pure and mixed stands in relation to the different plant arrangement and seeding ratio are reported in Table 1.

The yield of sole-cropped berseem was significantly higher than ryegrass in pure stand (9449 vs 4826; P<0.0001). The DM yield of mixture and pure stand clover were similar. No significant differences for DM yield were observed due to the arrangement of plants and to the different seeding ratio in the mixture.

The legumes competed very well in the mixed stands, producing on average 80% of the total yield. Both seeding ratio and plant arrangement had small effects on %clover on DM basis.

Annual ryegrass intercropped with berseem consistently had a higher nitrogen concentration in the dry matter than ryegrass grown in pure stand (P<0.001). This is in agreement with the

Berseem-annual ryegrass intercropping: effect of plant arrangement

reports of Jorgessen *et al.* (1999) and Malhi *et al.* (2002). Moreover, mixing the components within the row, compared to sowing in alternate rows, significantly increased the N content in annual ryegrass.

On the contrary, the N concentration of berseem was much greater in pure stand than in mixture; similar results have been found by Scarpello (2003) for berseem clover intercropped with annual ryegrass in a Mediterranean semi-arid environment.

In the mixture the N concentration (g kg^{-1} DM) of berseem and ryegrass forage increased significantly with increase in berseem portion of stand.

Table 1. Dry matter yield (kg ha^{-1}), N concentration (g kg^{-1} DM) and percent of N uptake derived from atmospheric nitrogen fixation (%Ndfa) for annual ryegrass and berseem clover in pure and mixed stands as affected by plant arrangement (alternate rows – BR_{AR}; same row – BR_{SR}) and seeding ratio

	Dry matter yield (kg ha^{-1})			N concentration (g kg^{-1} DM)		Ndfa (%)
	Berseem	Ryegrass	Total	Berseem	Ryegrass	Berseem
Berseem (B)	9449	–	9449	35.0	–	73.3
Ryegrass (R)	–	4826	4826	–	20.6	–
BR _{AR} (B ₂₅ -R ₇₅)	6746	2000	8746	31.6	29.9	81.1
BR _{AR} (B ₅₀ -R ₅₀)	7083	1707	8790	32.6	29.2	80.7
BR _{AR} (B ₇₅ -R ₂₅)	6948	1718	8666	33.5	31.3	78.8
BR _{SR} (B ₂₅ -R ₇₅)	6824	2458	9282	31.4	28.4	84.1
BR _{SR} (B ₅₀ -R ₅₀)	7980	1266	9246	33.7	33.3	80.2
BR _{SR} (B ₇₅ -R ₂₅)	8258	1438	9696	33.9	33.6	78.8
LSD _{0.05}	970	852	811	1.3	2.5	3.0
<i>Mean effect of plant arrangement</i>						
Alternate rows	6925	1809	8734	32.6	30.1 b	80.2
Same row	7688	1721	9409	33.0	31.8 a	81.0
<i>Mean effect of seeding ratio</i>						
BR (B ₂₅ -R ₇₅)	6785 b	2228 a	9013	31.5 b	29.1 b	82.6 a
BR (B ₅₀ -R ₅₀)	7531 a	1487 b	9018	33.1 a	31.2 a	80.4 ab
BR (B ₇₅ -R ₂₅)	7603 a	1579 b	9182	33.7 a	32.5 a	78.8 b

Different letters within the columns indicate significant different at the $P \leq 0.05$ level

On average, *Trifolium alexandrinum* achieved a higher proportion of its N from nitrogen fixation when grown in mixture with ryegrass than when it was grown in monoculture ($P < 0.0001$). In mixture ryegrass has been found to be stronger competitor for soil nitrogen (data not shown), and berseem relied more on N_2 fixation as N source. Similar observations were made by Papastylianou (1988), Loiseau *et al.* (2001), Carlsson and Huss-Danell (2003) and Stringi *et al.* (2004). This suggests that nitrogen competition plays a key role in intercropping legume-nonlegume systems and its knowledge can enable to manage better N inputs in order to make the intercropping system more efficient.

No significant differences for %Ndfa were observed due to the arrangement of plants in the mixture. The percentage of nitrogen derived from N_2 -fixation decreased with increase in berseem portion of stand. Such result seems related to the increase of soil N uptake by

Stringi et al.

ryegrass as its density increased in the mixture; this, on turn, encouraged the legume to obtain a greater proportion of its nitrogen through biological fixation.

The apparent transfer of fixed nitrogen from clover to grass was negligible in all treatment. In fact, no significant lower values for % atom ^{15}N of intercropped annual ryegrass compared to monocropped were observed, indicating that transfer of the nitrogen from legume to grass was marginal or nonexistent (Cowell *et al.*, 1989). This is in agreement with the reports of Peoples and Herridge (1990), Danso *et al.* (1993) and Stringi *et al.* (2004) that highlighted how in annual grass-legume intercropping N transfer may not be common phenomena.

Acknowledgements

This research was supported by Patti territoriali “Magazzolo-Platani”.

References

- Bulson H. A. J., Snaydon R. W. and Stopes C. E. (1997) Effects of plant density on intercropped wheat and field beans in an organic farming system. *Journal of Agricultural Science*, 128, 59–71.
- Carlsson G. and Huss-Danell K. (2003) Nitrogen fixation in perennial forage legumes in the field. *Plant and Soil*, 253, 353–372.
- Cowell L. E., Bremer E. and Van Kessel C. (1989) Yield and N_2 fixation of pea and lentil as affected by intercropping and N application. *Canadian Journal of Soil Science*, 69, 243–251.
- Danso S. K. A., Palmason F. and Hardason G. (1993) Is nitrogen transferred between field crops? Examining the question through a sweet-blue lupin (*Lupinus angustifolium* L.) – oats (*Avena sativa*) intercrop. *Soil Biology and Biochemistry*, 25, 1135–1137.
- Fried M. and Middleboe V. (1997) Measurements of amount of nitrogen fixed by a legume crop. *Plant and Soil*, 4, 713–715.
- Hogh-Jensen H. and Schjoerring J. K. (1994) Measurement of biological dinitrogen fixation in grassland: Comparison of the enriched ^{15}N dilution and the natural ^{15}N abundance methods at different nitrogen application rates and defoliation frequencies. *Plant and Soil*, 166, 154–164.
- Jorgensen F. V., Jensen E. S. and Schjoerring J. K. (1999) Dinitrogen fixation in white clover grown in pure stand and mixture with ryegrass estimated by the immobilized ^{15}N isotope dilution method. *Plant and soil*, 208, 293–305.
- Loiseau P., Soussanna J. F., Lounault F. and Delpy R. (2001) Soil N contributes to the oscillations of the white clover content in mixed swards of perennial ryegrass under conditions that simulate grazing over five years. *Grass and Forage Science*, 56, 205–217.
- Malhi S. S., Zentner R. P. and Heier K. (2002) Effectiveness of alfalfa in reducing fertilizer N input for optimum forage yield, protein concentration, returns and energy performance of brome-grass-alfalfa mixtures. *Nutrient cycling in agroecosystems*, 62, 219–227.
- Ofori F. and Stern W. R. (1987) Cereal-legume intercropping systems. *Advances in Agronomy*, 4, 41–90.
- Papastylianou I. (1988) The ^{15}N methodology in estimating N_2 fixation by vetch and pea grown in pure stand or in mixes with oat. *Plant and Soil*, 107, 183–188.
- Peoples M. B. and Herridge D. F. (1990) Nitrogen fixation by legumes in tropical and subtropical agriculture. *Advances in Agronomy*, 44, 155–223.
- Scarpello C. (2003) Efficienza biologica della consociazione trifoglio alessandrino-loiessa e stima dell'azotofissazione in rapporto a diversi livelli di azoto minerale nel suolo. *PhD thesis*, University of Palermo, Palermo, Italy.
- Stern W. R. (1993) Nitrogen fixation and transfer in intercrop systems. *Field Crops Research*, 34, 335–356.
- Stringi L., Giambalvo D., Scarpello C., Di Miceli G. and Attardo C. (2004) Nitrogen fixation by beerseem clover in pure and mixed stand in a semi-arid Mediterranean environment. In Jacobsen S.E., Jensen C.R. Porter J.R. (eds) *Proceedings of VIII ESA Congress: European agriculture in a global context* Copenhagen, Denmark 11–15 July, 671–672.
- Willey, R. W. (1979). Intercropping – its importance and research needs. Part 1. Competition and yield advantages. *Field Crop Abstracts*, 32, 1–10.