

**11<sup>th</sup> International Symposium on  
Applied Bioinorganic Chemistry**

**Barcelona, Spain**

**December 2<sup>nd</sup> – 5<sup>th</sup>, 2011**

### **International Advisory Committee**

David R. Williams (Chairman, Cardiff, UK)	Jan Reedijk (Co-chairman, Leiden, NL)
Ettore Benedetti (Naples, IT)	Franc Meyer (Goettingen, GE)
Sue Berners-Price (Nathan, AU)	Mikio Nakamura (Tokyo, JP)
Mercè Capdevila (Barcelona, SP)	Giovanni Natile (Bari, IT)
Maria Armenia Carrondo (Lisbon, PT)	Chris Orvig (Vancouver, CA)
John H. Dawson (SC, US)	João Costa Pessoa (Lisboa, PT)
Nick Hadjiliadis (Ioannina, GR)	Enrico Rizzarelli (Catania, IT)
Liang-Nian Ji (Guangzhou, CN)	Peter Sadler (Warwick, UK)
Nina Kasyanenko (St. Petersburg, RU)	Athanasios Salifoglou (Thessaloniki, GR)
Tamas Kiss (Szeged, HU)	Andrea Scozzafava (Florence, IT)
Bernhard Lippert (Dortmund, DE)	Helmut Sigel (Basel, CH)
Guillermo Mendoza-Díaz (Guanajuato, ME)	Imre Sóvágó (Debrecen, HU)

### **Local Organizing Committee**

Mercè Capdevila (Chair, UAB, Barcelona)

Olga Iranzo (Secretary, ITQB, Oeiras)

Sílvia Atrian (UB, Barcelona)

Gabriel Aullón (UB, Barcelona)

Gotzone Barandika (UPV/EHU, Vitoria/Gasteiz)

Roger Bofill (UAB, Barcelona)

Amparo Caubet (UB, Barcelona)

José M. Domínguez-Vera (UGR, Granada)

Antonio Donaire (UM, Murcia)

Virtudes Moreno (UB, Barcelona)

Òscar Palacios (UAB, Barcelona)

Joan Suades (UAB, Barcelona)

## DNA Binding and Biological Activity of Transition Metal Complexes


A. Terenzi,<sup>1</sup> G. Barone<sup>1</sup>

<sup>1</sup> Università degli Studi di Palermo, Dipartimento di Chimica "S. Cannizzaro", 90128-Palermo (Italy); [alessio.terenzi@unipa.it](mailto:alessio.terenzi@unipa.it)

The principal aim of our work is mostly focused on the synthesis of novel transition metal complexes able to non-covalently recognize DNA, using Schiff base, oxadiazole and dipyrrolyl derivatives as ligands. The coordination geometries of the molecules obtained range from tetracoordinate mononuclear complexes to binuclear metallo-supramolecular assemblies.

For instance, the DNA binding of Ni<sup>II</sup>, Cu<sup>II</sup> and Zn<sup>II</sup> complexes of Salphen[1,2] and oxadiazole[3,4] ligands mainly occurs by intercalation and groove-binding for the former and the latter, respectively. Moreover, Cu<sup>II</sup> and Zn<sup>II</sup> heteroleptic complexes of dipyrro[3,2-a:2',3'-c]phenazine (dppz) and amino acids are strong DNA-intercalators. Finally, a 2,7-diazapyrenium binuclear Pt<sup>II</sup> rectangular box-shaped metallacycle[5] is a major-groove binder that induces DNA coiling.

Interestingly, the DNA binding properties of the complexes find a positive feedback in their *in vitro* biological activity. For example, Cu<sup>II</sup> complexes of both 1,2,4- and 1,3,4-oxadiazole ligands as well as the Pt<sup>II</sup> metallacycle reduce the vitality of different human cancer cell lines in a dose- and time-dependent manner.



### Acknowledgements

Financial support from the C.I.R.C.M.S.B. (Consorzio Interuniversitario di Ricerca in Chimica dei Metalli nei Sistemi Biologici) is gratefully acknowledged.

### References

- [1] G. Barone, N. Gambino, A. Ruggirello, A. Silvestri, A. Terenzi, V. Turco Liveri, *J. Inorg. Biochem.* **2009**, *103*, 731-737.
- [2] G. Barone, A. Ruggirello, A. Silvestri, A. Terenzi, V. Turco Liveri, *J. Inorg. Biochem.* **2010**, *104*, 765-773
- [3] A. Terenzi, G. Barone, A. Palumbo Piccionello, G. Giorgi, A. Guarcello, A. Pace, *Inorg. Chim. Acta* **2011**, *373*, 62-67.
- [4] A. Terenzi, G. Barone, A. Palumbo Piccionello, G. Giorgi, A. Guarcello, P. Portanova, G. Calvaruso, S. Buscemi, N. Vivona, A. Pace, *Dalton Trans.* **2010**, *39*, 9140-9145.
- [5] V. Blanco, M.D. Garcia, A. Terenzi, E. Pia, A. Fernandez-Mato, C. Peinador, J.M. Quintela, *Chem. Eur. J.* **2010**, *16*, 12373-12380.