


Alzheimer's disease: amino acid levels and brain metabolic status

Maria Concetta Gueli & Gennaro Taibi

Neurological Sciences
Official Journal of the Italian
Neurological Society

ISSN 1590-1874

Neurol Sci
DOI 10.1007/s10072-013-1289-9



 Springer

Your article is protected by copyright and all rights are held exclusively by Springer-Verlag Italia. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your work, please use the accepted author's version for posting to your own website or your institution's repository. You may further deposit the accepted author's version on a funder's repository at a funder's request, provided it is not made publicly available until 12 months after publication.

Alzheimer's disease: amino acid levels and brain metabolic status

Maria Concetta Gueli · Gennaro Taibi

Received: 2 August 2012 / Accepted: 4 January 2013
© Springer-Verlag Italia 2013

Abstract To study brain free amino acids and their relation with dementia we measured, by high-performance liquid chromatography (HPLC), the concentration of eight free amino acids, amines and related compounds. We used temporal cortex (TC) samples obtained from 13 Alzheimer's disease (AD) patients and an equal number of age-matched controls (AC). The patterns of free amino acids, amines and related compounds showed significant quantitative changes in AD conditions with respect to healthy ones. In Alzheimer patients, lower levels of GABA were found in the TC (−57 %). Amino acids glutamate (Glu), and aspartate (Asp) concentrations, also appeared significantly reduced in the TC of AD patients (Glu: −30 %; Asp: −40 %) when compared with controls. The significant gap between methionine (Met: −30 %) and cystathionine (Cysta: +60 %) levels in TC of AD people to controls, might suggest an under/over activity of the transmethylation and transsulphuration pathways, respectively. Glutamine (Gln) and Urea were an exception to this trend because their content was higher in AD patients than in controls. Albeit these compounds may have particular physiological roles, including the possible mediation of synaptic transmission, changes in amino acid levels and related compounds (detected in steady state) suggest a modified metabolic status in brains of AD patients that reveals a reduced function of synaptic transmission. Because several evidences show that patients might display quite different concentrations of neurotransmitters in brain

areas, assessing metabolites in different and well-characterized AD stages should be investigated further.

Keywords Alzheimer's disease · Brain · Temporal cortex · Amino acids · Neurotransmitters

Introduction

Alzheimer's disease (AD) is a neurodegenerative disorder characterized by a global impairment of higher cognitive functions mostly related to failure of synaptic connections long before neurons die [1]. Studies conducted on antemortem cerebral biopsy in neocortex of patients with AD show not only that neurotransmitter systems (cholinergic, noradrenergic, serotonergic and glutamergic neurons) and pyramidal neurons are prematurely damaged in the course of the disease [2], but also that the loss of cholinergic and pyramidal neurons is related to the severity of dementia evolution [3]. The molecular etiology of AD is still incomplete. Neuronal death can occur by either apoptotic or necrotic mechanisms [4]. All studies on humans with AD indicate the amyloid- β -peptide (A β P) accumulation in the cerebral cortex as an early and invariant event in the development of AD pathology, preceding other brain lesions and clinical symptoms by many years or decades [5]. Despite extensive research, today the primary events underlying the neuropathological changes are still like pieces of a puzzle that cannot be fitted together. Thus, in view of the importance of collecting more useful information on the biochemical features characterizing the neurodegeneration in AD, we analyzed the variations in the patterns of free amino acids and related compounds in brain extracts in the disease with respect to AC. In the light of previous works on this matter [6], the described data are

M. C. Gueli (✉) · G. Taibi
Dipartimento di Biomedicina Sperimentale e Neuroscienze
Cliniche (BIONEC), Università di Palermo, Via del Vespro 129,
90127 Policlinico, PA, Italy
e-mail: mariac.gueli@unipa.it

also employed to establish a correlation between such amino acids variation and the modification of their function.

Materials and methods

Brain specimens in AC and AD neuropathologically confirmed cases (see Table 1) have been used to determine the amino acid level patterns and related compounds. Brain tissue samples (temporal cortex, grey matter) were received from the MRC London Brain Bank for Neurodegenerative Diseases, Department of Neuropathology, Institute of Psychiatry, London, UK. The AD patients fulfilled the National Institute of Neurological and Communicative Disorders and Stroke and Alzheimer's Disease and Related Disorders Association (NINCDS/ADRDA) criteria for probable AD [7]. The histological diagnosis of AD was established and was consistent with the CERAD criteria [8] for a sure diagnosis of AD. Control subjects had not any history of dementia, nor other neurologic disease. Neuropathologic examination confirmed the absence of any significant gross or microscopic brain abnormalities in controls. Specimens were transported to Palermo laboratory on dry ice and the freezing chain was never interrupted. The brain samples were coded and kept frozen at $-70\text{ }^{\circ}\text{C}$ until used.

Amino acid standards, protease inhibitors, derivatization reagents and other chemical reagents (A-grade or for chromatographic use) were purchased from Sigma (Milan, Italy).

AC and AD samples are weighed promptly with a microbalance (Mettler-H72, Zurich, Switzerland), thawed on ice, minced and then mechanically homogenized in an ice-cold solution of 10 mM Hepes buffer, pH 7.4 containing 137 mM NaCl; 4.6 mM KCl; 1.1 mM KH_2PO_4 ; 0.6 mM MgSO_4 using a Wheaton Dounce tissue grinder. To prevent artifactual proteolysis, the Protease Inhibitor Cocktail (Sigma) is added to the mixture along with 0.2 mM dithiothreitol. Homogenate is then centrifuged at 12,000g for 20 min at $4\text{ }^{\circ}\text{C}$ with a Beckman L8-70M

refrigerated centrifuge. Supernatants are removed, deproteinized with 100 μl of sulfosalicylic acid (50 % solution), and again centrifuged at 7,500g for 5 min at $4\text{ }^{\circ}\text{C}$ with an ALC 4239R refrigerated centrifuge. The clear supernatants are then withdrawn, filtered through Millipore Millex-HV (0.45 μm) and used for the amino acids and related compounds determination.

The free amino acids and related compounds contents of brain specimens have been analyzed through a Beckman's System 6300 High-Performance Amino Acid Analyzer (Beckman Coulter S.P.A. Milan, Italia) with the procedure described by Borum [9] using an Li high-performance column (10 cm) with the buffers Li A/Li B at a flow rate of 30 ml/min at 1,600 psi at a column temperature of $34\text{--}63\text{ }^{\circ}\text{C}$. It has been used ninhydrin as a colorimetric reagent. The analysis time is 53 min. Data integration has been performed by a Shimadzu Chromatopac C-R3A. Clinical and autopsy data are given in Table 1. Statistical analysis has been performed using Primer of Biostatistics (Stanton A. Glantz, McGraw-Hill-Libri, New York) and Prism (GraphPad Software, San Diego, USA). Differences between controls and pathological patients have been tested by independent two-sample *t* test with $P < 0.05$ accepted as significant.

Results

As shown in the Table 2, a total of 10 different compounds between free amino acids, amine species and related compounds are determined in the TC brain of the AC and AD patients. In general, almost all amino acids and ammonia levels are significantly lower in AD when compared with AC patients ones. It was found that Met levels showed a decrease in about 30 % in AD patients with respect of control values. When compared with AC, GABA, Glu, Asp and ornithine (Orn) levels are markedly and significantly decreased (-33 , -30 , -40 , -62 %, respectively) in TC of AD brain subjects. The change in Cysta, urea and Gln levels are an exception to this trend, their contents are greater in AD people ($+63$, $+124$, $+60$ %, respectively) than in AC cortex.

Another interesting evidence from chromatographic analysis in pathological brain extracts is the lowered peaks of ammonia (-25 %). The arginine (Arg) content in AD group is slightly lower (-27 %) than in the AC.

Discussion

Interesting considerations emerge from the evaluation of amino acids, amine patterns and related compounds from soluble extracts in brain TC of patients affected by AD

Table 1 Characterization of study groups by number, age, post-mortem interval, male-to-female ratio

	Alzheimer patients	Age-matched patients
<i>N</i>	13	13
Age (year)	71.1 ± 7.6	72.7 ± 9.7
Sex (m/f)	3/2	5/2
PMI (h)	30.09 ± 0.67	31.24 ± 0.58

Data are reported as mean \pm SEMs

PMI postmortem interval

Table 2 Levels of L-amino acids and related compounds found in human temporal cortex extracts of Alzheimer patients and age-matched patients

Compound	Alzheimer patients (μmol/g wwt)	Age-matched patients (μmol/g wwt)	Ratio Alzheimer/control
Urea	13.8 ± 11.72 ^a	6.16 ± 5.9	2.240
Aspartate (Asp)	1.80 ± 1.12 ^a	3.00 ± 1.26	0.600
Glutamate (Glu)	4.90 ± 1.69 ^a	7.02 ± 1.65	0.698
Glutamine (Gln)	2.68 ± 1.37 ^a	1.67 ± 0.61	1.600
Methionine (Met)	0.23 ± 0.09 ^a	0.32 ± 0.11	0.718
Cystathionine (Cysta)	0.80 ± 0.05 ^a	0.49 ± 0.33	1.632
GABA	0.60 ± 0.16 ^a	0.90 ± 0.28	0.666
Ammonia	0.15 ± 0.03 ^a	0.20 ± 0.07	0.750
Ornithine (Orn)	0.03 ± 0.01 ^a	0.08 ± 0.07	0.370
Arginine (Arg)	0.29 ± 0.14 ^a	0.40 ± 0.11	0.725

Data are given as mean ± standard deviation (SD) found in temporal cortex extracts from 13 Alzheimer patients and 13 normal subjects

Significance between groups was tested by two-sample independent *t* test with *P* < 0.05 accepted as significant

wwt wet weight tissue

^a Indicates values which are statistically significant between Alzheimer and control

when compared with AC. The data reveal some changes in almost all amino acids levels detected at steady state. The slightly decrease in Met levels in AD patients shows a related deficit of brain structural proteins [10]. As for amino acids, such as Asp, Glu and Met the damage consequent to their decrease will not only lead to a changed protein turnover in AD conditions, but also to a reduced availability of the special metabolites or neurotransmitters or precursors. When compared with the controls, Met levels are significantly reduced in AD group thus we think this drastic reduction could be the outcome of a metabolic pathway disorder, but it could also be a cause. Met, in the form of *S*-adenosylmethionine (SAM), is the principal donor of methyl groups in numerous methyltransferase reactions [11]. Met levels may be reduced through excessive utilization in polyamine biosynthesis. Previous studies have demonstrated markedly elevated SAM decarboxylase activity, ornithine decarboxylase (ODC) protein and spermidine levels so far [12]. As reported, SAM levels are severely decreased in AD patients' brain and CSF [13]. Moreover, MAT2A gene expression in human hepatocarcinoma cells was modulated by Met deprivation through its conversion into SAM [14]. Therefore, a Met decrease in AD patients' brain could also be an early signal or an adaptive response to the change of its modified status, and SAM level variation could be a key event for homeostasis of the brain too. The quite different pattern between Cysta levels in AD patients compared with AC caught our

attention. In fact, a high concentration of Cysta in the TC of AD patients in our experimental conditions, was an unexpected finding. The sharp Cysta levels could suggest the existence in AD brains of a particular relationship between the amount of the enzymes involved in the synthesis by cystathionine β-synthase activity and in the cleavage of the amino acid by γ-cystathionase activity. As reported, the human brains analysed by autopsy on normal subjects contain a high concentration of Cysta [15]. The presence of large amounts of Cysta in affected human brain and in developing brain has been demonstrated [16]. Thus, the surplus of Cysta level in AD brain could be used to increase cysteine pool for glutathione synthesis, the major redox buffer in mammalian cells [17, 18].

A decrease in Arg content, for instance, may have the slowdown of the citrulline-NO cycle as a consequence [19]. Furthermore, Arg decreases may partly explain a reduced local synthesis of creatine together with the markedly impaired activity of creatine kinase (CK) BB isoform, detected in AD brains [20].

A marked decrease is observed in GABA neurotransmitter in AD brain. GABA deficit in senile dementia is associated with a decreased glutamate decarboxylase activity [21]. It may be determined by a more active catabolism from the glial GABA transaminase, but it may also reflect the reduced availability of Glu, as our experimental observations have shown. Gln is an exception to this trend, its content being 1.6-fold greater in AD than in AC cortex, but it only partially contributes to the Glu homeostasis by the aging of the glutaminase activity. In fact, graded reductions of glutaminase (GA) are combined with increases in glutamine synthetase (GS) activity [22, 23]. Our results suggest an *in vivo* adaptation of the glutamate–glutamine cycle to avoid neuronal death. The increase in Gln level is a really expected result if the observed activation of glia cells in neurodegeneration is taken into consideration [24]. Hence, the GS activity should be monitored as it seems behaving like a switch under pathological conditions in brain. The decreased content of Glu in AD TC suggests its possible withdrawal to increase Gln levels in astrocytic cells. This agrees with the previous data on decreased Glu and higher Gln content reported for the same cortex area in AD postmortem brain [25, 26]. A lower Asp concentration, for instance, is detected in all AD temporal cortex specimens and we consider it may be the result of a reduced Glu concentration, as in brain the main route of Asp biosynthesis is that of glutamate–oxalacetate transamination.

Another interesting evidence from chromatographic analysis in pathological brain extracts is the lowered peaks of ammonia in AD brains. The lowered ammonia level, together with its involvement in Gln biosynthesis, might be the result of the remarkable increase in urea in brain

pathological group. This increased level could be a consequence of the altered urea diffusion in affected brain areas, determined by permeability change of the blood–brain barrier that causes its retention or a higher influx. It rather appears more realistic to assume arginine broken down by overexpressed arginase hetero-form to give unpredictable urea levels in the brain. This may contribute to a dangerous arginine depletion that may be a prerequisite for translational control of iNOS mRNA and NO generation [27]. A marked decrease in ornithine (Orn) in AD could be the virtual warning bell of a significantly increased inducible ODC activity. A recent study has shown how an overexpression of ODC had caused cellular transformation, suggesting that the key enzyme for polyamine biosynthesis should be regarded as a proto-oncogene product [28]. Recent observations have showed an attempted re-entry into the cell division cycle of the neurons in AD [29] and since neuronal maturation is associated with effective escape from the cell division cycle, emergence of quiescence may be deleterious. The current data analysis of free amino acids in brain temporal cortex of AD individuals once more points out some common aspects of the neuronal damage. Furthermore, they solicit the question whether such a dramatic amino acids variation may be almost exclusively associated with an increased protein catabolism by calpain and other proteases [30, 31], if it may also be due to a deficient protein biosynthesis or excess brain protein oxidation [10] (or to a long-term process imbalance).

The apparent deficit in amino acids and neurotransmitters in AD degenerating brains may be a protective device for a progressively reduced function, based on a long-term adaptation mechanism which might start early in life. Moreover, although not as widely appreciated, the regulation of cellular functions by major (carbohydrates, fatty acids, sterols) and minor (minerals, vitamins) dietary constituents (including the control of the gene expression), is well documented in prokaryotes and lower eukaryotes [32]. The idea amino acids can regulate gene expression in mammalian cells has been considered [33]. Mammalian cells can also adapt to dietary or pathological fluctuation in amino acids availability. It has been shown that a number of genes have specifically changed their expression levels following amino acids deprivation [34]. Under such limiting conditions, amino acids could be playing a signalling role in affected brains, conveying regulatory messages to the transcriptional machinery and affecting the turnover of specific mRNA and proteins. For instance, limitation of several amino acids greatly increases the expression of the gene encoding growth factors, binding proteins, enzymes, amino acids transport systems, retrograde tRNA transport [35, 36]. Another possibility is that this deficit could be the result of an ineffective attempt by brain tissues in repairing or gathering energy. These observations, taken together,

suggest that a condition of amino acid stress anyway caused can be crucial in starting changes in brain metabolic status. Moreover, such an adaptive process involves both neuron and glia populations, with an exaltation of glial function with respect to neuron function as a consequence of neuronal control loss [37]. To define if these present findings are trait or state markers of AD it needs further studies on brain samples obtained from AD patients with well-characterized stages of disease.

Acknowledgments We thank Prof Gert Lubec for critical reading of the manuscript.

References

- Hyman BT (1996) Anatomy of pathological alterations in Alzheimer's disease. In: Wasco W, Tanzi RE (eds) Molecular mechanisms of dementia. Humana Press Inc., Totowa, pp 219–223
- Nitsch RM (1996) From acetylcholine to amyloid: neurotransmitters and the pathology of Alzheimer's disease. *Neurodegeneration* 5:477–482
- Bowen DM, Allen SJ, Benton JS, Goodhardt MJ, Haan EA, Palmer AM, Sims NR, Smith CCT, Spillane JA, Esiri MM, Neary D, Snowden JS, Wilcock GK, Davison AN (1983) Biochemical assessment of serotonergic and cholinergic dysfunction and cerebral atrophy in Alzheimer's disease. *J Neurochem* 41:266–272
- Yuan J, Yankner BA (2000) Apoptosis in the nervous system. *Nature* 407:802–809
- Hardy JA, Higgins GA (1992) Alzheimer's disease: the amyloid cascade hypothesis. *Science* 25:184–185
- Tarbit I, Perry EK, Perry RH, Blessed G, Tomlinson BE (1980) Hippocampal free amino acids in Alzheimer's disease. *J Neurochem* 35:1246–1249
- Tierney MC, Fisher RH, Lewis AJ, Zorzitto ML, Snow WG, Reid DW, Nieuwstraten P (1988) The NINCDS-ADRDA work group criteria for the clinical diagnosis of probable Alzheimer's disease. *Neurology* 38:359–364
- Mirra SS, Heyman A, McKill D, Sumi SM, Crain BS, Brownlee LM, Vogel SS, Hughes JP, Van Belle G, Berg L (1991) The consortium to establish a registry for Alzheimer disease (CERAD II). Standardisation of the neuropathological assessment of Alzheimer's disease. *Neurology* 41:479–486
- Borum PR (1985) Manual for amino acid analysis of physiological samples. Atlanta GA American Association for Clinical Chemistry 37th National Meeting
- Smith CD, Carney JM, Starke-Reed PE, Oliver CN, Stadtman ER, Floyd RA, Markesbery WR (1991) Excess brain protein oxidation and enzyme dysfunction in normal aging and in Alzheimer disease. *Proc Natl Acad Sci USA* 88:10540–10543
- Cantoni L (1975) Biological methylation: selected aspects. *Annu Rev Biochem* 44:435–451
- Morrison LD, Bergeron C, Kish SJ (1993) Brain S-Adenosylmethionine decarboxylase activity is increased in Alzheimer's disease. *Neurosci Lett* 154:141–144
- Bottiglieri T, Godfrey P, Flynn T, Carney MWP, Toone BK, Reynolds EH (1990) Cerebrospinal fluid S-adenosylmethionine in depression and dementia: effects of treatment with parenteral and oral S-adenosylmethionine. *J Neurol Neurosurg Psychiatry* 53:1096–1098
- Martínez-Chantar ML, Latasa MU, Varela-Rey MLuC, García-Trevijano ER, Mato JM, Avila MA (2003) L-Methionine availability

- regulates expression of the methionine adenosyltransferase 2A gene in human hepatocarcinoma cells. *J Biol Chem* 278:19885–19890
15. Tallan HH, Moore S, Stein WH (1958) L-Cystathionine in human brain. *J Biol Chem* 230:707–716
 16. Okumura N, Kawai K (1961) Nitrogen metabolism of the brain. *Folia Psychiatr Neurol Jpn* 15:133–146
 17. Dringer R (2000) Metabolism and function of glutathione in brain. *Prog Neurol* 62:649–671
 18. Vitvitsky V, Thomas M, Ghorpade A, Gendelman HE, Banerjee R (2006) A functional trassulfuration pathway in the brain links to glutathione homeostasis. *J Biol Chem* 281:35785–35793
 19. Nussler AK, Billiar TR, Liu Z-Z, Morris MJ (1994) Coinduction of nitric oxide synthase and argininosuccinate synthetase in a murine macrophage cell line. *J Biol Chem* 269:1257–1261
 20. Aksenov M, Aksenova M, Butterfield DA, Markesbery WR (2000) Oxidative modification of creatine kinase bb in Alzheimer's disease brain. *J Neurochem* 74:2520–2527
 21. Bowen DM, Smith CD, White P, Dawson AN (1976) Neurotransmitter related enzymes and indices of hypoxia in senile dementia and other abiotrophies. *Brain* 99:459–496
 22. Ramonet D, Rodriguez MJ, Fredriksson K, Bernal F, Mahy N (2004) In vivo neuroprotective adaptation of the Glutamate/Glutamine cycle to neuronal death. *Hippocampus* 14:586–594
 23. Chen J, Herrup K (2012) Glutamine act as a neuroprotectant against DNA damage, be-amyloid and H₂O₂-induced stress. *PLoS One* 7(3):e33177
 24. Hamberger A, Jacobson I, Lindroth P, Mopper K, Nystrom B, Sandberg M, Molin S-O, Svanberg U (1981) Neuron-glia interactions in the biosynthesis and release of transmitter amino acids: in amino acid neurotransmitters. In: De Feudis FV, Mandel (eds) Raven Press, New York, pp 509–518
 25. Procter AW, Palmer AM, Francis PT, Lowe SL, Neary D, Murphy E, Doshi R, Bowen DM (1988) Evidence of glutamatergic denervation and possible abnormal metabolism in Alzheimer's disease. *J Neurochem* 50:790–802
 26. Fayed N, Modrego PJ, Rojas-Salinas G, Aguilar K (2011) Brain glutamate levels are decreased in Alzheimer's disease: a magnetic resonance spectroscopy study. *Am J Alzheimers Dis Other Dem* 26(6):450–456
 27. Lee J, Ryu H, Ferrante RJ, Morris SM, Ratan RR (2002) Translational control of inducible nitric oxide synthase expression by arginine can explain the arginine paradox. *PNAS* 100(8):4843–4848
 28. Auvinem M, Paasinem A, Andersson LC, Holta E (1992) Ornithine decarboxylase activity is critical for cell transformation. *Nature* 360:355–358
 29. Raina AK, Monteiro MJ, Mc Shea A, Smith MA (1999) The role of cell cycle mediated events in Alzheimer's disease. *Int Exp Path* 80:71–76
 30. LeBlanc A, Li H, Goodyer C, Bergeron C, Hammond J (1999) Caspase-6 role in apoptosis of human neurons, amyloidogenesis, and Alzheimer's disease. *J Biol Chem* 274:23426–23436
 31. Gervais FG, Xu D, Robertson GS, Vaillancourt JP, Zhu Y, Huang J, LeBlanc A, Smith D, Rigby M, Shearman MS, Clarke EE, Zheng H, Van Der Ploeg LHT, Ruffolo SC, Thornberry NA, Xanthoudakis S, Zamboni RJ, Roy S, Nicholson DW (1999) Involvement of caspases in proteolytic cleavage of Alzheimer's amyloid- β precursor protein and amyloidogenic A β peptide formation. *Cell* 97(3):395–406
 32. Fafournoux P, Bruhat A, Jousse C (2000) Amino acid regulation of gene expression. *Biochem J* 351:1–12
 33. Jousse C, Averous J, Bruhat A, Carraro V, Mordier S, Fafournoux P (2004) Amino acids as regulators of gene expression: molecular mechanisms. *BBRC* 313:447–452
 34. Bruhat A, Jousse C, Carraro V, Reimold AM, Ferrara M, Fafournoux P (2000) Amino acids control mammalian gene transcription: activating transcription factor 2 is essential for the amino acid responsiveness of the CHOP promoter. *Mol Cell Biol* 20:7192–7204
 35. Chen H, Pan Y-X, Dudenhausen EE, Kilberg MS (2004) Amino acid deprivation induces the transcription rate of the human asparagine synthetase gene through a timed program of expression and promoter binding of nutrient-responsive basic region/leucine zipper transcription factors as well as localised histone acetylation. *J Biol Chem* 279:50829–50839
 36. Shaheen HH, Horetsky RL, Kimball SR, Murthi A, Jefferson LS, Hopper AK (2007) Retrograde nuclear accumulation of cytoplasmic tRNA in rat hepatoma cells in response to amino acid deprivation. *PNAS* 104:8845–8850
 37. Van Eldik LJ (2001) Glia and Alzheimer's disease. *Neurochem Int* 39:329–331