

Cardiovascular health in migrants: current status and issues for prevention. A collaborative multidisciplinary task force report

Pietro A. Modesti^a, Stefano Bianchi^b, Claudio Borghi^c, Matteo Cameli^d, Giovambattista Capasso^e, Antonio Ceriello^f, Marco Matteo Ciccone^g, Giuseppe Germanò^h, Maria Maielloⁱ, Maria Lorenza Muiesan^j, Salvatore Novo^k, Luigi Padeletti^a, Pasquale Palmiero^l, Sergio Pillon^m, Carlo Maria Rotellaⁿ, Pier Sergio Saba^o, Pietro Scicchitano^g, Bruno Trimarco^p, Massimo Volpe^q, Roberto Pedrinelli^r, Matteo Di Biase^s, on behalf of Societa' Italiana di Cardiologia (SIC), Societa' Italiana dell'Ipertensione Arteriosa (SIIA), Societa' Italiana di Medicina Interna (SIMI), Societa' Italiana di Nefrologia (SIN), Associazione Medici Diabetologi (AMD), Associazione Italiana Aritmia e Cardiostimolazione (AIAC), Societa' Italiana dell'Obesità (SIO), Societa' Italiana di Prevenzione Cardiovascolare (SIPREC) and Osservatorio Nazionale della Sanità Elettronica e Telemedicina (ONSET)

Objectives To review information on cardiovascular health and migration, to stress the attention of researchers that much needs to be done in the collection of sound data in Italy and to allow policy-makers to identify this issue as an important public health concern.

Background In Italy, the rate of immigrants in the total number of residents increased from 2.5% in 1990 to 7.4% in 2010, and currently exceeds 10% in regions such as Lombardia, Emilia Romagna and Toscana.

Methods A consensus statement was developed by approaching relevant Italian national scientific societies involved in cardiovascular prevention. Task force members were identified by the president and/or the boards of each relevant scientific society or working group, as appropriate. To obtain a widespread consensus, drafts were merged and distributed to the scientific societies for local evaluation and revision by as many experts as possible. The ensuing final draft was finally approved by scientific societies.

Results In several western European countries, the prevalence of hypertension, diabetes, chronic kidney disease, obesity and metabolic syndrome was found to be higher among immigrants than in the native population. Although migrants are often initially healthier than non-migrant populations in their host countries, genetic factors, and changing environments with lifestyle changes, social exclusion and insufficient medical control may expose them to health challenges. Cultural reasons may also hamper

both the dissemination of prevention strategies and migrant communication with healthcare providers. However, great diversity exists across and within different groups of migrants, making generalizations very difficult and many countries do not collect registry or survey data for migrant's health.

Conclusions In the present economic context, the European Union is placing great attention to improve data collection for migrant health and to support the implementation of specific prevention policies aimed at limiting the future burden of cardiovascular and renal disease, and the consequent load for health systems. Wider initiatives on the topic are awaited in Italy.

J Cardiovasc Med 2014, 15:683–692

Keywords: cardiovascular risk, ethnic, ethnicity, global health, immigration, migrant, minorities

^aDepartment of Medicina Sperimentale e Clinica, University of Florence, Florence, ^bDepartment of Medicina Interna, Nefrologia e Dialisi, Ospedali Riuniti di Livorno, Livorno, ^cDepartment of Scienze Mediche e Chirurgiche, S.Orsola-Malpighi University Hospital, Bologna, ^dDepartment of Malattie Cardiovascolari, University of Siena, Siena, ^eDepartment of Nephrology, Second University of Naples, Naples, Italy, ^fInstitute d'Investigacions Biomèdiques August Pi i Sunyer, Barcelona, Spain, ^gDepartment of Emergenza e dei trapianti d'Organo DETO, Sezione di Malattie dell'Apparato Cardiovascolare, University of Bari, Bari, ^hDepartment of Scienze Cardiovascolari, Respiratorie, Geriatriche e Nefrologiche, University 'La Sapienza', Rome, ⁱAS Department of Cardiology, Brindisi District, ^jDepartment of Clinical and Experimental Sciences, University of Brescia, Brescia, ^kDivision of Cardiology, Dipartimento di Medicina Interna, Malattie Cardiovascolari e Nefrourologiche, University of Palermo, Palermo, ^lDivision of Cardiology, ASL BR, Brindisi, ^mUOD Telemedicina, Dipartimento Cardiovascolare, A.O.San Camillo-Forlanini, Roma, ⁿDivision of Cardiology,

Department of Fisiopatologia Clinica–Sezione di Endocrinologia, University of Florence, Florence, ⁹Division of Cardiology, AOU Sassari, Sassari, ¹⁰Department of Advanced Biomedical Sciences, Federico II University, Naples, ¹¹Division of Cardiology, Department of Medicina Clinica e Molecolare, Facoltà di Medicina e Psicologia Università di Roma 'Sapienza' – Azienda Ospedaliera Sant'Andrea, and IRCCS Neuromed, Rome, ¹²Department Cardio Toracico e Vascolare, University of Pisa, Pisa and ¹³Department of Scienze Mediche e Chirurgiche, University of Foggia, Foggia, Italy

Correspondence to Prof. Pietro Amedeo Modesti, MD, PhD, Dept. Medicina Sperimentale e Clinica, University of Florence, Largo Brambilla 3, 50134 Florence, Italy
Tel: +39 055 7949376; e-mail: pamodesti@unifi.it


Received 20 September 2013 Revised 21 February 2014
Accepted 21 February 2014

Introduction

Europe, a source of emigrants in the 19th century and first half of the 20th century, is now a net recipient of migrants.¹ In 2010 there were 47.3 million foreign-born residents in Europe, corresponding to 9.4% of the total population (Fig. 1).² Germany, France and the United Kingdom are the Member States of Europe with the highest numbers of immigrants.³ However, the migrant influx to these countries from 2001 has become relatively modest compared with Spain and Italy where the percentage of international migrants as a percentage of the population increased from 2.1 and 2.5% in 1990 to 14.1% and 7.4% in 2010, respectively.³

In Italy, migrants are an expanding global population of growing social, demographic and political importance. The rate of immigrants in the total number of residents currently exceeds 10% in regions such as Lombardia, Emilia Romagna and Toscana.⁴ The incidence is higher among minors (1 million) and among young adults (18–44 years), with consequent higher visibility at school (over 700 000) and on the labor market (over 2 million). According to 'The Immigration Statistics Dossier 2011'⁵ the foreign workers currently account for one-tenth of total employment. Foreigners, in fact, have a younger average age than Italians (32 years against 44 years for the total population),¹ with an incidence of 22.0% of minors (5.1 percentage points higher than the average) and people of working age of 75.7% (10 percentage points higher), whereas among the migrants people aged 65 years and older are only 2.4% (minus 17.9 percentage points).

Fig. 1



Population age structure by citizenship in Europe (EU-27, 2010).²

Similar age distribution was recorded in other European countries. Probably because of the young age of migrants, the attention of most European member states on migrant health is mainly focused on emergency care. Italy is one of the five countries (the Netherlands, France, Italy, Portugal and Spain) with explicit entitlements for specific services (e.g. primary care, maternity care), and/or for specific groups (e.g. children, pregnant women). In Italy, the pursuit of employment or registration as unemployed (current employment centres) give to foreign citizens legally resident (regular immigrants) the right of compulsory registration to the Servizio Sanitario Regionale (SSR).⁶ Those regular immigrants residing in Italy for a period exceeding 3 months (for study, au pairs, religious personnel), who are not entitled to compulsory registration, are required to insure against the risk of sickness, accident and maternity benefits, by entering into an insurance policy with an Italian or foreign insurance company, valid on the national territory or, alternatively, they may request voluntary registration to SSR, against payment of a contribution. Undocumented migrants in Italy are entitled to receive: emergency care, maternity care until 6 months postpartum, children's care and care and prevention of infectious diseases (including HIV and tuberculosis and children's vaccination among others). Undocumented migrants will be normally requested to pay a user's fee unless he/she officially declares to be poor and unable to pay. In this case, care is provided free of charge. Only four European Member States (the Netherlands, France, Portugal and Spain) offered undocumented migrants access to the same range of services as nationals of that country as long as they met certain preconditions, such as proof of identity or residence.^{3,7–9}

Disparities often exist between a migrant population's place of origin and its destination, particularly with relation to health determinants.¹⁰ Migration across health and disease disparities may influence the epidemiology of certain diseases globally and in nations receiving migrants. Traditionally, migration health activities have been focused on the recognition, identification and management of communicable diseases such as tuberculosis, leprosy or syphilis at the time and place of the migrant arrival.¹¹ These border control practices were designed for national application through policies of exclusion and lack an integrated international perspective. Current dynamics of migration and rates of population mobility are indeed creating new health challenges for existing policy.^{8,12} Non-communicable diseases (in particular diabetes and hypertension) have sharply increased during

the last decades in the poor areas of the planet where culture of prevention is not yet spread at population level.¹³ Lifestyle-associated health issues, including tobacco use and alcohol consumption, as well as the lack of implementation of prevention strategies among the migrant communities may also importantly affect the long-term consequences of movement between locations with different health determinants.^{12,14,15} Early studies performed mainly in Great Britain and the Netherlands revealed an increased risk of type 2 diabetes, hypertension^{16–18} obesity and metabolic syndrome in some ethnic minority groups. Cultural reasons may obstacle the dissemination of prevention strategies among immigrants, and communication between healthcare providers and migrant clients remains poor. In the present European economic context, the lack of implementation of specific national migrant health prevention policies can thus increase the burden of cardiovascular (myocardial infarction and stroke) and renal disease (dialysis) in the next few decades.¹⁹

Migrants from low and middle-income countries (LMIC) may also develop cardiovascular complications in the medium to long term as a consequence of the natural history of disease acquired in native countries and almost forgotten in Europe. The issue is relevant for rheumatic heart disease, which remains a major burden in different areas of the world.^{20–22} To be effective in the long term, the management of health issues resulting from population mobility will require an integration of national and global health initiatives.²³ For medical knowledge to improve health, health workers and patients need to hear about it, and prevention strategies and relevant treatment must be available and used appropriately. The aims of the present collaborative multidisciplinary initiative report are to collect information on different aspects of health and migration to make the health services research community and policy-makers more aware of the potential future consequent load for health systems, and to stress the importance of improving data collection for migrant health in Italy.

Health status of migrants and ethnic populations

Hypertension and stroke

Consistently, high mortality and incidence stroke rates have been observed for migrants of West African origin. Age-standardized and sex-specific stroke mortality rates for 1979–83, 1989–93 and 1999–2003 were higher for all migrant groups in each time period. More precisely in 1999–2003, stroke mortality was almost 200% higher among male migrants from West Africa, and almost 100% higher among those from the Caribbean.²⁴ These findings bear similarities with the high stroke mortality of black Surinamese and Antillean-born residents in the Netherlands, and the higher stroke mortality of black people in the United States.^{25,26} This pattern has been

attributed to high rates of hypertension among people of western African origin. There is consensus that among people of African origin hypertension is three-fold to four-fold more prevalent than in the European population.^{16,18,27–32} This is true for men and women and is present at any age, at least in adulthood. Finally, the higher rates of hypertension in people from African descent in Europe are confirmatory of the USA African-American and white comparisons.^{17,33,34}

It is uncertain whether this is mostly because of genetic or environmental factors.³⁵ A predominant role of the environment is suggested by studies that found a higher prevalence of hypertension of West African migrants compared with people in their countries of origin.²⁷ Body weight and sodium intake increases were found to be the main determinants of blood pressure (BP) increase in subjects who migrated to Israel and Italy from Ethiopia³⁶ and Somalia,³⁷ respectively. More precisely, 30% of normotensive people migrated to Italy from Somalia were hypertensive after a 6-month stay in Florence.³⁷ In the same subjects, a reduced sensitivity of the afferent pathway of the cardiopulmonary reflex with resultant defective inhibitory afference to the sympathetic outflow was also shown.³⁸ Sodium sensitivity and the reduced response of the afferent pathway of the cardiopulmonary reflex³⁸ are in agreement with the better response of hypertensive blacks to calcium blockers and diuretics than to ACE inhibitors and β -adrenergic blockers.³⁹ Further studies evaluating the efficacy of tailored dietary and pharmacologic treatment in ethnic minorities living in Europe are needed. The importance of genetic factors associated with hypertension in African blacks is also evident because differences in BP and hypertension prevalence between ethnic groups are known to exist.⁴⁰ Chinese-origin people have slightly higher BP and prevalence of hypertension than white individuals.^{41,42} Overall in South Asians, BP values were found to be similar, but there is heterogeneity among subgroups living in the same European country,¹⁷ and between the same subgroups living in different European countries.^{30,43} Increased stroke mortality levels in ethnic minorities²⁴ may also be partly linked to other factors, such as problems in timely diagnosis and care for patients with hypertension or stroke. Although the evidence on this issue is still inconclusive, there are indications of ethnic differences in the control and treatment of high BP.^{16,30,43}

Ischemic heart disease

Fischbacher *et al.*⁴⁴ examined the incidence of fatal and non-fatal myocardial infarction through a record-linked, retrospective cohort study of 4.6 million people that linked individual ethnic groups from the 2001 census to Scottish hospital discharge and mortality data from 2001 to 2003. The incidence of acute myocardial infarction (fatal combined with non-fatal) was higher in South

Asian compared with non-South Asian men, with a similar picture reported in women. Forouhi *et al.*⁴⁵ found a higher incidence of coronary death in South Asian men recruited in west London compared with European men. Trends in coronary heart disease (CHD) mortality in migrants to England and Wales were recently investigated using anonymized death records for 1979–83, 1989–93 and 1999–2003.²⁴ In all three time periods, compared with men born in England and Wales, men born in Jamaica, other Caribbean countries, or West Africa had significantly lower coronary mortality, whereas those born in East Africa, India, Pakistan or Bangladesh had higher mortality. Although there was a general pattern of decline in age-standardized and sex-specific death rates between 1979 and 2003, the declines were smaller for many migrant groups than for people born in England and Wales.²⁴ As a result, the groups with mortality higher than people born in England and Wales remained consistently so over the period. The disparity increased for coronary mortality in men born in Pakistan and Bangladesh. Other English studies confirmed the high rates of CHD mortality among residents born in the Indian subcontinent.^{24,46} A striking feature of these trends is the erosion of low coronary disease mortality in some groups. Women from Jamaica and men from Hungary now have coronary death rates higher than those born in England and Wales. In all of these groups, modifiable risk factors such as smoking, obesity and lack of sufficient physical activity are related to cardiovascular risk.²⁴ Likewise in Sweden, the incidence of myocardial infarction was 50% higher among men born in south Asia, being 20–30% lower among men born in North Africa.⁴⁷ Similar patterns are observed in other European countries.^{48–51} The causes of the higher CHD rates are still uncertain, and may also be related to genetic factors. South Asian migrants have greater CHD risk at similar levels of body mass index.

Cultural difference in symptom presentation needs to be recognized by clinicians if mortality rates in this high-risk group are to be reduced. Over 18-year follow-up of London healthy civil servants in the Whitehall II study, South Asians had higher cumulative frequencies of angina (17.0 vs. 11.3%, $P < 0.001$) compared with white individuals.⁵² However, South Asians are less likely than White British to present with classic symptoms of myocardial infarction, and some argue that this makes diagnosis difficult and possibly delays essential treatment. Bangladeshi patients with acute myocardial infarction often present with atypical symptoms, which may lead to slower triage in the casualty department and delay in essential treatment.⁵³ Design of health services must indeed recognize differential presentation according to ethnicity to improve access. This was highlighted by a UK study using the Rose Angina questionnaire, which has been extensively used in different cultural settings and epidemiological studies of CHD. In this study,

Fischbacher *et al.*⁵⁴ found that definite Rose angina showed lower sensitivity for other measures in South Asians than in Europeans: sensitivity for a doctor's diagnosis was 21% in South Asian and 37% in European men. Thus, the performance of the Rose angina questionnaire was sufficiently inconsistent to warrant further work to achieve greater cross-cultural validity.

The impact of differential access to healthcare on ethnic differences in cardiovascular mortality is unclear. Current recommendations for primary prevention of CVD in high-risk subjects include the use of antiplatelet therapy (aspirin), β -blockers, ACE inhibitors and statins. In general, there are limited data in relation to different effectiveness in the ethnicity, although there are some data in relation to different usage of these drugs, or uptake of prevention measures. For example, an audit of primary care angina management in England⁵⁵ reported smoking cessation advice in 97.1% of white Europeans, compared with 46.0% of non-white subjects. Beta-blocker use, as well as weight and exercise advice, was less common among non-white subjects. More data are available in the non-UK setting. Brown *et al.*⁵⁶ reported that African Americans and Hispanics are less likely to take aspirin than their white European counterparts, and this was not accounted for by differences in sociodemographic characteristics and CVD risk factors among ethnic minorities.

Diabetes and overweight

With few exceptions, diabetes mellitus incidence,⁵⁷ prevalence^{58–60} and mortality rates^{25,61,62} are much higher among migrants than among locally born residents. In the Netherlands, diabetes mellitus prevalence rates were two times higher among migrants (i.e. those born in Turkey, Morocco, Suriname or the Antilles) than among the locally born population.⁶³ Even larger differences were observed in terms of diabetes mellitus mortality (i.e. deaths for which diabetes mellitus was identified as the 'underlying' cause of death), with three times higher rates among migrant men, and four times higher rates among migrant women compared with the locally born population. Migrants of Surinamese origin had the highest prevalence and mortality rates.²⁵ Although the higher CAD mortality rates in migrants are closely linked to higher diabetes mellitus prevalence rates,²⁵ differences in case fatality might also play a role. However, direct evidence on case fatality among migrants in Europe is scarce. In an English follow-up survey diabetic patients from South Asia had higher diabetes mortality rates than European diabetic patients, especially at younger ages.⁵⁷ Using mortality data, a European study reviewed the extent to which diabetes mellitus mortality was more common among migrants.⁶¹ The study considered data on 30 migrant groups in seven European countries. For the majority of migrant groups, diabetes mellitus mortality was found to be much higher than for locally born residents. On average, diabetes

mellitus mortality among migrants was 90% higher for men and 120% higher for women. These findings illustrate that among non-communicable diseases with sufficient data available, diabetes mellitus is the only disease that is much more common in virtually all migrant groups⁶¹ (Table 1).

According to the Health Survey for England,⁶⁴ the prevalence of diabetes among migrants was more than five-fold that of all women and increased by 16% (compared with 2% for all women) among women aged 55 years and older. However, few national data source of risk factor exposures for ethnic groups are available. Prevalence of diabetes mellitus is closely linked to overweight and obesity. The prevalence of overweight including obesity in Bangladeshis, although lower than for the general population, increased between 1999 and 2004 by 8% for men and 14% for women (compared with 4% for all men and 3% for all women), and the prevalence of diabetes was highest in Bangladeshi men, four-fold that of men in general, in both years. High prevalence of diabetes and obesity were also observed in migrants from the middle East area.^{65–68} A certain susceptibility to insulin resistance and abdominal adiposity, the intrauterine environment and biological imprinting all act synergistically to increase the risk of diabetes mellitus in migrant populations.⁶⁹ In addition to genetic predisposition, there are several other interacting causes of the increased diabetes mellitus risk of migrants, including changing environments and insufficient medical control.^{70,71} Metabolic control is poor among migrant groups with diabetes, and HbA1c (the amount of glucose that is being carried by the red blood cells in the body) in migrants is generally higher than in the locally born population, increasing the risk of diabetic complications.⁷² A more general explanation suggests that migrants' excess diabetes mellitus mortality is due to a dramatic change from a poor to an affluent environment.⁶⁰ Lifestyle factors that are thought to be involved include both physical inactivity and unhealthy diet.

Table 1 MMRs by country-of-birth groups in Europe adjusted for age and diabetes mortality level of countries of residence⁶¹

Regression model	Age and country-adjusted MRRs ^a	
	Men MRR (95% CI)	Women MRR (95% CI)
All migrants	1.9 (1.8–2.0)	2.2 (2.1–2.3)
North African ^b	1.4 (1.3–1.4)	1.7 (1.5–1.8)
Turkish	1.7 (1.4–2.0)	2.1 (1.7–2.6)
South Asian	3.7 (3.4–4.0)	4.4 (4.0–4.9)
GDP of COB		
1Q (<\$2400)	3.8 (3.5–4.2)	3.9 (3.4–4.5)
2Q (\$2400–\$5300)	2.8 (2.6–2.9)	3.5 (3.2–3.7)
3Q (\$5300–\$9500)	1.8 (1.7–2.0)	2.2 (2.0–2.4)
4Q (>\$9500)	1.0 (0.9–1.2)	0.9 (0.8–1.1)

CI, confidence interval; COB, country of birth; GDP, gross domestic product; Q (1Q–4Q), quartile (first–fourth quartile); MMR, mortality rate ratio. ^a Derived from Poisson regression model with local-born populations as reference category. ^b Refers to migrants originating either from Algeria, Egypt, Libya, Morocco, Sudan or Tunisia.

Their joint effect is to raise the prevalence of overweight and obesity in many migrant populations as compared with locally born residents.^{60,73} Obesity may be especially important as a factor contributing to increased insulin concentrations and decreased insulin sensitivity, underlining the need for migrant-sensitive health promotion activities.^{74,75} However, higher rates of obesity and diabetes in ethnic minority groups in European countries are not fully explained by traditional risk factors (obesity, adiposity, diet and physical activity cannot explain this risk of diabetes).^{76,77} A better understanding of why this risk is increased is imperative in order to develop interventions and policies specifically in those populations.⁷⁶


The high prevalence of diabetes mellitus among migrants in Europe is also crucial for the design of screening strategies. According to a survey conducted in Amsterdam in 1999–2000,⁷⁸ the high prevalence of diabetes mellitus among Hindustani Surinamese (25.6%) and to a lesser extent among African Surinamese (12.7%) (Dutch 6.8%) suggests that universal rather than selective screening may be indicated. If a choice for selective screening is made however, an ethnicity-specific approach is required.

Renal disease

In the United Kingdom, the incidence of end-stage renal failure among individuals from ethnic communities is three to five times greater than among white Caucasians.^{79,80} High incidence rates in age-specific ethnic groups were observed for the whole of England (data from 1991–1992),⁸¹ and for the Coventry population (1992–1995 rates).⁸² The review of the London Thames area population in the early 1990s⁸³ also revealed that the relative risk of renal replacement therapy (RRT) for Asians compared with white patients aged 16–54 years was 3.0, and rose with age to 8.0 for those aged over 65 years. A similarly alarming picture was seen nationwide.⁵¹ The high incidence rates in age-specific ethnic groups observed in the years 1990–1995 were recently confirmed for the years 1999–2004.⁸⁴ There was a consistent increase of ~40% in all ethnic groups between the 1991 and 2001 cohorts (Fig. 2).⁸⁴ The increase in patient numbers was because of rising ethnic minority populations and increasing acceptance rates, especially in the elderly. Higgins *et al.*⁸² did not observe difference in the proportions of white Europeans and Indo-Asians receiving dialysis or placed on a waiting list for transplant surgery. However, proportionately more Indo-Asian patients, relative to white Europeans, started dialysis and were waiting for kidney transplantation,⁸⁵ and 33% of Indo-Asians remained on the waiting list for more than 2 years without receiving a transplant, as compared with 19% of white Europeans ($P < 0.025$). The take-on rate is likely to rise disproportionately as the migrant population ages over the next 10 years.

Organ donation was lower in the Indo-Asian population than in the white European population, with 40.1 kidneys

Fig. 2



Annualized incidence rates for renal replacement therapy by age and ethnicity (white, Indo-Asian, black) in the adult population of Birmingham Health Authority in two 5-year periods (1990–1995 and 1999–2004) using the 1991 and 2001 UK population census datasets and local programs data.⁸⁴

donated per million population per annum among white Europeans, compared with 9.3 among Indo-Asians. Factors leading to a low rate of organ donation included age distribution, the low number of Indo-Asians admitted to intensive care and religious views about organ donation from cadavers.

Knowledge of outcomes of RRT in ethnic minorities has been very limited. A study of Indo-Asians from Bradford, United Kingdom, raised the possibility of worse long-term outcomes after renal transplantation in this group compared with non-Asians, owing to a higher rate of death with graft function.⁸⁶

Patterns of renal disorder and the cause of renal failure are different in Indo-Asians compared with Caucasians, with higher rates of diabetic nephropathy, interstitial nephritis and focal segmental glomerulosclerosis.^{80,87,88} Two studies did suggest that idiopathic interstitial nephritis as well as focal, segmental glomerulosclerosis, chronic pyelonephritis and systemic lupus erythematosus was over-represented in this group.^{79,80} UK studies have demonstrated a significant correlation between the incidence/prevalence of RRT and social deprivation.^{89,90} Large studies in Australia, New Zealand and the United States have also demonstrated this link.^{91,92} The effect of deprivation is complicated by the association this has with ethnicity. The UK Renal Registry published data in 2003 showing a clear association between deprivation and ethnicity.⁹³ Healthcare practitioners should also be aware of the high prevalent use of alternative therapies and should be proactive in obtaining this information from patients.⁹⁴

The issue of reducing the excess renal disease and renal failure in ethnic communities needs to be addressed at several levels.⁹⁵ Interventions are classified as highly

cost-effective (the cost of the intervention per disease adjusted life-year saved is less than the gross domestic product per person), cost-effective (one to three times the gross domestic product per person) or not cost-effective (more than three times the gross domestic product per person). Screening for chronic kidney disease is cost-effective in people with diabetes (selective screening), whereas cost-effectiveness of screening for chronic kidney disease in the general population is unclear.⁹⁶ Data are insufficient, however, to make generalizations especially to immigrants from LMIC. The integration of screening for chronic kidney disease with cardiovascular health programs might indeed increase health awareness among migrant communities.^{97–100} Several recent studies are providing evidence that most forms of renal impairment can be prevented or delayed.^{101,102} It is thus important to ensure that best practice with respect to BP and diabetes care is implemented for all patients.^{102,103}

Issue for prevention

In spite of the unprecedented scientific advances in biomedical research over the past several decades, many under-served communities in the developed nations have not benefited equally. A complex interaction of genetic, biological, cultural and socioeconomic factors seems to be responsible for a significant proportion of the health disparities in many communities.¹⁰⁴

A number of studies have revealed important barriers in healthcare access.^{105,106} According to a recent retrospective analysis of administrative healthcare records in Italy, all quality management indicators for diabetes, heart failure, and ischemic heart disease were lower for citizens from high-pressure migration countries than for citizens from highly developed countries and Italians.¹⁰⁷ Furthermore, people from high-pressure migration countries were also more likely to contact emergency services and less likely to visit specialist doctors or use preventive care.¹⁰⁷ Importantly, patients from highly developed countries did not differ from Italians in their adherence to disease management schemes. Generally, the impact of the culture of origin diminishes with increasing length of stay.¹⁰⁸ Although the increased risk of developing CHD declined somewhat in second-generation Finnish immigrants in Sweden, the persisting risk suggests the combination of environmental and genetic factors.^{108,109}

Effective communication is essential in establishing a diagnosis and treatment plan.¹¹⁰ Although it is well recognized that our ability to provide optimal care importantly relies on our communication skills, cultural differences between the physician and the patient may create communication barriers so significant that our patients leave the office not knowing what we are telling them. Culture has been defined as 'beliefs and behaviors that are learned and shared by members of a group'.¹¹¹ Gestures are not universal; personal space, the space between individuals during a one-on-one private


conversation, is much closer in many cultures (i.e. Latin American or Middle Eastern); direct or prolonged eye contact with the patient may be thought to signify disrespect causing unnecessary miscommunications.¹¹⁰ The way a patient experiences his or her biomedical ailment is also influenced by specific cultural norms, or personal health belief systems. As a consequence, certain conditions may be 'acceptable' or even desired. In low-income countries, women have high odds of obesity and abdominal obesity.⁹⁸ The importance of culture is evident because gender differences start at puberty.¹¹² Abdominal obesity was found in nearly 80% of Iranian women living both in Iran and in Sweden.³² Likewise in the Netherlands, the prevalence of central obesity was higher in women than in men, sex differences being particularly larger in ethnic minority groups, especially in African-Surinamese than in white Dutch.¹¹³ Given the beneficial effects of physical activity on weight reduction, promotion of physical activity may help to bridge the sex gap in the metabolic syndrome, particularly in ethnic minority women in whom obesity is highly prevalent, but physical activity levels are low.¹¹³ Investment in women's education may present an important long-term investment in obesity prevention.¹¹⁴

Even while attempting to learn about specific cultural norms, the dangers of stereotyping need to be acknowledged, realizing that factors such as socioeconomic status, educational level, occupation and family values and belief systems may have as important a role in determining culture as does ethnicity.

The efficiency of prevention strategies implementation among immigrant communities requires special consideration of the importance of the social issue. By outlining an approach that examines health beliefs, decisions and behaviours within the context of culture, the Perceptions, Enablers, Nurturers (PEN-3) model seeks to empower communities through their intrinsic positive and unique qualities so that culturally appropriate interventions can be planned, implemented and evaluated.^{115–118} The PEN-3 model consists of three interrelated domains: cultural empowerment, relationships and expectations and cultural identity. Each of the domains consists of three components (Fig. 3). The first two domains, cross-tabulated in a 3 × 3 table,^{115,119} serve as the assessment tool kit to inform the intervention, whereas the last domain, cultural identity, determines the point of entry or entries for intervention (person, extended family and/or neighbourhood, i.e. community including entities such as the healthcare system), thereby increasing its effectiveness. The intervention should thus not be focused only on the individual but rather on the context and community, otherwise change may not be sustainable.

Less is known about the enabling factors, whereas insight into these is equally important in generating solutions to enhance healthcare access, CVD prevention and BP

Fig. 3



The Perceptions, Enablers, Nurturers (PEN-3) model.

control. In line with this, intervention efforts should be sensitive to the sociocultural contexts of communities in order to be effective. Television has the remarkable potential to influence people's behaviours and norms. When used for good, television can sometimes have an even bigger positive impact than a doctor's advice. For the migrant populations, such information might give insights into how exposure to different environmental situations might influence their health outcomes.

Conclusion

Changes in lifestyle and diet habits following urbanization are greatly contributing to increase the prevalence of hypertension, obesity, diabetes^{13,120–122} and kidney damage^{123,124} in LMIC. The effects of these global changes are now perceived also in Italy where although information regarding cardiovascular risk in migrants is scarce, the retrospective analysis of administrative healthcare records has been indicating that diabetes and hypertension are more prevalent among some ethnic minority groups than European descents.¹⁰⁷ The aging of ethnic minority populations can lead to a future burden of cardiovascular (myocardial infarction and stroke) and renal disease (dialysis) in the next few decades, which will increase local need for health services significantly. Therefore, in Italy, much needs to be done especially in the collection of sound data, which is the first step to allow central governments identifying this issue as an important public health concern. Specific measures to control diabetes, hypertension and secondary complications in susceptible populations will contribute both to safeguarding health and to economies in care spending. Healthcare professionals have a crucial role to play here.

Acknowledgements

The authors jointly conceived the structure for the paper. P.A.M. wrote the first draft, subsequently revised by task force members identified by the president and/or the boards of each relevant Scientific Society, as appropriate. To obtain a widespread consensus, drafts were merged and distributed to the Scientific Societies for local evaluation and revision by as many experts as possible. The

ensuing final draft was finally approved by task force members and Scientific Societies.

Conflict of interest: There are no conflicts of interest.

References

- Vasileva K. Foreigners living in the EU are diverse and largely younger than the nationals of the EU Member States. Statistics in focus. In: Eurostat, editor. Luxembourg: European Commission; 2010.
- Vasileva K. Population and social conditions. 6.5% of the EU population are foreigners and 9.4% are born abroad. Statistics in Focus. In: Eurostat, editor. Luxembourg: European Commission; 2011.
- Rechel B, Mladovsky P, Ingleby D, et al. Migration and health in an increasingly diverse Europe. *Lancet* 2013; **381**:1235–1245.
- Statistica ISd. La popolazione straniera residente in Italia. Istat, comunicato settembre 2011.
- Migrantes C. *Immigrazione. Dossier Statistico 2012*. Rome: Edizioni Idos; 2012; XXII Rapporto 2011.
- Lo Scalzo A, Donatini A, Orzella L, et al. Italy: health system review. *Health Syst Transit* 2009; **11**:1–216.
- Cuadra CB. Right of access to healthcare for undocumented migrants in EU: a comparative study of national policies. *Eur J Public Health* 2011; **22**:267–271.
- Mackenbach JP, Karanikolos M, McKee M. The unequal health of Europeans: successes and failures of policies. *Lancet* 2013; **381**:1125–1134.
- Greer SL, Hervey TK, Mackenbach JP, McKee M. Health law and policy in the European Union. *Lancet* 2013; **381**:1135–1144.
- Nygren-Krug H, World Health Organization. *International migration, health & human rights*. Geneva: World Health Organization; 2003.
- Gensini GF, Yacoub MH, Conti AA. The concept of quarantine in history: from plague to SARS. *J Infect* 2004; **49**:257–261.
- Gushulak BD, MacPherson DW. The basic principles of migration health: population mobility and gaps in disease prevalence. *Emerg Themes Epidemiol* 2006; **3**:3.
- Lozano R, Naghavi M, Foreman K, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; **380**:2095–2128.
- Fine P, Victora CG, Rothman KJ, et al. John Snow's legacy: epidemiology without borders. *Lancet* 2013; **381**:1302–1311.
- Anderson GM, Bronskill SE, Mustard CA, et al. Both clinical epidemiology and population health perspectives can define the role of healthcare in reducing health disparities. *J Clin Epidemiol* 2005; **58**:757–762.
- Cappuccio FP, Cook DG, Atkinson RW, Strazzullo P. Prevalence, detection, and management of cardiovascular risk factors in different ethnic groups in south London. *Heart* 1997; **78**:555–563.
- Agyemang C, Bhopal RS. Is the blood pressure of South Asian adults in the UK higher or lower than that in European white adults? A review of cross-sectional data. *J Hum Hypertens* 2002; **16**:739–751.
- Cappuccio FP, Oakeshott P, Strazzullo P, Kerry SM. Application of Framingham risk estimates to ethnic minorities in United Kingdom and implications for primary prevention of heart disease in general practice: cross sectional population based study. *BMJ* 2002; **325**:1271–1276.
- Raleigh VS. Diabetes and hypertension in Britain's ethnic minorities: implications for the future of renal services. *BMJ* 1997; **314**:209–213.
- Carapetis JR, Steer AC, Mulholland EK, Weber M. The global burden of group A streptococcal diseases. *Lancet Infect Dis* 2005; **5**:685–694.
- Carapetis JR, McDonald M, Wilson NJ. Acute rheumatic fever. *Lancet* 2005; **366**:155–168.
- Carapetis JR. Rheumatic heart disease in developing countries. *N Engl J Med* 2007; **357**:439–441.
- Geneau R, Stuckler D, Stachenko S, et al. Raising the priority of preventing chronic diseases: a political process. *Lancet* 2010; **376**:1689–1698.
- Harding S, Rosato M, Teyhan A. Trends for coronary heart disease and stroke mortality among migrants in England and Wales, 1979–2003: stroke declines notable for some groups. *Heart* 2008; **94**:463–470.
- Stirbu I, Kunst AE, Bos V, Mackenbach JP. Differences in avoidable mortality between migrants and the native Dutch in The Netherlands. *BMC Public Health* 2006; **6**:78.
- Keppel KG, Pearcy JN, Heron MP. Is there progress toward eliminating racial/ethnic disparities in the leading causes of death? *Public health reports (Washington, D C: 1974)* 2010; **125**:689–697.
- Cooper R, Rotimi C, Ataman S, et al. The prevalence of hypertension in seven populations of West African origin. *Am J Public Health* 1997; **87**:160–168.
- Primates P, Bost L, Poulter NR. Blood pressure levels and hypertension status among ethnic groups in England. *J Hum Hypertens* 2000; **14**:143–148.
- Agyemang C, Bhopal R. Is the blood pressure of people from African origin adults in the UK higher or lower than that in European origin white people? A review of cross-sectional data. *J Hum Hypertens* 2003; **17**:523–534.
- Agyemang C, Bindraban N, Mairuhu G, et al. Prevalence, awareness, treatment, and control of hypertension among Black Surinamese, South Asian Surinamese and White Dutch in Amsterdam, The Netherlands: the SUNSET study. *J Hypertens* 2005; **23**:1971–1977.
- Pudarcic S, Sundquist J, Johansson SE. Major risk factors for cardiovascular disease in elderly migrants in Sweden. *Ethn Health* 2000; **5**:137–150.
- Koochek A, Mirmiran P, Azizi T, et al. Is migration to Sweden associated with increased prevalence of risk factors for cardiovascular disease? *Eur J Cardiovasc Prev Rehabil* 2008; **15**:78–82.
- Agyemang C, Bhopal R, Bruijnzeels M. Negro, Black, Black African, African Caribbean, African American or what? Labelling African origin populations in the health arena in the 21st century. *J Epidemiol Comm Health* 2005; **59**:1014–1018.
- Twagirumukiza M, De Bacquer D, Kips JG, et al. Current and projected prevalence of arterial hypertension in sub-Saharan Africa by sex, age and habitat: an estimate from population studies. *J Hypertens* 2011; **29**:1243–1252.
- Agyemang C, Kunst A, Bhopal R, et al. A cross-national comparative study of blood pressure and hypertension between English and Dutch South-Asian- and African-origin populations: the role of national context. *Am J Hypertens* 2010; **23**:639–648.
- Goldbourt U, Khoury M, Landau E, et al. Blood pressure in Ethiopian immigrants: relationship to age and anthropometric factors, and changes during their first year in Israel. *Isr J Med Sci* 1991; **27**:264–267.
- Modesti PA, Tamburini C, Hagi MI, et al. Twenty-four-hour blood pressure changes in young Somalian blacks after migration to Italy. *Am J Hypertens* 1995; **8**:201–205.
- Modesti PA, Hagi MI, Corsoni V, et al. Impaired adaptation of cardiopulmonary receptors to western diet in normotensive black immigrants. *Am J Hypertens* 1999; **12**:145–150.
- Brewster LM, Seedat YK. Why do hypertensive patients of African ancestry respond better to calcium blockers and diuretics than to ACE inhibitors and beta-adrenergic blockers? A systematic review. *BMC Med* 2013; **11**:141.
- Regidor E, Astasio P, Calle ME, et al. The association between birthplace in different regions of the world and cardiovascular mortality among residents of Spain. *Eur J Epidemiol* 2009; **24**:503–512.
- Anand SS, Yusuf S, Vuksan V, et al. Differences in risk factors, atherosclerosis, and cardiovascular disease between ethnic groups in Canada: the Study of Health Assessment and Risk in Ethnic groups (SHARE). *Lancet* 2000; **356**:279–284.
- Harland JO, Unwin N, Bhopal RS, et al. Low levels of cardiovascular risk factors and coronary heart disease in a UK Chinese population. *J Epidemiol Comm Health* 1997; **51**:636–642.
- Glenday K, Kumar BN, Tverdal A, Meyer HE. Cardiovascular disease risk factors among five major ethnic groups in Oslo, Norway: the Oslo Immigrant Health Study. *Eur J Cardiovasc Prev Rehabil* 2006; **13**:348–355.
- Fischbacher CM, Steiner M, Bhopal R, et al. Variations in all cause and cardiovascular mortality by country of birth in Scotland, 1997–2003. *Scott Med J* 2007; **52**:5–10.
- Forouhi NG, Sattar N, Tillin T, et al. Do known risk factors explain the higher coronary heart disease mortality in South Asian compared with European men? Prospective follow-up of the Southall and Brent studies, UK. *Diabetologia* 2006; **49**:2580–2588.
- Bhopal R. What is the risk of coronary heart disease in South Asians? A review of UK research. *J Public Health Med* 2000; **22**:375–385.
- Hedlund E, Lange A, Hammar N. Acute myocardial infarction incidence in immigrants to Sweden. Country of birth, time since immigration, and time trends over 20 years. *Eur J Epidemiol* 2007; **22**:493–503.
- Razum O, Zeeb H. Risk of coronary heart disease among Turkish migrants to Germany: further epidemiological evidence. *Atherosclerosis* 2000; **150**:439–440.
- Regidor E, de La Fuente L, Martinez D, et al. Heterogeneity in cause-specific mortality according to birthplace in immigrant men residing in Madrid, Spain. *Ann Epidemiol* 2008; **18**:605–613.
- Ho L, Bos V, Kunst AE. Differences in cause-of-death patterns between the native Dutch and persons of Indonesian descent in the Netherlands. *Am J Public Health* 2007; **97**:1616–1618.

- 51 Albin B, Hjelm K, Ekberg J, Elmstahl S. Higher mortality and different pattern of causes of death among foreign-born compared to native Swedes 1970–1999. *J Imm Minor Health* 2006; **8**:101–113.
- 52 Zaman MJS, Shipley MJ, Stafford M, *et al.* Incidence and prognosis of angina pectoris in South Asians and Whites: 18 years of follow-up over seven phases in the Whitehall-II prospective cohort study. *J Public Health (Oxf)* 2011; **33**:430–438.
- 53 Barakat H, Barakat H, Baaj MK. CVD and obesity in transitional Syria: a perspective from the Middle East. *Vasc Health Risk Manag* 2012; **8**:145–150.
- 54 Fischbacher CM, Bhopal R, Unwin N, *et al.* The performance of the Rose angina questionnaire in South Asian and European origin populations: a comparative study in Newcastle, UK. *Int J Epidemiol* 2001; **30**:1009–1016.
- 55 Stewart A, Rao J, Osho-Williams G, *et al.* Audit of primary care angina management in Sandwell, England. *J R Soc Promot Health* 2002; **122**:112–117.
- 56 Brown DW, Shepard D, Giles WH, *et al.* Racial differences in the use of aspirin: an important tool for preventing heart disease and stroke. *Ethn Dis* 2005; **15**:620–626.
- 57 Mather HM, Chaturvedi N, Fuller JH. Mortality and morbidity from diabetes in South Asians and Europeans: 11-year follow-up of the Southall Diabetes Survey, London, UK. *Diabet Med* 1998; **15**:53–59.
- 58 Ujic-Voortman JK, Schram MT, Jacobs-van der Bruggen MA, *et al.* Diabetes prevalence and risk factors among ethnic minorities. *Eur J Public Health* 2009; **19**:511–515.
- 59 Carlsson AC, Wandell PE, Hedlund E, *et al.* Country of birth-specific and gender differences in prevalence of diabetes in Sweden. *Diabetes Res Clin Pract* 2013; **100**:404–408.
- 60 Misra A, Ganda OP. Migration and its impact on adiposity and type 2 diabetes. *Nutrition* 2007; **23**:696–708.
- 61 Vandenheede H, Deboosere P, Stirbu I, *et al.* Migrant mortality from diabetes mellitus across Europe: the importance of socio-economic change. *Eur J Epidemiol* 2012; **27**:109–117.
- 62 Patrick D, Sylvie G. Adult migrant mortality advantage in Belgium: evidence using census and register data. *Population* 2005; **60**:765–811.
- 63 Bos M, Agyemang C. Prevalence and complications of diabetes mellitus in Northern Africa, a systematic review. *BMC Public Health* 2013; **13**:387.
- 64 Stamatakis E, Ekelund U, Wareham NJ. Temporal trends in physical activity in England: the health survey for England 1991 to 2004. *Prev Med* 2007; **45**:416–423.
- 65 Bader A, Musshausser D, Sahin F, *et al.* The Mosque Campaign: a cardiovascular prevention program for female Turkish immigrants. *Wien Klin Wochenschr* 2006; **118**:217–223.
- 66 Daryani A, Berglund L, Andersson A, *et al.* Risk factors for coronary heart disease among immigrant women from Iran and Turkey, compared to women of Swedish ethnicity. *Ethn Dis* 2005; **15**:213–220.
- 67 Porsch-Oezcueromez M, Bilgin Y, Wollny M, *et al.* Prevalence of risk factors of coronary heart disease in Turks living in Germany: The Giessen Study. *Atherosclerosis* 1999; **144**:185–198.
- 68 Tran AT, Straand J, Diep LM, *et al.* Cardiovascular disease by diabetes status in five ethnic minority groups compared to ethnic Norwegians. *BMC Public Health* 2011; **11**:554.
- 69 Ramachandran A, Ma RCW, Snehalatha C. Diabetes in Asia. *Lancet* 2010; **375**:408–418.
- 70 Cruickshank JK, Mbanya JC, Wilks R, *et al.* Sick genes, sick individuals or sick populations with chronic disease? The emergence of diabetes and high blood pressure in African-origin populations. *Int J Epidemiol* 2001; **30**:111–117.
- 71 Knight TM, Smith Z, Whittles A, *et al.* Insulin resistance, diabetes, and risk markers for ischaemic heart disease in Asian men and non-Asian in Bradford. *Br Heart J* 1992; **67**:343–350.
- 72 Lanting LC, Joung IMA, Mackenbach JP, *et al.* Ethnic differences in mortality, end-stage complications, and quality of care among diabetic patients: a review. *Diabetes Care* 2005; **28**:2280–2288.
- 73 Agyemang C, Owusu-Dabo E, de Jonge A, *et al.* Overweight and obesity among Ghanaian residents in The Netherlands: how do they weigh against their urban and rural counterparts in Ghana? *Public Health Nutr* 2009; **12**:909–916.
- 74 Vlaar EMA, van Valkengoed IGM, Nierkens V, *et al.* Feasibility and effectiveness of a targeted diabetes prevention program for 18 to 60-year-old South Asian migrants: design and methods of the DH!AAN study. *BMC Public Health* 2012; **12**:371.
- 75 van de Vijver S, Oti S, Addo J, *et al.* Review of community-based interventions for prevention of cardiovascular diseases in low- and middle-income countries. *Ethn Health* 2012; **17**:651–676.
- 76 Franco M, Cooper RS, Bilal U, Fuster V. Challenges and opportunities for cardiovascular disease prevention. *Am J Med* 2011; **124**:95–102.
- 77 Abubakari AR, Lauder W, Jones MC, *et al.* Prevalence and time trends in diabetes and physical inactivity among adult West African populations: the epidemic has arrived. *Public Health* 2009; **123**:602–614.
- 78 Bindraban NR, van Valkengoed IG, Mairuhu G, *et al.* Prevalence of diabetes mellitus and the performance of a risk score among Hindustani Surinamese, African Surinamese and ethnic Dutch: a cross-sectional population-based study. *BMC Public Health* 2008; **8**:271.
- 79 Lightstone L, Rees AJ, Tomson C, *et al.* High incidence of end-stage renal disease in Indo-Asians in the UK. *QJM* 1995; **88**:191–195.
- 80 Ball S, Lloyd J, Cairns T, *et al.* Why is there so much end-stage renal failure of undetermined cause in UK Indo-Asians? *QJM* 2001; **94**:187–193.
- 81 Roderick PJ, Raleigh VS, Hallam L, Mallick NP. The need and demand for renal replacement therapy in ethnic minorities in England. *J Epidemiol Comm Health* 1996; **50**:334–339.
- 82 Higgins RM, Edmunds ME, Dukes DC. End-stage renal failure in Indo-Asians. *QJM* 1995; **88**:523–524.
- 83 Roderick PJ, Jones I, Raleigh VS, *et al.* Population need for renal replacement therapy in Thames regions: ethnic dimension. *BMJ* 1994; **309**:1111–1114.
- 84 Lambie M, Richards N, Smith S. Ethnicity, age and incidence rates for renal replacement therapy (RRT) in Birmingham, UK: 1990–2004. *Nephrol Dial Transplant* 2008; **23**:3983–3987.
- 85 Perera S, Mamode N. South Asian patients awaiting organ transplantation in the UK. *Nephrol Dial Transplant* 2011; **26**:1380–1384.
- 86 Jeffrey RF, Woodrow G, Mahler J, *et al.* Indo-Asian experience of renal transplantation in Yorkshire: results of a 10-year survey. *Transplantation* 2002; **73**:1652–1657.
- 87 Ball S, Cook T, Hulme B, *et al.* The diagnosis and racial origin of 394 patients undergoing renal biopsy: an association between Indian race and interstitial nephritis. *Nephrol Dial Transplant* 1997; **12**:71–77.
- 88 Pazianas M, Eastwood JB, MacRae KD, Phillips ME. Racial origin and primary renal diagnosis in 771 patients with end-stage renal disease. *Nephrol Dial Transplant* 1991; **6**:931–935.
- 89 Roderick P, Clements S, Stone N, *et al.* What determines geographical variation in rates of acceptance onto renal replacement therapy in England? *J Health Serv Res Pol* 1999; **4**:139–146.
- 90 Maheswaran R, Payne N, Meechan D, *et al.* Socioeconomic deprivation, travel distance, and renal replacement therapy in the Trent Region, United Kingdom 2000: an ecological study. *J Epidemiol Comm Health* 2003; **57**:523–524.
- 91 Byrne C, Nedelman J, Luke RG. Race, socioeconomic status, and the development of end-stage renal disease. *Am J Kidney Dis* 1994; **23**:16–22.
- 92 Cass A, Cunningham J, Wang Z, Hoy W. Social disadvantage and variation in the incidence of end-stage renal disease in Australian capital cities. *Aust N Z J Public Health* 2001; **25**:322–326.
- 93 Ansell D. Summary of findings in the 2006 UK Renal Registry report (chapter 1). *Nephrol Dial Transplant* 2007; **22** (Suppl 7):vii1–2.
- 94 Luyckx VA, Naicker S. Acute kidney injury associated with the use of traditional medicines. *Nat Clin Pract Nephrol* 2008; **4**:664–671.
- 95 Ali O, Mohiuddin A, Mathur R, *et al.* A cohort study on the rate of progression of diabetic chronic kidney disease in different ethnic groups. *BMJ Open* 2013; **3**:e001855.
- 96 Manns B, Hemmelgarn B, Tonelli M, *et al.* Population based screening for chronic kidney disease: cost effectiveness study. *BMJ* 2010; **341**:c5869.
- 97 Modesti PA, Bamoshmoosh M, Rapi S, *et al.* Relationship between hypertension, diabetes and proteinuria in rural and urban households in Yemen. *J Hum Hypertens* 2013; **27**:572–579.
- 98 Modesti PA, Rapi S, Bamoshmoosh M, *et al.* Impact of one or two visits strategy on hypertension burden estimation in HYDY, a population-based cross-sectional study: implications for healthcare resource allocation decision making. *BMJ Open* 2012; **2**:e001062.
- 99 Cravedi P, Sharma SK, Bravo RF, *et al.* Preventing renal and cardiovascular risk by renal function assessment: insights from a cross-sectional study in low-income countries and the USA. *BMJ Open* 2012; **2**:2.
- 100 Modesti PA, Bamoshmoosh M, Rapi S, *et al.* Epidemiology of hypertension in Yemen: effects of urbanization and geographical area. *Hypertens Res* 2013; **36**:711–717.
- 101 Ruggenenti P, Schieppati A, Remuzzi G. Progression, remission, regression of chronic renal diseases. *Lancet* 2001; **357**:1601–1608.
- 102 Kasiske BL, Wheeler DC. KDIGO clinical practice guideline for the evaluation and management of chronic kidney disease foreword. *Kidney Int Suppl* 2013; **3**:2–12.

- 103 Li PK, Chow KM, Matsuo S, *et al.* Asian chronic kidney disease best practice recommendations: positional statements for early detection of chronic kidney disease from Asian Forum for Chronic Kidney Disease Initiatives (AFCKDI). *Nephrology* 2011; **16**:633–641.
- 104 Stengel B, Tarver-Carr ME, Powe NR, *et al.* Lifestyle factors, obesity and the risk of chronic kidney disease. *Epidemiology* 2003; **14**:479–487.
- 105 Lien E, Nafstad P, Rosvold EO. Nonwestern immigrants' satisfaction with the general practitioners' services in Oslo, Norway. *Int J Equity Health* 2008; **7**:7.
- 106 Morgan M, Figueroa-Munoz JI. Barriers to uptake and adherence with malaria prophylaxis by the African community in London, England: focus group study. *Ethn Health* 2005; **10**:355–372.
- 107 Buja A, Gini R, Visca M, *et al.* Prevalence of chronic diseases by immigrant status and disparities in chronic disease management in immigrants: a population-based cohort study, Valore Project. *BMC Public Health* 2013; **13**:504.
- 108 Sundquist K, Li X. Coronary heart disease risks in first- and second-generation immigrants in Sweden: a follow-up study. *J Intern Med* 2006; **259**:418–427.
- 109 Tarnutzer S, Bopp M, Grp SS. Healthy migrants but unhealthy offspring? A retrospective cohort study among Italians in Switzerland. *BMC Public Health* 2012; **12**:1104.
- 110 Misra-Hebert AD. Physician cultural competence: cross-cultural communication improves care. *Cleve Clin J Med* 2003; **70**:289–303.
- 111 Galanti G-A. *Caring for patients from different cultures*. 4th ed. Philadelphia: University of Pennsylvania Press; 2008.
- 112 Bamoshmoosh M, Massetti L, Aklan H, *et al.* Central obesity in Yemeni children: a population based cross-sectional study. *World J Cardiol* 2013; **5**:295–304.
- 113 Agyemang C, van Valkengoed IG, van den Born BJ, *et al.* Heterogeneity in sex differences in the metabolic syndrome in Dutch white, Surinamese African and South Asian populations. *Diabet Med* 2012; **29**:1159–1164.
- 114 Aitsi-Selmi A, Chen R, Shipley MJ, Marmot MG. Education is associated with lower levels of abdominal obesity in women with a nonagricultural occupation: an interaction study using China's four provinces survey. *BMC Public Health* 2013; **13**:769.
- 115 Airhihenbuwa C, Okoror T, Shefer T, *et al.* Stigma, culture, and HIV and AIDS in the Western Cape, South Africa: an application of the PEN-3 cultural model for community-based research. *J Black Psychol* 2009; **35**:407–432.
- 116 Airhihenbuwa CO. A conceptual model for culturally appropriate health education programs in developing countries. *Int Q Community Health Educ* 1990; **11**:53–62.
- 117 Airhihenbuwa CO. On being comfortable with being uncomfortable: centering an Africanist vision in our gateway to global health. *Health Educ Behav* 2007; **34**:31–42.
- 118 Airhihenbuwa CO. Lessons of yesterday, promise of tomorrow: framing new approaches to health communications globally. *J Health Commun* 2012; **17**:629–630.
- 119 BelLue R, Okoror TA, Iwelunmor J, *et al.* An overview of cardiovascular risk factor burden in sub-Saharan African countries: a socio-cultural perspective. *Global Health* 2009; **5**:10.
- 120 Ford ES, Ajani UA, Croft JB, *et al.* Explaining the decrease in U.S. deaths from coronary disease, 1980–2000. *N Engl J Med* 2007; **356**:2388–2398.
- 121 Kearney PM, Whelton M, Reynolds K, *et al.* Global burden of hypertension: analysis of worldwide data. *Lancet* 2005; **365**:217–223.
- 122 Lim SS, Vos T, Flaxman AD, *et al.* A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; **380**:2224–2260.
- 123 Feehally J. Chronic kidney disease: health burden of kidney disease recognized by UN. *Nat Rev Nephrol* 2012; **8**:12–13.
- 124 Jha V, Garcia-Garcia G, Iseki K, *et al.* Chronic kidney disease: global dimension and perspectives. *Lancet* 2013; **382**:260–272.