

Survey of neonatal respiratory care and surfactant administration in very preterm infants in the Italian Neonatal Network

Roberto Bellù¹, Luigi Gagliardi², Paolo Tagliabue³, Carlo Corchia⁴, Valentina Vendettuoli⁵, Fabio Mosca⁵, Rinaldo Zanini¹; for the Italian Neonatal Network⁶

¹Neonatal Intensive Care Unit, Ospedale "A Manzoni", Lecco, Italy; ²Pediatrics and Neonatology Division, Woman and Child Health Department, Ospedale Versilia, Lido di Camaiore, Italy; ³Neonatology and Neonatal Intensive Care Unit, Fondazione MBBM – Ospedale San Gerardo Monza, Italy; ⁴International Centre on Birth Defects and Prematurity, Rome, Italy; and ⁵Department of Maternal and Pediatric Sciences, University of Milan; Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico, Milan, Italy

Abstract. *Introduction:* Variation of respiratory care is described between centers around the world. The Italian Neonatal Network (INN), as a national group of the Vermont-Oxford Network (VON) allows to perform a wide analysis of respiratory care in very low birth weight infants. *Methods:* We analyzed the dataset of infants enrolled in the INN in 2009 and 2010 and, for surfactant administration only, from 2006 to 2010 from 83 participating centers. All definitions are those of the (VON). A questionnaire analysis was also performed with a questionnaire on centers practices. *Results:* We report data for 8297 infants. Data on ventilator practices and outcomes are outlined. Variation for both practices and outcome is found. Trend in surfactant administration is also analyzed. **Conclusions.** The great variation across hospitals in all the surveyed techniques points to the possibility of implementing potentially better practices with the aim of reducing unwanted variation. These data also show the power of large neonatal networks in identifying areas for potential improvement. (www.actabiomedica.it)

Key words: preterm infants, ventilation, ventilatory support

Introduction

Since 2005, the Italian Neonatal Network (INN) is collecting data following the Vermont-Oxford Network (VON) as national group. This program allows for a very wide monitoring of most important outcomes and procedures for very low birth weight infants. Among neonatal procedures, ventilation and respiratory care are of paramount importance. Survival increased in the last 30 years and advances in respiratory care are among contributors of these better outcomes. Nevertheless, variation of care is described be-

tween centers around the world (1-5); moreover, compliance with more recent standards of care deserve attention and monitoring. For these reasons, it is worth describing recent data on respiratory care in a wide national setting like the INN, with particular focus on ventilator support and surfactant treatment. A dual approach was adopted: (1) an analysis of the data of the respiratory support and care offered to individual very preterm neonates enrolled in the INN, and (2) a brief survey of the protocols in use in the neonatal units adhering to the INN.

Methods

We analyzed the dataset of infants enrolled in the INN in 2009 and 2010 and, for surfactant administration only, from 2006 to 2010. The INN is the Italian branch of the VON. It includes preterm neonates up to 29 weeks + 6 days of gestational age, or neonates with a birth weight <1501 g, admitted to any of the centers within 1 month of age. In 2009, 73 centers participated, and in 2010 a further 10 contributed data (Total = 83). The neonatal units participating in the INN care vast majority of very preterm infants in Italy, and in some Italian regions the coverage is complete. Therefore, the INN represents an important source of data that reflect neonatal management in Italy. All definitions are those of the VON (www.vtoxford.org); briefly, in addition to anthropometric measures such as gestational age (GA), birth weight (BW), sex, etc, the following variables are collected for each infant: respiratory management in the delivery room (use of oxygen, bag-and-mask ventilation, intubation, surfactant administration) and in the ward. As far as ventilator support is concerned, the database collects data on procedures such as oxygen supplementation; use of N-CPAP, of N-CPAP before intubation; of high flow nasal cannulae (HFNC); of nasal intermittent mandatory ventilation (N-IMV)/synchronized intermittent mandatory ventilation (SIMV); of conventional mechanical (CMV) ventilation; of HFV; and surfactant administration in the ward. Bronchopulmonary dysplasia (BPD) refers to the oxygen requirement at 36 weeks post conceptional age. The data of the INN is compared with the data from the whole VON in 2010 (more than 57 000 infants with the same characteristics), obtained from the web-based system "Nightingale". Questionnaire analysis was performed with a questionnaire prepared by PT and sent by email to all chiefs of neonatal units adhering to the INN, with reminders in case of non-response. It included questions on the existence in the unit of protocols detailing the approach to respiratory diseases in the neonates, and which type of respiratory support was chosen as first line therapy and whether HFV was used for rescue in the event of a failure of conventional mechanical ventilation. Fifty-seven hospitals (69%) responded.

Results

We report data for 8297 infants (3981 in 2009, and 4316 in 2010), treated in 83 hospitals (73 infants dying in the delivery room were excluded). The mean GA was 29.3 weeks (SD = 2.96) and the mean BW was 1089 g (SD = 302). Mortality was 14.2%, and BPD 13.9% (15.5% in infants <33 weeks).

Table 1 shows the percentage of infants receiving any of the listed procedures, comparing the data with the data from the VON.

The frequency of use of most procedures is lower in Italian hospitals in comparison to the VON. As INN recruits infants above 30 weeks GA only if they are below 1501 g BW, these procedures are displayed in Table 2 for infants of 23–29 weeks only (the data for this population is complete in the database). The GA-associated decrease in the use of procedures is clearly apparent.

Table 1. Percentage of infants receiving any of the listed procedures.

INN	VON	
Delivery room procedures		
Supplemental oxygen	76.2	85.3
Bag-and-mask ventilation	61.3	61.8
Tracheal intubation	41.4	50.8
Surfactant administration	16.8	32.3
After delivery room		
Oxygen	80.8	88.2
CMV	53.1	62.7
HFV	15.8	22.1
N-CPAP	71.5	68.2
N-CPAP before ETT	41.9	40.0
Any surfactant	56.1	63.1
HFNC	8.2	49.9
NIMV/NSIMV	17.6	18.7
Any mechanical ventilation	55.5	65.1
Non-invasive ventilation	71.7	–
Any respiratory assistance ^a	84.5	–
NIV only ^b	34.0	–

INN, Italian Neonatal Network; VON, Vermont-Oxford Network, 2010; ETT, endotracheal intubation; HFNC, high flow nasal cannula; NIMV, nasal intermittent mechanical ventilation; NIV, non-invasive ventilation; HFV, high frequency ventilation

^aVentilation or NIV, excluding oxygen supplementation.

^bIn infants receiving any respiratory assistance, excluding oxygen supplementation

Table 2. Percentage of use of selected methods of respiratory support according to GA, 23–29 weeks (%)

	Gestational age (weeks)							Total
	23	24	25	26	27	28	29	
DR surfactant	51.8	41.8	42.6	34.8	27.6	18.2	11.3	26.9
Any Surfactant	96.6	98.1	94.3	89.0	82.0	74.2	61.9	80.0
MV	99.6	99.4	96.0	90.6	80.8	67.4	53.0	76.6
NIV	36.4	56.9	70.8	83.6	87.7	89.0	87.2	80.0
Any Resp. assistance	99.6	100.0	99.5	99.6	99.5	97.4	93.7	97.7
NIV only	0.0	0.5	3.5	9.0	18.7	30.7	42.9	21.5

Table 3 provides details regarding surfactant administration; i.e. a comparison in the time to surfactant administration in infants of 23–29 weeks between INN and rest of the VON. For all GA weeks considered, the percentage of infants treated with surfactant >2 h of age is always greater in Italy than in the VON, even for very preterm infants, where surfactant is almost universally administered.

Among the different procedures and practices, a wide variation was observed among centers; the coefficient of variation was lower for non-invasive ventilation (0.17), ventilation (0.26) and delivery room intubation (0.36) than for CPAP before endotracheal intubation (ETT) (0.58), and was highest for HFV (0.86) and delivery room surfactant (0.89). It is difficult to ascribe this broad variation to differences in case-mix, and it likely represents a “physician-driven” variation.

Looking at annual trends for surfactant administration (data from 2006 to 2010), it is clear than in INN surfactant administration is lower than in VON. In the VON there is a slightly decrease of surfactant administration over time. In the INN surfactant administration is stable in the whole time period.

Table 4 shows comparative data on ventilator procedures (2006–2010). Considering other respiratory interventions in an homogeneous period (2006–2010, with no change in baseline risk), after adjusting for confounders, infants receiving invasive mechanical ventilation decreased ([OR]=0.74, 95% [CI] 0.58–0.97)

Table 3. Surfactant administration (%) in INN and VON

	2006	2007	2008	2009	2010
INN	55,2	55,2	56,7	55,9	56,0
VON	64,2	64,0	63,8	63,6	62,9

Table 4. Frequency of selected ventilatory procedures

	2006	2010	p
Delivery room intubation	56.1%	48.1%	<0.001
nCPAP before intubation	28.8 %	45.2%	<0.001
Any ventilation	94.3%	96.7%	0.003
Mechanical ventilation	73.3%	66.4%	< 0.001
HFV	17.6%	22.9%	0.001
CMV	70.8%	64.7%	0.001
NIMV or SIMV	13.9%	25.18%	<0.001
NIV	76.9%	84.1%	<0.001
nCPAP	77.1%	84.5%	<0.001
HFNC	5.8%	16.6%	<0.001

HFV, high frequency ventilation; NIMV, nasal intermittent mechanical ventilation; NIV, non-invasive ventilation; HFNC, high flow nasal cannula

while NIV increased (OR= 1.75, 95% CI 1.38–2.21); intubation in delivery room decreased (OR=0.61, 95%CI 0.5–0.75). Considering outcomes, there was a significant reduction in mortality (OR=0.71, 95% CI 0.54–0.94) and in the combined outcome mortality or BPD (OR=0.89, 95%CI 0.69–1.1).

As for the results of the questionnaire, of the 57 hospitals that responded to the questionnaire, six said they did not have a protocol for respiratory assistance and were excluded from further analysis; all the others said that they used “non-invasive” techniques first.

In the event of needing tracheal intubation and mechanical ventilation as a first-intention treatment, two hospitals said they used IPPV/IMV; 20 synchronized IPPV/IMV (SIPPV/SIMV); 25 “volume guarantee” ventilation; and 10 hospitals used HFV (in five hospital for all infants, and in other five hospitals, depending upon BW/GA).

As for rescue HFV, seven hospitals (among those that did not always use HFV as a first intention) re-

sponded that they did not use HFV as a rescue, while all the others did use HFV as rescue.

Conclusions

This study represents a large scale, current survey of actual techniques used to provide respiratory assistance to very preterm infants, admitted to neonatal intensive care units in Italy in recent years. Our conclusions rest on more than 8000 very preterm/ very small neonates assisted in 83 centers in 2009–2010.

One major conclusion, in light of the current trend towards a lower level of invasiveness, is the number of infants who are managed without “invasive” (i.e. without endotracheal tube positioning) techniques. While invasive ventilation is always used at the lower extreme of viability, our data show that at 29 weeks, only half of the infants require intubation and receive at least one dose of surfactant, while the other half is managed non-invasively. The comparison with the data of VON shows that, on the whole, Italian neonates are treated less invasively.

In addition to the current description of use of various techniques, this study shows great variations in their use. This variation has been frequently reported (1–5]) in studies on ventilation. Despite the long time that HFV, CMV, and nasal continuous positive airway pressure (NCPAP) have been available, the differences between the centers are striking, and probably reflect different approaches to the preterm infants with respiratory distress, i.e. center-driven variation, rather than differences in case-mix.

The survey conducted among chiefs of neonatal units demonstrates that none of the respondents considers NIV unsuitable for a trial in infants. In practice, however, CPAP before intubation is used from 0% to about 80% of infants. As for the choice of invasive (with intubation) ventilation, only two hospitals used a non-synchronized “conventional” ventilation, five used HFV always, and the other five used HFV in some (smaller/more immature) infants only. The vast majority of units used synchronized ventilation, with or without volume guarantee.

It is worth noting that despite a stable baseline risk, from 2006 to 2010, we observed a lower level of invasiveness, a reduction of mechanical ventilation and an increase of NIV use, and this was accompanied by a decrease in risk-adjusted mortality and BPD.

These data form a basis for recognizing areas for potential improvement in ventilation given to preterm neonates.

The great variation across hospitals in all the surveyed techniques points to the possibility of implementing potentially better practices with the aim of reducing unwanted, center-related variation.

Last, these data show the power of large neonatal networks in identifying areas for potential improvement.

References

1. Horbar JD, McAuliffe TL, Adler SM, Albersheim S, Casady G, Edwards W, Jones R, et al. Variability in 28-day outcomes for very low birth weight infants: an analysis of 11 neonatal intensive care units. *Pediatrics* 1988; 82: 554–559.
2. Van Marter LJ, Allred EN, Pagano M, Sanocka U, Parad R, Moore M, Susser M, et al. Do clinical markers of barotrauma and oxygen toxicity explain interhospital variation in rates of chronic lung disease? The Neonatology Committee for the Developmental Network. *Pediatrics* 2000; 105: 1194–1201.
3. Lee SK, McMillan DD, Ohlsson A, Pendray M, Synnes A, Whyte R, Chien LY, Sale J. Variations in practice and outcomes in the Canadian NICU network: 1996–1997. *Pediatrics* 2000; 106: 1070–1079.
4. Vohr BR, Wright LL, Dusick AM, Perritt R, Poole WK, Tyson JE, Steichen JJ, et al.; Neonatal Research Network. Center differences and outcomes of extremely low birth weight infants. *Pediatrics* 2004; 113: 781–789.
5. Gagliardi L, Bellù R, Lista G, Zanini R; Network Neonatale Lombardo Study Group. Do differences in delivery room intubation explain different rates of bronchopulmonary dysplasia between hospitals? *ArchDis Child Fetal Neonatal Ed* 2011; 96: F30–F35.

Correspondence:
Roberto Bellù, MD,
Neonatal Intensive Care Unit,
Ospedale “A Manzoni”,
Via Eremo 9, 23900 Lecco, Italy,
Tel. +39-0341-489231
Fax +39-0341-489247
E-mail: r.bellu@ospedale.lecco.it

6 Appendix: Participants to the Italian Neonatal Network

Lecco (M Condò, D Turolì); Milano Mangiagalli (Gaia Weissmann, Mara Vanzati); Trento (Giuseppe De Nisi); Como S. Anna (E Villa); Milano Macedonio Melloni (Guido Moro); Ferrara (Pietro Guerrini, Raffaella Contiero); Udine (Serena Ellero, Riccardo Furlan); Monza (Laura Abbiati, Cesarina Borroni); Torino S. Anna (Giovanna Prandi, Claudio Fabris, Francesca Vielmi); Varese (Massimo Agosti, Francesco Tandoi, Guidali); Roma. Umberto I (Mario De Curtis, Claudio Tozzi, Renato Lucchini); Milano Buzzi (Marina Battaglioli, Gian Luca Lista, Paola Introvini); Modena. (Fabrizio Ferrari, Claudio Gallo); Palermo Cervello (Elvira Bel-lante); Mantova (Chiara Bottura, Francesca Pasquali, Simona Boccacci); Brindisi (Giuseppe Latini, Raffaella Giannuzzi); Milano Niguarda (Stefano Martinelli, Alberto Brunelli, Giuliana Motta); Campobasso (Maria Lucia Di Nunzio, Antonio Vendemiati); Camposampiero (Carlo Zorzi, Giovanna Carli, Alfiero); Rimini (Nicola Romeo, Palma Mammoliti); Torino (Emanuele Mastretta, Laura Barberis, Daniele Farina); Cuneo (Gianpaolo Gancia, Cristina Dalmazzo); Napoli Villa Betania (Marcello Napolitano, Francesco Messina); Foggia. (Giuseppe Rinaldi, Rosario Magaldi, Matteo Rinaldi, Rosangela Litta); Padova (Paola Lago, Vincenzo Zanardo, Lino Chiandetti); Treviso (Stefano Visentin); Tricase (Giuseppe Presta, Davide Cella); Cremona (Carlo Poggiani, Daniela Ferrari, Sara Parati); Siracusa (Francesco Lombardo, Rosetta Grigorio); Milano San Raffaele (Graziano Barera, Margherita Franco, Antonella Poloniatto); Bergamo (Angelo Colombo, Giampiero Burgio, Elena Sala); Messina (Ignazio Barberi, Venera Tiralongo, Alessandro Arco); Firenze Careggi (Carlo Dani, Simone Pratesi); Bologna S. Orsola (Valeria Mignatti, Gina Ancora, Giacomo Faldella, Sara Grandi); Pavia (Mauro Stronati, Gianfranco Perotti); Brescia (Gaetano Chirico, Claudio Migliori); Trieste (Sergio De Marini, Vincenzo Forleo); Napoli Federico II (Roberto Paludetto, Letizia Capasso, Giuseppina Mansi, Francesco Raimondi); No-vara (Gianni Bona, Iliaria Stucchi, Savastio, Federica Ferrero, Andrea Parola); Verona Borgo Trento (Ezio Maria Padovani, Elena Vi-viani, Laura Pecoraro); Roma. Fatebenefratelli Isola Tiberina (Rocco Agostino, Camilla Gizzi, Luca Massenzi); Bolzano (Hubert Messner, Alex Staffler); Napoli Buon Consiglio – F.B.F. (Gennaro Salvia); Acquaviva (Luigi Esposito, Vincenzo Forziati, Giusep-pe Latorre); Bologna Ospedale Maggiore (Fabrizio Sandri, Stefania Alati, Fabrizio Demaria); Cosenza (Oreste Lombardi, Carmine Deni Costabile, Gianfranco Scarpelli); Parma (Claudio Cavalli, Enrico Volante, Dott.ssa Proietti); Palermo Buccheri La Ferla F.B.F. (Olga Ganguzza, Bartolomeo Spinella); Roma S. Pietro F.B.F. (Cristina Haass, Eleonora Scapillati, Chiara Consigli); S. Giovanni Rotondo (Alberto Gatta, Pasqua Quitadamo); Pisa (Antonio Boldrini, Marco Vuerich, Emilio Sigali, Paolo Ghirri); Firenze Mey-er (Patrizio Fiorini, Letizia Petrucci, Marco Moroni); Perugia (Patrizia Braghetta, Paola Casucci, Liliana Minelli, Daniele Mezzetti); Benevento (Luigi Orfeo, Maria Gabriella De Luca); Bari Policlinico (Nicola Laforgia, Alessandro Grassi); Roma Bambino Gesù (Andrea Dotta, Ferdinando Savignoni); Siena (Franco Bagnoli, Claudio De Felice, Silvia Badii); Cesena (Augusto Biasini, Antonio Belluzzi, Marcello Stella); Roma Gemelli (Costantino Romagnoli, Enrico Zecca, Giovanni Barone); Vicenza (Paolo Colleselli, Lu-ca Vecchiato, Dott.ssa Nicolussi); Napoli Monaldi (Paolo Giliberti, Giovanni Chello, Silvana Rojo); Rho (Chiara Giovanettoni, Clelia A. Colnaghi); Versilia (Elena Verucci, Giulia Placidi); Lodi (Cesare Belloni, Giuseppe Carrera, Chiara Zambetti); Verona Os-pedale Civile Maggiore (Paolo Biban, Alessandra Serra); Ancona (Virgilio Carnielli, Annalisa Pedini); Benevento Ospedale Fatebenefratelli (Gennaro Vetranò, Giuseppe Furcolo, Brigida Pasquariello); Caserta (Luigi Falco, Gaetano Ausanio, Italo Bernar-do); Nocera Inferiore (Giuseppe Marchesano, Norberto Nosari, Paola Sarnelli); Como Valduce (Mario Maccabruni, Daniele Mer-azzi); Alessandria (Diego Gazzolo, Francesca Temporini Miriam Sabatini, Micaela Colivicchi); Bari Di Venere (Antonio Del Vec-chio, Marzia Tarantino); Reggio Emilia (Giancarlo Gargano, Simona Pedori); Thiene (Massimo Bellettato, Roberta Pesavento, Alessandra Cesaro); Reggio Calabria (Antonio Nicolò, Isabella Mondello); Napoli Cardarelli (Aniello Pugliese, Claudio Iervolino); Palermo Policlinico Giaccone (Giovanni Corsello, Mario Giuffrè); Catania Policlinico Rodoligo, Catania (Maria Betta, Mario Giuseppe Romeo, Alessandro Saporito); Pescara (Carmine D'Incecco, Barbara Perrone); Catania. Ospedale Santo Bambino (Maria Grazia Leone, Alessandro Rodonò); Genova Gaslini (Giovanni Serra, Alessia Franceschi, Francesco M. Riso); Napoli SS Annun-ziate (Mauro Carpentieri, Maria Pia Cigliano); Roma Casilino (Piermichele Paolillo, Simonetta Picone); Moncalieri (Antonio Mar-ra, Dott.ssa Rossetti, Tiziana Testa); Lecce (Fernando Del Cuore).