

Barotraumatic blowout fracture of the orbit after sneezing: Cone beam CT demonstration.

e-Poster: SP-31

Congress: ESHNR 2013

Type: Scientific Poster Presentation

Topic: Head and Neck

Authors: A. Lo Casto, P. Purpura, L. Galbo, F. Barreca, G. La Tona; Palermo/IT

Keywords: Orbit, Blowout Fracture, CBCT, Sneezing

Any information contained in this pdf file is automatically generated from digital material submitted to e-Poster by third parties in the form of scientific presentations. References to any names, marks, products, or services of third parties or hypertext links to third-party sites or information are provided solely as a convenience to you and do not in any way constitute or imply ESHNR 2013's endorsement, sponsorship or recommendation of the third party, information, product, or service. ESHNR 2013 is not responsible for the content of these pages and does not make any representations regarding the content or accuracy of material in this file.

As per copyright regulations, any unauthorised use of the material or parts thereof as well as commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is strictly prohibited.

You agree to defend, indemnify, and hold ESHNR 2013 harmless from and against any and all claims, damages, costs, and expenses, including attorneys' fees, arising from or related to your use of these pages.

Please note: Links to movies, ppt slideshows and any other multimedia files are not available in the pdf version of presentations.

no congress homepage

1. Purpose

To describe a case of barotraumatic blowout fracture of the right orbit after sneezing diagnosed by CBCT in a patient with no history of trauma or sinus surgery.

2. Materials and Methods

A forty-year-old man complained sudden right diplopia after vigorous sneezing. No history of trauma or previous sinus surgery was reported by the patient. The patient was submitted elsewhere to a MRI study for persisting diplopia, with administration of i.v. gadolinium contrast. One week later the patient was submitted to a maxillofacial CBCT (Scanora 3D, Soredex, Tuusula, Finland) with 90 kVp and 13 mA, 20 s rotation time, FOV 13 x 14.5 cm, 0.25 x 0.25 mm pixel size, at our Institution.

3. Results and Discussion

Results.

A reduced size and a different shape of the right maxillary sinus with respect to the left maxillary sinus, a slightly increased vertical diameter of the right orbit with respect to the left orbit, and a round shape of the inferior rectus muscle on coronal sections were present on MRI study. A maxillary sinus effusion that was hyperintense on T1 with shading on T2 images was addressed as haemosinus (Fig. 1). After gadolinium i.v. administration a strong peripheral enhancement of the maxillary sinus wall was noted (Fig. 1). A blowout fracture of the orbital floor and lateral orbital wall with an intact orbital rim and a maxillary sinus effusion with an air liquid level were detected at the CBCT study (Fig. 2). Our conclusion was that the patient had a barotraumatic, isolated, pure blowout fracture of the right orbit as a consequence of the episode of vigorous sneezing. The patient was thus treated by cortisone and antibiotic therapy to reduce the risk of sinusitis and orbital cellulitis, invited to possibly avoid nose blowing, sneezing, coughing, and vomiting. Diplopia resolved after two weeks.

Discussion.

Blowout fracture of the orbit is usually linked to facial trauma injuries (1). Rarely this type of orbital fracture is related to sneezing in patients with no history of trauma or sinus surgery, and to date just one case with associated orbital emphysema has been reported in the literature (2). Cone beam CT (CBCT) is widely used in the diagnosis of hard tissue anomalies of the maxillofacial district, however its use in maxillofacial trauma is limited and just one extensive study is available only for minor injuries as nasal fractures (3). Blowout fractures of the orbit occur when the fracture fragments extend beyond the orbit into the maxillary or ethmoid sinus due to relative weakness of the inferior and medial wall of the orbit. An orbital blowout fracture is defined as pure if the orbital rim is intact, while the impure type involves the orbital rim and may be associated to zygomaticomaxillary fractures (1, 4). Orbital fractures are commonly related to blunt trauma occurring during motor vehicle accidents, falls, assaults, sport injuries and other traumatic events (5). Isolated pure orbital wall fracture has a relatively low incidence, in fact it is diagnosed in only 13.3% of all facial bone fractures (1). A blowout fracture of the orbit is far less commonly related to sneezing (2) or nose blowing (6-10). Two theories are reported to explain the pathogenesis of orbital blowout fractures (10). The hydraulic theory assesses that fractures of the thin orbital floor are brought about by hydraulic forces, which cause posterior globe displacement and increased orbital pressure (11). The buckling theory states that a direct trauma to the inferior orbital rim causes the floor to buckle (12). Because no trauma was associated with the case we described, the hydraulic theory seems to explain its occurrence. In this case the compressed air forced through the nasal cavity should have energy enough to fracture the thin orbital walls as previously hypothesized by other authors (7, 10). High pressure can be in fact generated during nose blowing or sneezing as experimentally demonstrated both in healthy subjects and in patients with

chronic sinusitis or septal deviation (13, 14), and the amount of energy produced is comparable to that needed to fracture the orbital floors obtained from human cadavers in an experimental study (10,15). As far as we know, only one case of orbital blowout fracture after a protracted episode of vigorous sneezing has been previously reported in the literature, but it involved the medial wall of the orbit and was associated with orbital emphysema, ecchymosis of the right eye, pressure within the orbit, periorbital swelling, inflation of periorbital region and crepitus around the eye, but no pain or vision disturbance (2). In the patient we observed none of these symptoms and among them especially orbital emphysema was present, the blowout fracture involved the floor and lateral wall of the right orbit and main symptom complained was diplopia. Furthermore this is the first case studied by CBCT. The value of CBCT in the diagnosis of minor facial injuries as nasal fractures was previously demonstrated (3). In our patient a deformity of the right maxillary sinus with haemosinus and a slightly increased vertical diameter of the right orbit were detected by MRI, but a clear assessment of the thin orbital wall was not possible. The diplopia our patient showed, which resolved after two weeks, could be related to the involvement of the right inferior rectus muscle, that, although not herniated in the maxillary sinus, showed a round shape on coronal MR images. As previously reported, this shape could be related to the involvement of the fascial sling of the inferior rectus muscle in the orbital floor defect (5). The incarceration of extraocular muscles within an orbital fracture is the most commonly reported cause for extraocular movement limitation and diplopia. In orbital blowout fracture diplopia is more frequently associated to a fracture of the orbital floor rather than of the lamina papyracea (1). However it was postulated that traumatic haemorrhage causes swelling of the posterior inferior orbital fat, which contains a network of venules and bands of fibrous tissue, connecting the inferior rectus and inferior oblique muscles to the periosteum of the orbital floor. The swelling creates strain in the fibrous connective tissue of the posterior inferior orbital fat and thus on inferior rectus and inferior oblique muscles with consequent diplopia (16). This mechanism could explain the resolution of diplopia after some weeks in our patient and in other series, where no incarceration of the inferior rectus muscle was observed although an orbital floor fracture was evident (1, 16).

References

1. Park MS, Kim YJ, Kim H, Nam SH, Choi YW. Prevalence of Diplopia and Extraocular Movement Limitation according to the Location of Isolated Pure Blowout Fractures. *Arch Plast Surg* 2012;39:204-208
2. Khader QA, Abdul-Baqi KJ. Orbital emphysema after a protracted episode of sneezing in a patient with no history of trauma or sinus surgery *Ear Nose Throat J* 2010; 89:E12-3
3. Bremke M, Wiegand S, A.M. Sesterhenn, Eken M, Bien S, Werner JA. Digital volume tomography in the diagnosis of nasal bone fractures. *Rhinology* 2009; 47:126-131
4. Lo Casto A, Priolo GD, Garufi A, Purpura P, Salerno S, La Tona G, Coppolino F. Imaging evaluation of facial complex strut fractures. *Semin Ultrasound CT MR* 2012; 33:396-409
5. Hopper RA, Salemy S, Sze RW. Diagnosis of midface fractures with CT: what the surgeon needs to know. *Radiographics* 26:783-793, 2006, Kim JJ, Huoh K. Maxillofacial (midface) fractures. *Neuroimag Clin N Am* 2010; 20:581-596
6. Oluwole M, White P. Orbital floor fracture following nose blowing. *Ear Nose Throat J* 1996;75:169-170.
7. Suzuki H, Furukawa M, Takahashi E, Matsuura K. Barotraumatic blowout fracture of the orbit. *Auris Nasus Larynx* 2001;28:257-259
8. Zechmann C, Giesel FL, Nolden M, Meinzer HP, Düber C. Einseitige lidschwellung nach

einfachem nase-schnäuzen. Radiologe 2004; 44:1026-1028

9. García de Marcos JA, del Castillo-Pardo de Vera JL, Calderón-Polanco J. Orbital floor fracture and emphysema after nose blowing. Oral Maxillofac Surg 2008;12:163-165

10. Watanabe T, Kawano T, Kodama S, Suzuki M. Orbital blowout fracture caused by nose blowing. Ear Nose Throat J 2012;91:24-25

11. Smith B, Regan WF Jr. Blow-out fracture of the orbit; mechanism and correction of internal orbital fracture. Am J Ophthalmol 1957; 44: 733-739

12. Fujino T. Experimental "blow-out" fracture of the orbit. Plast Reconstr Surg 1974; 54: 81-82

13. Gwaltney JM, Handley JO, Philips CD, Bass CR, Mygind N, Winther B. Nose blowing propels nasal fluid into the paranasal sinuses. Clin Infect Dis 2000;30:387-391

14. Clement P, Chovanova H. Pressures generated during nose blowing in patients with nasal complaints and normal test subjects. Rhinology 2003; 41: 152-158

15. Bullock JD, Warwar RE, Ballal DR, Ballal RD. Mechanisms of orbital floor fractures: A clinical, experimental, and theoretical study. Trans Am Ophthalmol Soc 1999; 97:87-110

16. Putterman AM. Management of orbital floor blowout fractures. Adv Ophthalmic Plast Reconstr Surg 1987;6:281-5

Figures

Fig. 1. MRI study. Barotraumatic pure blowout fracture of the right orbit. A) Coronal TSE T2 image. Ovalar shape of the right inferior rectus muscle. Hyperintense rim of the right maxillary sinus with an intermediate intensity liquid effusion. B) Coronal TSE T2 image at a more anterior level than A). The vertical diameter of the right orbit is slightly increased with respect to the left one. C) Coronal SE T1 after Gd i.v. administration. Enhancing rim of the right maxillary sinus. D) Axial SE T1 image. Hyperintense content due to haemosinus with air liquid level in the right maxillary sinus.

1a.jpg

1b.jpg

1a - 1b

1c.jpg

1c

1d.jpg

1d

Fig. 2. CBCT study. Barotraumatic pure blowout fracture of the right orbit. A) Coronal MPR image. The right orbital floor is collapsed. Air liquid level in the right maxillary sinus. B) Coronal MPR image. Fracture of the lateral wall of the right maxillary sinus. C) Axial MPR image. The right lamina papyracea is intact. D) 3D image. The right orbital rim is intact.

2a.jpg

2a

2b.jpg

2b

2c.jpg

2c

2d.jpg

2d

4. Conclusion

CBCT correctly allows to detect the isolated pure orbital blowout fracture, involving the floor and lateral wall of the right orbit, and effusion with an air liquid level in the maxillary sinus. Radiologists and more generally doctors should be aware that a barotraumatic fracture of the orbit after sneezing should also be included among the causes of sudden onset of diplopia.

5. Mediafiles

1a.jpg

1b.jpg

1c.jpg

1d.jpg

2a.jpg

2b.jpg

2c.jpg

2d.jpg