

# Evidence of a Chromosomal Polymorphism Unique to Cercopithecini. A Key Factor in the Tribe Definition ?

Sineo Luca<sup>1</sup>, Lo Bianco Stefania<sup>1</sup> and Picone Barbara<sup>2\*</sup>

<sup>1</sup>University of Palermo, School of Basic and Applied Sciences, Palermo (Italy)

<sup>2</sup>Department of Genetics, Stellenbosch University, Private Bag X1, Matieland 7602, South Africa

\*Corresponding author: Barbara Picone, Department of Genetics, Stellenbosch University, Private Bag X1, Matieland 7602, South Africa, Tel: +3909123891817; E-mail: [barbara.picone@gmail.com](mailto:barbara.picone@gmail.com)

Rec date: October 27, 2015; Acc date: October 29, 2015; Pub date: November 05, 2015

Copyright: © 2015 Sineo et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Editorial

There are obvious gaps in the comprehension of genomic mechanisms leading to speciation [1]. This arise the question: "Are genetic differences and polymorphism highly responsible in the species isolation mechanisms? Despite many efforts, it might be fundamental collect evidences and bringing more examples in the discussion. Nevertheless, the existence of alternative and incompatible species concepts reflects a basic disagreement about the nature of species. By introducing the biological species concept, Mayr [2] stated that the species is an assemblage of interbreeding natural populations reproductively isolated. Cracraft [3] proposed the phylogenetic species concept affirming that a species is a monophyletic cluster of organisms, a prelude to the ontological concept of Christoffersen [4] that defines the species as the product of a single lineage evolution in a interbreeding history. Species, as concrete biogeographical entity, has a genetic base that limit its essence and protect its existence. Genetic and chromosomal barriers are important in the limitation of fertility and in the arose of varieties and species in animals and plants [5-7]. Conversely, several examples, like Tribe Papionini in Primates, indicate that species differentiation is not necessarily related to genomic differences and/or levels of genetic variability and chromosomal polymorphism. Both variability and polymorphism are highly scrutinized by the natural selection. According to human genetics studies (the largest body of information on genetic variability we have assembled analysing ourselves), humans (HSA) have a level of endured polymorphism in their karyotype. This polymorphism is restricted to the quantitative expression of autosomal heterochromatic traits in pericentromeric areas, or in the expression of NOR's districts, or in the HSA specific Y chromosome q-arm; polymorphism associated to balanced inversions are rare, and unbalanced translocations are strongly selected. Intrinsic genetic mechanisms work well in the protection of species identity.

On the other side, intrinsic genetic barriers are evidently not necessary in potentially interfertile sympatric species, that are reproductively isolated by behavioural mechanisms (sympatric *Macaca* species with identical genome organization in 42 homologous chromosomes, maintain their reproductive isolation). Even if it is possible to describe "local" break down in reproductive isolation either in captivity and/or in nature, there is still a open discussion on the (rare) efficacy of hybridization in species creation.

Karyotype assemblage and chromosome features assume a potential importance in speciation. Molecular chromosome studies are certainly a tool in phylogenetic studies [8-10].

If in Papionini (see above discussion) the speciation event is not related to chromosomal changes, and species maintain their isolation, using rigid behavioural schemes, in Cercopithecini, chromosomal rearrangements have played a fundamental role in the birth and separation of populations. *Cercopithecus* and related species are important models for the study of primate evolution during the Plio-Pleistocene. The group includes genera and species characterised by facial patterns, various ecological distribution, and a vivacious reshuffling of chromosomal features in the different species or in different Species-Group [11]. Some plesiomorphic character like "catamenial swelling" in *Allenopithecus* or *Miopithecus* challenge the classical composition of the tribe and indicate a weak phylogenetic relationship of these genera with real guenons. It is known that Cercopithecini Tribe are characterised by a fission-fusion model of chromosomal evolution, and that some synteny disruption occurred in different species with different mechanisms [12]. The intense chromosomal diversity among species has been interpreted as a motif of speciation (reproductive barriers), together with biogeographical clusterization in the African geography.

What if the Cercopithecini in addition to an highly diploid variation, have a "tribal-specific" chromosomal polymorphism? Cercopithecini tribe is characterised by a HSA20/21 syntenic association [13], an apomorphic trait acquired during the primate evolution. Interestingly, this synteny, constructed by two small human hortologous, can be found in two forms (heteromorphy), in homozygosity or frequently in heterozygosity in different species.

The first description of this chromosomal polymorphism was in occasion of a GTG banding on a *C. nictitans* from the Zoo of Barcelona [14]. Subsequently, a second report was on a *C. nictitans* stampflii from the Zoo of Zurich [15]. This polymorphism was then investigated in other guenons of the same Superspecies group with widely dispersed distribution: *C. mitis maesi* and *C. albogularis labiatus*, animals coming from South African facilities and from the Transvaal forests respectively [16].

On the base of GTG banding this polymorphism was described as a pericentric inversion followed by a paracentric inversion. The pericentric inversion hypothesis seems confirmed by molecular cytogenetics data on other guenons like *C. petaurista* and other species in the Campbelli and Cephus species group (unpublished data), although a more intriguing hypothesis of a Evolutionary New Centromere (i.e. the activation of centromeric silenced sequences) has been recently proposed by Stanyon et al. [17].

The scrutiny of literature data suggests the presence of the same polymorphis trait even in other Cercopithecoidea related to the genus *Cercopithecus* [18,19].

The presence of this polymorphism in different species could be a symptom of particular chromosomal dynamics for this syntenic association, and not a result of speciation and genetic drift [20]. Chromosome 21 in humans is a gene poor chromosome with vast areas of duplication and, interestingly, does not contain Fragile Sites. Chromosome 22 is more rich in important genes and has only one rare Fragile Site in its length. The inversion (or the ENC, if defined in future by molecular approaches), has presumably derived from a common ancestor in Cercopithecoidea. In the hypothesis of an inversion, these mechanisms spread because they prevent recombination from breaking apart sets of alleles that work well in an ecological or sexual setting. Since recombination continues normally within the populations of inverted and uninverted chromosomes, inversions may escape many of the deleterious consequences suffered by other genetic mechanisms that shut down recombination entirely [21]. In many species, including plants, fungi, insects and humans, there is evidence that inversions respond to natural selection; however few genes or other chromosomal features that are the results of selections have been ambiguously identified. Thus the mechanisms that affect most of the inversions remains unknown [22].

A part from the molecular mechanism involved in this heteromorphism, our interest is in the phyletic meaning of this peculiarity, in the hypothesis that this polymorphism predate speciation, a very interesting situation that could be symptom of a gene flow and of a phylogenetic and chronologic vicinity within the species in the Tribe. The sporadic but distributed frequency of this polymorphism is unusual and seems the first case of chromosomal polymorphism involving different related-species in a Primates genus and in its sister genera.

Tribe Cercopithecini offers a great variety of interesting parameters that should be carefully interpreted in the light of a multidisciplinary approach and of a re-defined ecological framework to its evolution.

## References

1. Sehausen O, Butlin RK, Keller I, Wagner CE, Boughman JW, et al. (2014) Genomics and the origin of species. *Nature Reviews Genetics* 15: 176-192.
2. Mayr E, Ashlock PD (1969) "Principles of systematic zoology", McGraw-Hill, New York.
3. Cracraft J (1989) Species and its ontology: the empirical consequences of alternative species concepts for understanding patterns and processes of differentiation. In: Otte and Endler (eds). *Species and its consequences* 28-59.
4. Christoffersen ML (1995) Cladistic taxonomy, phylogenetic systematics and evolutionary ranking. *Syst Biol* 44: 440-454.
5. Solano E, Castiglia R, Capanna E (2009) Chromosomal evolution of the house mouse, *Mus musculus domesticus*, in the Aeolian Archipelago (Sicily, Italy). *Biological Journal of the Linnean Society* 96: 194-202.
6. Lowry DB, Willis JH (2010) A widespread chromosomal inversion polymorphism contributes to a major life-history transition, local adaptation, and reproductive isolation. *PLoS Biology*.
7. Silva AA, Braga LS, Guedes RN, Tavares MG (2015) Cytogenetic analyses using C-banding and DAPI/CMA3 staining of four populations of the maize weevil *Sitophilus zeamais* Motschulsky, 1855 (Coleoptera, Curculionidae). *Comp Cytogenet* 9: 89-102.
8. Nadeau JH, Taylor BA (1984) Lengths of chromosomal segments conserved since divergence of man and mouse. *Proc Natl Acad Sci* 81: 814-818.
9. Pevzner P, Tesler G (2003) Human and mouse genomic sequences reveal extensive breakpoint reuse in mammalian evolution. *Proc Natl Acad Sci* 100: 7672-7677.
10. Murphy WJ, Larkin DM, Everts-van der Wind A, Bourque G, Tesler G, et al. (2005) Dynamics of mammalian chromosome evolution inferred from multispecies comparative maps. *Science* 309: 613-617.
11. Hill OWC (1966) *Primates Comparative Anatomy and Taxonomy*. Cercopithecoidea, Edinburgh Univ Press.
12. Romagno D, Chiarelli B, Guarducci S, Uzielli MLG, Sineo L (2000) Chromosome mapping of GABRB3 and PML loci in *Macaca* and *Cercopithecus* indicates the mechanism of evolution of human chromosome 15. *Chromosome Res* 8: 747-749.
13. Stanyon R, Rocchi M, Capozzi O, Roberto R, Misceo D, et al. (2008) Primate chromosome evolution: Ancestral karyotypes, marker order and neocentromeres. *Chromosome Res* 16:17-39.
14. Ponsà M, Estop A, Egozcue J, Garcia M, Miro R (1981) Chromosomes in the species *Cercopithecus pogonias* and *Cercopithecus nictitans*. *Int J primatol* 1: 525-533.
15. Sineo L, Scheffrahn W, Glaser D, Maurer A (1986) The banded chromosomes of *Cercopithecus nictitans stampflii*. *International Journal of Anthropology* 1: 369-373.
16. Sineo L (1990) The banded karyotype of the samango monkey compared with the karyotype of the maesi monkey and the stampflii. *International Journal of Primatology* 11: 541-552.
17. Rocchi M, Archidiacono N, Schempp W, Capozzi O, Stanyon R (2012) Centromere repositioning in mammals. *Heredity* 108: 59-67.
18. O'Brien SJ, Menninger JC, Nash WJ (2006). *Atlas of Mammalian Chromosomes*.
19. Moulin S, Gerbault-Seureau M, Dutrillaux B, Richard FA (2008) Phylogenomics of African guenons. *Chromosome Res* 16: 783-799.
20. Kirkpatrick M, Barton N (2006) Chromosome inversions, local adaptation and speciation. *Genetics* 173: 419-434.
21. Otto SP, Lenormand T (2002) Evolution of sex: Resolving the paradox of sex and recombination. *Nature Reviews Genetics* 3: 252-261.
22. Hoffmann AA, Rieseberg LH (2008) Revisiting the impact of inversions in evolution: from population genetic markers to drivers of adaptive shifts and speciation?. *Annu Rev Ecol Evol Syst* 39: 21-42.