

LIFE EVENTS AND PRIMARY MONOSYMPTOMATIC NOCTURNAL ENURESIS: A PEDIATRIC PILOT STUDY

LUCIA PARISI^{*}, MARIA ANTONIETTA FARALDO^{**}, MARIA RUBERTO^{***}, MARGHERITA SALERNO^{****}, AGATA MALTESE^{*}, ANNABELLA DI FOLCO^{*}, GIOVANNI MESSINA^{*****}, TERESA DI FILIPPO^{*}, MICHELE ROCCELLA^{*}

^{*}Department of Psychological, Pedagogical and Educational Sciences, University of Palermo, Italy - ^{**}Clinic of Child and Adolescent Neuropsychiatry, Department of Mental Health and Physical and Preventive Medicine; Second University of Naples, Italy - ^{***}Department of Medical-Surgical and Dental Specialties, Second University of Naples, Italy - ^{****}Sciences for Mother and Child Health Promotion, University of Palermo, Italy - ^{*****}Department of Experimental Medicine, Section of Human Physiology and Unit of Dietetics and Sports Medicine, Department of Clinical and Experimental Medicine, University of Foggia, Foggia, Italy

**Lucia Parisi, Maria Antonietta Faraldo and Maria Ruberto equally contributed to the manuscript*

ABSTRACT

Introduction: The association between primary monosymptomatic nocturnal enuresis (PMNE) and behavioral disorders was highlighted frequently, suggesting psychiatric origin. On the other hand, there is no difference between the incidence of mental disorders in children with PMNE and controls, although the psychological genesis could provide additional secondary forms, in which the child might react to stressful events with the resumption of involuntary urination at night, with a possible alteration of bowel control due to a high vulnerability to stressors.

The purpose of this study is assessing the stressful events of life in a sample of children with PMNE.

Materials and methods: 56 subjects with PMNE, (37 males and 19 females) (mean 10.87 years; SD \pm 1.68) were recruited consecutively. All subjects were evaluated for the presence of stressful events (ICU) with test Coddington Life Events Scales (CLES). The control group consists of 98 typically developing children (65 males, 33 females) ($p = 0.855$) (mean 11.3 years; SD \pm 1.85; $p = 0.594$).

Results: Individuals with enuresis do not show a significant difference in the prevalence of stressful events than the control group (42.85% vs 41.83%; Chi-square = 0.002; $p = 0.963$) (table 1).

Conclusions: These results for the first time show that PMNE can be regarded as an independent state by stress factors, suggesting that it itself represents a stress factor that can affect the proper psychological and neuropsychological development in children.

Keywords: primary monosymptomatic nocturnal enuresis, PMNE; Coddington Life Events Scales, life adverse events.

DOI: 10.19193/0393-6384_2017_1_003

Received September 30, 2016; Accepted November 02, 2016

Introduction

The physiology of micturition is a complex system that involves three important organs, whose proper functionality is a key requirement: autonomic nervous system (SNA), kidneys and bladder.

The bladder control allows the alternation of long and slow phases of filling and emptying phases of short, in opposition with each other for reflex mechanism, with a rate of filling of the bladder of approximately 50 ml of urine/hour⁽¹⁻⁵⁾.

In pediatric age, Primary Monosymptomatic Nocturnal Enuresis (PMNE) occurs when the mic-

turition control falls due to one of the following three different factors: nocturnal polyuria; overactivity of the detrusor muscle; low arousability rate⁽¹⁻⁵⁾.

In the past decades, the association between PMNE and behavioral disorders was highlighted, leading researchers to believe that the disease may be due to a psychiatric origin, and linking the symptom to an altered parent-child relationship, based on the detection of pathological results achieved, both by mothers of children affected, in rating scales⁽¹⁻⁵⁾.

Subsequent studies have shown conflicting data with this theory, up to its complete abandonment. In fact, there is no difference between the incidence of mental disorders in children with PMNE and controls, although the psychological genesis could be envisaged more for secondary forms, in which the child would react to stressful events with the resumption of involuntary urination at night, with a possible alteration of the sphincter control linked to a 'high vulnerability to stressors'⁽¹⁻⁵⁾.

The aim of the present study is evaluating the stressful life events in a sample of children affected by PMNE.

Materials and methods

According to DSM-5 criteria, 56 subjects affected by PMNE, (37 males and 19 females) (mean age 10.87; SD \pm 1.68) were consecutively recruited.

All subjects were evaluated for the presence of stressful life events (LCU) with tests Coddington Life Events Scales (CLES). All subjects in both groups were recruited within the same urban area, all Caucasian and homogeneous for socioeconomic level.

Exclusion criteria were the following: mental retardation (IQ <70), borderline intellectual functioning, genetic syndromes (eg. Down syndrome, Prader-Willi syndrome, fragile X syndrome), hypothyroidism, metabolic and celiac diseases, psychiatric disorders (autism, schizophrenia, mood disorders, ADHD), movement and neuromuscular disorders, epilepsy, obesity, primary headache, sleep related breathing disorders, sleep troubles and pharmacological and/or complementary treatment⁽⁶⁻⁴⁴⁾.

All evaluations were performed after informed consent from the parents, and when

appropriate also by children. Control group was composed by 98 typical developing healthy children (65 males, 33 females) ($p = 0.855$) (mean age 11.3; SD \pm 1.85; $p = 0.594$).

Coddington life events scales (CLES)

Stressful LEs were measured using the Coddington Life Events Scales (CLES), which measure the occurrence of 53 LEs. Respondents indicate for each item describing a specific LE and the number of times the event has occurred in the last 3 months, 4-6 months earlier, 7-9 months earlier, or 10-12 months earlier. The frequency of occurrence is taken into account in the calculation of Life Change Units (LCUs) which also reflect the amount of stress inherent to the event and how long ago it happened. We used the original LCUs, which were obtained from ratings provided by teachers, paediatricians, and child psychiatrists. A total LCU score can be calculated for each respondent as a weighted sum of all the LCU scores (range of LCUs for one LE: 5-216).

We used the Italian version of CLES. Life events were classified according to two different typologies: desirable (e.g. "Graduating from high school") vs. undesirable events (e.g. "Divorce of parents") and family-related (e.g. "Loss of a job of your father or mother") vs. extra-family events (e.g. "Going on the first date"). Each LE was classified accordingly into one of the two categories in each typology, except nine which were classified in only one LE typology (e.g. "Being hospitalized for illness or injury" was undesirable but was not classified in the 'family' typology because it was neutral with regard to that particular typology).

Statistical analysis

The t-Student's test and Chi-square tests were applied to determine the differences among two groups. P values <0.05 were considered statistically significant. For statistical analysis it used the software STATISTICA (data analysis software system, version 6, StatSoft, Inc. (2001).

Results

The subjects with enuresis do not show a significant difference in the prevalence of stressful events than the control group (42.85% vs. 41.83%; Chi-square = 0.002; $p = 0.963$) (table 1).

	PMNE (n=56)	Controls (n=98)	<i>p</i>
Age	10.87 ± 1.68	11.3 ± 1.85	0.594
Sex (M/F)	37/19	65/33	0.855
z-BMI	0.71 ± 0.12	0.69 ± 0.18	0.459
Nocturnal enuresis frequency/week	7	-	-
Stressful events %	42.85	41.83	0.963

Table 1: shows comparisons among means and/or percentages among children affected by primary monosymptomatic nocturnal enuresis (PMNE) and healthy controls for age, sex distribution, z-BMI, and stressful events. t-Test and Chi-square test analysis were performed when appropriated. $p < 0.05$ values were considered as statistical significant.

Discussion

In the past decades scientific reports have extensively analyzed and confirmed the association between adverse life events during the past and/or during the last twelve months and the onset of psychopathological disorders in both preschool and school aged children and adolescents⁽⁴⁵⁻⁵⁰⁾.

Therefore, psychiatric/psychological distress disorder is currently regarded more and more as a secondary to PMNE and or as its comorbidities. Undoubtedly, PMNE can determine both social and personal difficulties in children affected with significant impact on emotional state, social development and self-esteem, as well as on the whole family⁽⁴⁵⁻⁵⁰⁾.

PMNE may be a devastating experience for children and young adults, inducing feelings of guilt, shame, embarrassment, sense of difference from others and a low self-esteem⁽⁴⁵⁻⁵⁰⁾.

PMNE children and adolescents tend to show pathological scores in the outsourcing stairs (aggression, attention deficit, which results in low academic performance), with a lower incidence of the internalization disorders than healthy controls⁽⁴⁵⁻⁵⁰⁾.

On the other hand, PMNE seems to occur as main comorbidity with other psychiatric disorders such as bipolar disorder, oppositional defiant and ADHD. However, not all children and adolescents who have experienced adverse life events will develop a psychopathological disorder. The event response is subjective and individual and depends on the combination of risk factors and protective factors. Pathogenesis of this relationship is not well studied although we could speculate the role of many different neurotransmitters⁽⁴⁵⁻⁵⁰⁾.

In conclusion, children affected by PMNE have not a higher prevalence of stressful events later in life compared to normal subjects. These results for the first time show that the PMNE may be considered as an independent state by stressors, suggesting that it itself represents a stressor that can affect the correct psychological and neuropsychological development in children.

References

- 1) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Mazzotta G, Carotenuto M. Primary nocturnal enuresis as a risk factor for sleep disorders: an observational questionnaire-based multicenter study. *Neuropsychiatr Dis Treat.* 2013; 9: 437-43. doi:10.2147/NDT.S43673.
- 2) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Mazzotta G, Patriciello G, Precenzano F, Carotenuto M. Visuomotor competencies and primary monosymptomatic nocturnal enuresis in prepubertal aged children. *Neuropsychiatr Dis Treat.* 2013; 9: 921-6. doi: 10.2147/NDT.S46772.
- 3) Carotenuto M, Esposito M, Pascotto A. Facial patterns and primary nocturnal enuresis in children. *Sleep Breath.* 2011 May; 15(2): 221-7. doi: 10.1007/s11325-010-0388-6.
- 4) Esposito M, Carotenuto M, Roccella M. Primary nocturnal enuresis and learning disability. *Minerva Pediatr.* 2011 Apr; 63(2): 99-104.
- 5) Carotenuto M, Esposito M, Pascotto A. Migraine and enuresis in children: An unusual correlation? *Med Hypotheses.* 2010 Jul;75(1):120-2. doi: 10.1016/j.mehy.2010.02.004.
- 6) Precenzano F, Lombardi P, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Della Valle I, Magliulo RM, Messina G, Roccella M. Internalizing symptoms in children affected by childhood absence epilepsy: a preliminary study. *Acta Medica Mediterranea*, 2016, 32: 1749; DOI: 10.19193/0393-6384_2016_6_158.
- 7) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Della Valle I, Visco G, Magliulo Rm, Messina G, Roccella M. ADHD-like symptoms in children affected by obstructive sleep apnea syndrome: a case-control study. *Acta Medica Mediterranea*, 2016, 32: 1755; DOI: 10.19193/0393-6384_2016_6_159.
- 8) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Grappa MF, Magliulo RM, Messina G, Roccella M. Borderline intellectual functioning and parental stress: an italian case-control study. *Acta Medica Mediterranea*, 2016, 32: 1761; DOI: 10.19193/0393-6384_2016_6_160.

- 9) Ruberto M, Precenzano F, Parisi L, Salerno M, Maltese A, Messina G, Roccella M. Visuomotor integration skills in children affected by obstructive sleep apnea syndrome: a case-control study. *Acta Medica Mediterranea*, 2016, 32: 1659; DOI: 10.19193/0393-6384_2016_5_146.
- 10) Parisi L, Ruberto M, Precenzano F, Di Filippo T, Russotto C, Maltese A, Salerno M, Roccella M. The quality of life in children with cerebral palsy. *Acta Medica Mediterranea*, 2016, 32: 1665; DOI: 10.19193/0393-6384_2016_5_147.
- 11) Parisi L, Di Filippo T, Roccella M. The child with autism spectrum disorders (ASD): behavioral and neurobiological aspects. *Acta Medica Mediterranea*, 2015, 31: 1187.
- 12) Esposito M, Gallai B, Roccella M, Marotta R, Lavano F, Lavano SM, Mazzotta G, Bove D, Sorrentino M, Precenzano F, Carotenuto M. Anxiety and depression levels in prepubertal obese children: a case-control study. *Neuropsychiatr Dis Treat*. 2014 Oct 3; 10: 1897-902. doi: 10.2147/NDT.S69795.
- 13) Carotenuto M, Esposito M, Parisi L, Gallai B, Marotta R, Pascotto A, Roccella M. Depressive symptoms and childhood sleep apnea syndrome. *Neuropsychiatr Dis Treat*. 2012; 8: 369-73. doi: 10.2147/NDT.S35974.
- 14) Perillo L, Esposito M, Caprioglio A, Attanasio S, Santini AC, Carotenuto M. Orthodontic treatment need for adolescents in the Campania region: the malocclusion impact on self-concept. *Patient Prefer Adherence*. 2014 Mar 19; 8: 353-9. doi: 10.2147/PPA.S58971.
- 15) Carotenuto M, Esposito M, Di Pasquale F, De Stefano S, Santamaria F. Psychological, cognitive and maternal stress assessment in children with primary ciliary dyskinesia. *World J Pediatr*. 2013 Nov; 9(4): 312-7. doi:10.1007/s12519-013-0441-1.
- 16) Esposito M, Parisi L, Gallai B, Marotta R, Di Dona A, Lavano SM, Roccella M, Carotenuto M. Attachment styles in children affected by migraine without aura. *Neuropsychiatr Dis Treat*. 2013; 9: 1513-9. doi: 10.2147/NDT.S52716.
- 17) Esposito M, Roccella M, Gallai B, Parisi L, Lavano SM, Marotta R, Carotenuto M. Maternal personality profile of children affected by migraine. *Neuropsychiatr Dis Treat*. 2013; 9: 1351-8. doi: 10.2147/NDT.S51554.
- 18) Perillo L, Esposito M, Contiello M, Lucchese A, Santini AC, Carotenuto M. Occlusal traits in developmental dyslexia: a preliminary study. *Neuropsychiatr Dis Treat*. 2013; 9: 1231-7. doi: 10.2147/NDT.S49985.
- 19) Esposito M, Marotta R, Gallai B, Parisi L, Patriciello G, Lavano SM, Mazzotta G, Roccella M, Carotenuto M. Temperamental characteristics in childhood migraine without aura: a multicenter study. *Neuropsychiatr Dis Treat*. 2013;9:1187-92. doi:10.2147/NDT.S50458.
- 20) Esposito M, Gallai B, Parisi L, Castaldo L, Marotta R, Lavano SM, Mazzotta G, Roccella M, Carotenuto M. Self-concept evaluation and migraine without aura in childhood. *Neuropsychiatr Dis Treat*. 2013; 9: 1061-6. doi: 10.2147/NDT.S49364.
- 21) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Gritti A, Mazzotta G, Carotenuto M. Maternal stress and childhood migraine: a new perspective on management. *Neuropsychiatr Dis Treat*. 2013; 9: 351-5. doi:10.2147/NDT.S42818.
- 22) Esposito M, Ruberto M, Pascotto A, Carotenuto M. Nutraceutical preparations in childhood migraine prophylaxis: effects on headache outcomes including disability and behaviour. *Neurol Sci*. 2012 Dec; 33(6): 1365-8. doi: 10.1007/s10072-012-1019-8.
- 23) Carotenuto M, Esposito M, Precenzano F, Castaldo L, Roccella M. Cosleeping in childhood migraine. *Minerva Pediatr*. 2011 Apr; 63(2): 105-9.
- 24) Guzzetta A, Pizzardi A, Belmonti V, Boldrini A, Carotenuto M, D'Acunto G, Ferrari F, Fiori S, Gallo C, Ghirri P, Mercuri E, Romeo D, Roversi MF, Cioni G. Hand movements at 3 months predict later hemiplegia in term infants with neonatal cerebral infarction. *Dev Med Child Neurol*. 2010 Aug; 52(8): 767-72. doi:10.1111/j.1469-8749.2009.03497.x.
- 25) Carotenuto M, Santoro N, Grandone A, Santoro E, Pascotto C, Pascotto A, Perrone L, del Giudice EM. The insulin gene variable number of tandem repeats (INS-VNTR) genotype and sleep disordered breathing in childhood obesity. *J Endocrinol Invest*. 2009 Oct; 32(9): 752-5. doi: 10.3275/6398.
- 26) Esposito M, Carotenuto M. Borderline intellectual functioning and sleep: the role of cyclic alternating pattern. *Neurosci Lett*. 2010 Nov 19; 485(2): 89-93. doi:10.1016/j.neulet.2010.08.062.
- 27) Carotenuto M, Esposito M, D'Aniello A, Ripa CD, Precenzano F, Pascotto A, Bravaccio C, Elia M. Polysomnographic findings in Rett syndrome: a case-control study. *Sleep Breath*. 2013 Mar; 17(1): 93-8. doi: 10.1007/s11325-012-0654-x.
- 28) Esposito M, Antinolfi L, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Mazzotta G, Precenzano F, Carotenuto M. Executive dysfunction in children affected by obstructive sleep apnea syndrome: an observational study. *Neuropsychiatr Dis Treat*. 2013; 9: 1087-94. doi: 10.2147/NDT.S47287.
- 29) Esposito M, Parisi P, Miano S, Carotenuto M. Migraine and periodic limb movement disorders in sleep in children: a preliminary case-control study. *J Headache Pain*. 2013 Jul 1; 14: 57. doi: 10.1186/1129-2377-14-57.
- 30) Carotenuto M, Gimigliano F, Fiordelisi G, Ruberto M, Esposito M. Positional abnormalities during sleep in children affected by obstructive sleep apnea syndrome: the putative role of kinetic muscular chains. *Med Hypotheses*. 2013 Aug; 81(2): 306-8. doi: 10.1016/j.mehy.2013.04.023.
- 31) Carotenuto M, Esposito M. Nutraceuticals safety and efficacy in migraine without aura in a population of children affected by neurofibromatosis type I. *Neurol Sci*. 2013 Nov; 34(11): 1905-9. doi: 10.1007/s10072-013-1403-z.
- 32) Morandi A, Bonnefond A, Lobbens S, Carotenuto M, Del Giudice EM, Froguel P, Maffei C. A girl with incomplete Prader-Willi syndrome and negative MS-PCR, found to have mosaic maternal UPD-15 at SNP array. *Am J Med Genet A*. 2015 Nov; 167A(11): 2720-6. doi: 10.1002/ajmg.a.37222.
- 33) Esposito M, Carotenuto M. Intellectual disabilities and power spectra analysis during sleep: a new perspective on borderline intellectual functioning. *J Intellect Disabil Res*. 2014 May; 58(5): 421-9. doi: 10.1111/jir.12036.

- 34) Carotenuto M, Gallai B, Parisi L, Roccella M, Esposito M. Acupressure therapy for insomnia in adolescents: a polysomnographic study. *Neuropsychiatr Dis Treat*. 2013; 9: 157-62. doi: 10.2147/NDT.S41892.
- 35) Esposito M, Pascotto A, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Gritti A, Mazzotta G, Carotenuto M. Can headache impair intellectual abilities in children? An observational study. *Neuropsychiatr Dis Treat*. 2012; 8: 509-13. doi:10.2147/NDT.S36863.
- 36) Esposito M, Verrotti A, Gimigliano F, Ruberto M, Agostinelli S, Scuccimarra G, Pascotto A, Carotenuto M. Motor coordination impairment and migraine in children: a new comorbidity? *Eur J Pediatr*. 2012 Nov; 171(11): 1599-604. doi:10.1007/s00431-012-1759-8.
- 37) Carotenuto M, Esposito M, Cortese S, Laino D, Verrotti A. Children with developmental dyslexia showed greater sleep disturbances than controls, including problems initiating and maintaining sleep. *Acta Paediatr*. 2016 Sep; 105(9): 1079-82. doi: 10.1111/apa.13472.
- 38) Matricardi S, Spalice A, Salpietro V, Di Rosa G, Balistreri MC, Grosso S, Parisi P, Elia M, Striano P, Accorsi P, Cusmai R, Specchio N, Coppola G, Savasta S, Carotenuto M, Tozzi E, Ferrara P, Ruggieri M, Verrotti A. Epilepsy in the setting of full trisomy 18: A multicenter study on 18 affected children with and without structural brain abnormalities. *Am J Med Genet C Semin Med Genet*. 2016 Sep; 172(3): 288-95. doi: 10.1002/ajmg.c.31513.
- 39) Pasquali D, Carotenuto M, Leporati P, Esposito M, Antinolfi L, Esposito D, Accardo G, Carella C, Chiovato L, Rotondi M. Maternal hypothyroidism and subsequent neuropsychological outcome of the progeny: a family portrait. *Endocrine*. 2015 Dec; 50(3): 797-801. doi: 10.1007/s12020-015-0564-3.
- 40) Santamaria F, Esposito M, Montella S, Cantone E, Mollica C, De Stefano S, Mirra V, Carotenuto M. Sleep disordered breathing and airway disease in primary ciliary dyskinesia. *Respirology*. 2014 May; 19(4):570-5. doi: 10.1111/resp.12273.
- 41) Esposito M, Ruberto M, Gimigliano F, Marotta R, Gallai B, Parisi L, Lavano SM, Roccella M, Carotenuto M. Effectiveness and safety of Nintendo Wii Fit Plus™ training in children with migraine without aura: a preliminary study. *Neuropsychiatr Dis Treat*. 2013; 9: 1803-10. doi: 10.2147/NDT.S53853.
- 42) Di Filippo T, Orlando MF, Concialdi G, La Grutta S, Lo Baido R, Epifanio MS, Esposito M, Carotenuto M, Parisi L, Roccella M. The quality of life in developing age children with celiac disease. *Minerva Pediatr*. 2013 Dec; 65(6): 599-608.
- 43) Esposito M, Gimigliano F, Barillari MR, Precenzano F, Ruberto M, Sepe J, Barillari U, Gimigliano R, Militerni R, Messina G, Carotenuto M. Pediatric selective mutism therapy: a randomized controlled trial. *Eur J Phys Rehabil Med*. 2016 Nov 10. [Epub ahead of print] PubMed PMID: 27830922.
- 44) Esposito M, Gimigliano F, Ruberto M, Marotta R, Gallai B, Parisi L, Lavano SM, Mazzotta G, Roccella M, Carotenuto M. Psychomotor approach in children affected by nonretentive fecal soiling (FNRFs): a new rehabilitative purpose. *Neuropsychiatr Dis Treat*. 2013; 9: 1433-41. doi: 10.2147/NDT.S51257.
- 45) Messina A, De Fusco C, Monda V, Esposito M, Moscatelli F, Valenzano A, Carotenuto M, Viggiano E, Chieffi S, De Luca V, Cibelli G, Monda M, Messina G. Role of the Orexin System on the Hypothalamus-Pituitary-Thyroid Axis. *Front Neural Circuits*. 2016 Aug 25; 10: 66. doi: 10.3389/fncir.2016.00066.
- 46) Moscatelli F, Valenzano A, Petito A, Ivano Triggiani A, Anna Pia Ciliberti M, Luongo L, Carotenuto M, Esposito M, Messina A, Monda V, Monda M, Capranica L, Messina G, Cibelli G. Relationship between blood lactate and cortical excitability between taekwondo athletes and non-athletes after hand-grip exercise. *Somatosens Mot Res*. 2016 Jun; 33(2): 137-44. doi: 10.1080/08990220.2016.1203305.
- 47) Entesar Foumany GhH, Salehi J. The relationship between emotional intelligence and life satisfaction and the mediatory role of resiliency and emotional balance among the students of zanjaan university. *Acta Medica Mediterranea*, 2015, 31: 1351.
- 48) Nabi Amjad R, Navab E, Nikbakht Nasrabad A. Parents of children with epilepsy captured by epilepsy: a qualitative study. *Acta Medica Mediterranea*, 2016, 32: 1303.
- 49) Shams S, Omolbanin Mohammadian S, Monajemzadeh M, Emamgholipour S, Aghi Haghi Ashtiani M, Irani H, Shafeghat L. Evaluation of serum transferrin receptors in children with iron deficiency anemia. *Acta Medica Mediterranea*, 2016, 32: 1555.
- 50) Yongli Z, Chunting L. Problems and countermeasures of pediatric emergency nursing security. *Acta Medica Mediterranea*, 2016, 32: 1177.

Corresponding author
MICHELE ROCCELLA, MD, PhD
Department of Psychological
Pedagogical and Educational Sciences
University of Palermo
(Italy)