


AUTHOR QUERY FORM

| | | |
|---|--|--|
|  | Journal: EPSL Article Number: 13747 | Please e-mail your responses and any corrections to: E-mail: corrections.essd@elsevier.vtex.lt |
|---|--|--|

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'Q' link to go to the location in the proof.

| Location in article | Query / Remark: click on the Q link to go Please insert your reply or correction at the corresponding line in the proof |
|---------------------|--|
| Q1 | Your article is registered as a regular item and is being processed for inclusion in a regular issue of the journal. If this is NOT correct and your article belongs to a Special Issue/Collection please contact <k.stair@elsevier.com> immediately prior to returning your corrections. (p. 1/ line 1) |
| Q2 | The author names have been tagged as given names and surnames (surnames are highlighted in teal color). Please confirm if they have been identified correctly and are presented in the desired order. (p. 1/ line 15) |
| Q3 | Please indicate which author(s) should be marked as 'Corresponding author'. If the corresponding author has two or more affiliations please specify the address for correspondence. (p. 1/ line 16) |
| Q4, Q5, Q6, Q7 | Figs. 1-3, 5 will appear in black and white in print and in color on the web. The captions of these figures contain references to color or the colors are mentioned in the main text. Based on this, the respective figure captions have been updated. Please check, and correct if necessary. (p. 2/ line 34; p. 3/ line 21; p. 5/ line 26; p. 6/ line 22) |
| Q8 | Please check if sponsor names have been identified correctly and correct if necessary. (p. 12/ line 5) |
| Q9 | Please correct verb-subject agreement in the sentence 'Thus, the...for eruption.'. (p. 1/ line 105) |
| Q10 | Reference(s) given here were noted in the reference list but are missing from the text - please position each reference in the text or delete it from the list. All references in the main reference list must be cited in the main text. If the uncited reference appears only in the supplementary material, please either (1) move the supplemental reference to the supplemental file and send the corrected supplemental file, or (2) provide approval to move the uncited reference to a "Further Reading" section. (p. 9/ line 115) |
| Q11, Q12, Q13 | Please provide grant numbers for the sponsors 'Deep Carbon Observatory Deep Earth Carbon Degassing program', 'CNE Costa Rica', 'USAID' (if available) and include this number into the main text where appropriate. (p. 9/ line 92,95,96) |

| Location in article | Query / Remark: click on the Q link to go Please insert your reply or correction at the corresponding line in the proof |
|--------------------------------|---|
| | <div data-bbox="646 140 1188 245" style="border: 1px solid black; padding: 5px; width: fit-content; margin: auto;"><p data-bbox="667 163 1057 224">Please check this box if you have no corrections to make to the PDF file</p><input data-bbox="1089 163 1167 224" type="checkbox"/></div> |

1Q1

2

3

4

5

6

7

8

9

10

11

12

13

14

15Q2

16Q3

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66



ELSEVIER

Contents lists available at ScienceDirect

Earth and Planetary Science Letters

www.elsevier.com/locate/epsl



Short-period volcanic gas precursors to phreatic eruptions: Insights from Poás Volcano, Costa Rica

J.M. de Moor^{a,b,c}, A. Aiuppa^{c,d}, J. Pacheco^a, G. Avard^a, C. Kern^e, M. Liuzzo^d,
M. Martínez^a, G. Giudice^d, T.P. Fischer^b

^a Observatorio Vulcanológico y Sismológico de Costa Rica, Universidad Nacional, Heredia, Costa Rica

^b Department of Earth and Planetary Sciences, University of New Mexico, Albuquerque, NM, USA

^c Dipartimento DiSTeM, Università di Palermo, Palermo, Italy

^d Istituto Nazionale di Geofisica e Vulcanologia, Sezione di Palermo, Italy

^e Cascades Volcano Observatory, United States Geological Survey, Vancouver, WA, USA

ARTICLE INFO

Article history:

Received 6 November 2015

Received in revised form 19 February 2016

Accepted 29 February 2016

Available online xxxx

Editor: T.A. Mather

Keywords:

volcanic gas

phreatic eruption

eruption precursor

volcanic lake

hydrothermal system

Poás volcano

ABSTRACT

Volcanic eruptions involving interaction with water are amongst the most violent and unpredictable geologic phenomena on Earth. Phreatic eruptions are exceptionally difficult to forecast by traditional geophysical techniques. Here we report on short-term precursory variations in gas emissions related to phreatic blasts at Poás volcano, Costa Rica, as measured with an in situ multiple gas analyzer that was deployed at the edge of the erupting lake. Gas emitted from this hyper-acid crater lake approaches magmatic values of SO₂/CO₂ 1–6 days prior to eruption. The SO₂ flux derived from magmatic degassing through the lake is measureable by differential optical absorption spectrometry (sporadic campaign measurements), which allows us to constrain lake gas output and input for the major gas species during eruptive and non-eruptive periods. We can further calculate power supply to the hydrothermal system using volatile mass balance and thermodynamics, which indicates that the magmatic heat flux into the shallow hydrothermal system increases from ~27 MW during quiescence to ~59 MW during periods of phreatic events. These transient pulses of gas and heat from the deeper magmatic system generate both phreatic eruptions and the observed short-term changes in gas composition, because at high gas flux scrubbing of sulfur by the hydrothermal system is both kinetically and thermodynamically inhibited whereas CO₂ gas is always essentially inert in hyperacid conditions. Thus, the SO₂/CO₂ of lake emissions approaches magmatic values as gas and power supply to the sub-limnic hydrothermal system increase, vaporizing fluids and priming the hydrothermal system for eruption. Our results suggest that high-frequency real-time gas monitoring could provide useful short-term eruptive precursors at volcanoes prone to phreatic explosions.

© 2016 Published by Elsevier B.V.

1. Introduction

Volcanic eruptions involving explosive vaporization of meteoric water are common occurrences on our wet planet. Of the ~18,000 volcanic eruptions listed in the Global Volcanism Database since 1900, 822 (~5%) are considered phreatic in nature (Global Volcanism Program, 2013). Eruptions that occur through surface water account for ~8% of historic eruptions but have claimed a disproportionate number of human lives (~20% of total deaths) (Mastin and Witter, 2000). The diverse hazards related to wet eruptions contribute to their deadliness, and include base surges, ballistics,

lahars, tsunamis, and toxic gases. Lahars generated by explosive ejection of crater lake water are the deadliest hazard associated with volcanic lakes and have killed >16,000 people in historical times (Mastin and Witter, 2000).

Phreatic eruptions are exceptionally challenging to forecast because many do not directly involve the movement of magma, which generates distinct geophysical signals useful in eruption forecasting, and may occur simply due to subtle changes in shallow hydrothermal systems (Rouwet et al., 2014a). Documented precursors of phreatic activity show long to medium term changes in geophysical (i.e. seismicity, deformation, and gravimetry) or geochemical (i.e. lake water and gas compositions) parameters (Sano et al., 2015; Barberi et al., 1992; Rouwet et al., 2014b), but accurate short-term (days) eruption forecasting remains exception-

E-mail address: maartenjdemoor@gmail.com (J.M. de Moor).

<http://dx.doi.org/10.1016/j.epsl.2016.02.056>

0012-821X/© 2016 Published by Elsevier B.V.




Fig. 1. The Poás crater and general characteristics of gas emissions. Multi-GAS measurement sites (Lake and Dome Fumaroles), the CRPO seismic station, the webcam, and DOAS traverse path are indicated. Webcam images show that the fumarolic plume rises buoyantly and travels to the SW for >95% of the time (Supplementary Materials; Fig. S6; Movie S2). The gas plume from the lake is diffuse and transparent, and rises buoyantly to the NE and NW (Movie S2). Inserts A and B show example Multi-GAS datasets collected on 11 March 2014 during a period of low eruptive activity and on 16 May 2014 during a period of intense phreatic activity, respectively. The fumarolic (red diamonds) and lake gas (yellow circles) compositions are clearly distinguishable on 11 March, but similar on 16 May. Insert C shows example DOAS traverses (column amounts in ppm·m) in which lake and fumarole gas plumes are distinguished. The traverse in gray was conducted on 30 September 2014, 5 days before a climax in phreatic eruptions, and shows a strong SO₂ plume from the lake. The traverse in black was conducted on October 16, after the climax and at the beginning of a period with no eruptions and shows a much weaker lake SO₂ plume. Background image from Google Earth. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

ally challenging. For example, at some well-monitored volcanoes with crater lakes, precursory seismicity has been observed to occur just minutes before phreatic eruptions (Kaneshima et al., 1996; Jolly et al., 2010) or is entirely absent (Christenson et al., 2007).

Published cases of precursory changes in gas geochemistry prior to phreatic eruptions are rare. This is probably because most attention from the scientific community has focused on volcanoes prone to large magmatic eruptions. However, the 27 September 2014 eruption at Ontake-San Volcano (Japan) claimed 57 victims (with 6 still missing), highlighting the need for more focused research on this relatively poorly studied class of volcanic activity and the wet volcanoes where frequent phreatic activity can dominate restless behavior, without apparent movement or eruption of magma. Recently, Sano et al. (2015) reported a decadal-scale change in He isotope compositions towards more magmatic values at Ontake prior to the fatal phreatic blast. White Island in New Zealand is a well-monitored volcano with an ephemeral crater lake, high temperature fumaroles, and is prone to phreatic eruptions. Werner et al. (2008) recognized large changes in bulk plume gas chemistry over a period of years (2002–2006) after the formation of a crater lake in 2003. Both SO₂ flux and SO₂/CO₂ approached magmatic values and were accompanied by a drastic increase in seismic activity; however no eruptions directly followed the escalation. Ruapehu is another well-monitored volcano in New Zealand that hosts an active crater lake. Airborne gas monitoring revealed that SO₂ flux and SO₂/CO₂ both increased in the year to months prior to the 25 September 2007 phreatomagmatic eruption (Christenson et al., 2010). Nevado del Ruiz volcano in Columbia also had a highly active crater lake before the 1985 Plinian eruption. Giggenbach et al. (1990) considered this eruption to be between magmatic and phreatic in character, because the blast ejected only older mag-

matic rocks (Williams et al., 1986) and was driven by magmatic gas pressure buildup below a sealed carapace produced by the sublimic hydrothermal system. The expulsion of lake water and melting of summit ice during this “pneumato-magmatic” (Giggenbach et al., 1990) eruption generated an acidic lahar that killed >20,000 people from the town of Armero. In contrast to the cases at White Island (Werner et al., 2008) and Ruapehu (Christenson et al., 2010), where gases became more magmatic with increasing activity, gas compositions at Nevado del Ruiz became more hydrothermal in composition a few months before the eruption (Giggenbach et al., 1990). These studies suggest that decadal to monthly scale precursory changes in gas compositions likely occur at many volcanoes prone to phreatic (and related) eruptions. However, the cause of the changes and the specific conditions triggering sudden and potentially deadly phreatic explosions not associated with magmatic movement are poorly understood. The recognition of associated short-term changes in gas chemistry is essentially absent for these complex magmatic-hydrothermal systems and their eruptions, but could be essential in forecasting individual explosive blasts.

Our study focuses on Poás volcano (N 10.1968°; W 84.2305°) in Costa Rica, which entered a period of phreatic activity in 2006. The eruptive period culminated in a series of relatively large phreatic blasts (columns >300 m) between June and October of 2014, after which the volcano entered a period of repose that continues until present time. Poás volcano hosts a warm acidic lake (Laguna Caliente; $T \sim 50^\circ\text{C}$, $\text{pH} \sim 0$ at the time of study) of about 270 m diameter, as well as high-temperature (up to $\sim 800^\circ\text{C}$) fumaroles adjacent to the lake (Dome Fumaroles; Fig. 1). The Poás crater represents one of the most chemically extreme environments on Earth and Poás Volcano National Park was visited by >200,000 tourists in 2014. About 60 seismically registered phreatic blasts oc-




Fig. 2. Time series lake gas SO_2/CO_2 and SE of eruptions. The yellow dashed line is the median of lake gas SO_2/CO_2 (1.64). Total error in Multi-GAS measurements is conservatively estimated at 20% from experimental results (± 0.33 for SO_2/CO_2 of 1.64; see Methods), whereas the average reproducibility of the lake gas measurements is 0.18 (error bars shown at left). The composition of the dome fumaroles was measured on 16 May, and the fumarolic compositional trend is constrained by Multi-GAS campaigns between January and July of 2014 (Fig. S3). The sizes of the blue diamonds correspond to phreatic eruption groups as defined by SE (Fig. 3). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

curred from the lake during the same year, ranging from minor “gas bursts” to highly explosive jets ejecting ballistics, sediments, vapor and lake water to >400 m above the lake surface (Movie S1). Poás explosions are a type example of phreatic eruptions through a crater lake, displaying a vertical jet of solids and fluids accompanied by a radial base surge of vapor (Fig. S1, Movie S1), and juvenile magmatic material is erupted. Short-term geophysical precursors heralding an imminent eruption are absent. Long-term (2001 to 2014) and sporadic fumarole gas chemistry monitoring recently showed how the dynamics of the volcano-hydrothermal system are reflected in changes in gas discharges (Williams et al., 1986). Here we focus on the short-term (days to weeks) variations in plume gas compositions that provide new insights on the processes that lead to phreatic eruptions.

2. Materials and methods

2.1. Field methods

We conducted an experiment to assess high-frequency temporal variations in lake gas composition at Poás from April to June 2014, which was a period of intense phreatic activity. A fully autonomous Multi-Component Gas Analyzer System (Multi-GAS) (Giggenbach et al., 1990) was installed at the shore of the lake (Fig. 1). The fixed “lake” Multi-GAS instrument records concentrations of CO_2 , SO_2 , H_2S , and H_2O at a rate of 0.1 Hz. The instrument is programmed to acquire data for 30 min four times a day, which allowed us to derive a time-series of gas ratios (Fig. 2, Table S1) that are independent of mixing with air. Gas ratios are determined by linear regression through gas concentration data (Williams et al., 1986), as shown in Fig. S2. The site of the lake Multi-GAS station was carefully chosen to target pure lake gas emissions, and was located in a sheltered cove on the upwind side of the lake, where lake gas emissions tend to form a local eddy below a ~ 10 m cliff (see Supplementary Text). Data gaps in the time series are due to conditions in which not enough gas reached the station for robust determination of SO_2/CO_2 (i.e. SO_2 concentrations <5 ppm) or where a programmed analysis session failed due to lack of charge on the battery (i.e. persistent low light conditions, or solar panel damaged by 2 June eruption). The near-real time data from this fixed lake station (Fig. 2; Table S1) were supplemented with sporadic campaign measurements conducted with a portable Multi-GAS to characterize the chemical composition of the dome fumaroles (Fig. 1; Fig. S3).

Fourteen seismically registered eruptions occurred during the 44 days of data acquisition by the lake Multi-GAS station. The largest phreatic eruption of 2014 occurred on June 2, ejecting ballistics and sediments to >400 m height, which destroyed peripheral components of the lake station. The station continued to record data until June 12, at which point the battery was drained. During campaign measurements, fumarole temperature was measured with infrared camera from the eastern lake shore, and walking traverses with ultraviolet spectrometers for Differential Optical Absorption Spectroscopy (DOAS) determination of SO_2 emission rate (Fischer et al., 2015) were conducted along the western crater rim (Fig. 1). The lake plume is distinguishable from the fumarolic plume by DOAS under typical weather conditions (Fig. 1 inset C).

2.2. Uncertainties in multi-GAS measurements

Errors in Multi-GAS measurements of gas ratios are paramount in determining whether the time series trends in SO_2/CO_2 (Fig. 2) are confidently resolved. As we are also comparing data collected with two different instruments, it is also necessary to take into account errors associated with each instrument. The overall uncertainty (combined accuracy and precision) in SO_2/CO_2 for the type of Multi-GAS instruments was reported to be $\leq 20\%$ by Aiuppa et al. (2007). Based on laboratory experiments, Aiuppa et al. (2013) reported the Multi-GAS error in SO_2/CO_2 to be $\leq 15\%$.

We assess analytical errors associated with Multi-GAS measurements in two ways. First, we conducted a series of experiments using gas mixtures of known SO_2/CO_2 to assess the total error (accuracy and reproducibility) of both of our Multi-GAS stations (see Supplementary Text, Figs. S4 and S5, Tables S2 and S3). The gas mixtures have a range in composition similar to those observed at Poás, from 2.54 to 0.16, and the experiments were conducted in ambient outdoor conditions in Heredia, Costa Rica, to mimic field conditions. These results confirm that total error is better than 20% (at 95% confidence level) for SO_2/CO_2 for both instruments (Supplementary Text, Figs. S4 and S5, Tables S2 and S3).

Of fundamental importance to this study is whether the trends in SO_2/CO_2 shown in Fig. 2 are resolved with confidence. In this regard, instrumental precision or reproducibility is of paramount importance. Thus, the second assessment of errors was conducted using the acquired time series field data by calculating multiple SO_2/CO_2 values for each 30 minute data acquisition period to quantify reproducibility (Table S1). Each analysis includes subtraction of background CO_2 signal (i.e. from air), and optimization

of the R^2 of linear regression, typically using >30 individual gas CO_2 and SO_2 concentration data points. SO_2/CO_2 values were calculated by multiple linear regression determinations on subsets of data within each half hour analytical session, allowing quantification of reproducibility for many of the individual datapoints shown in Fig. 2. This approach indicates that the average SO_2/CO_2 reproducibility is 0.18 (or 11% of the median observed SO_2/CO_2 value; See Supplementary Text for further details). The measured variations in SO_2/CO_2 are far greater than the 20% error margin or the precision/reproducibility as determined by multiple linear regression assessments within datasets. We also note that the 20% analytical errors margins are greater than the precision values in the vast majority (86%) of cases, suggesting that the gases measured by the lake Multi-GAS station are **homogeneous** on very short timescales (i.e. within the 30 min analysis interval).

2.3. Intercontamination of gas sources

Changes in environmental conditions can also affect the robustness of Multi-GAS measurements. For our purposes, the most important effect to consider is that of wind direction and the possibility of intercontamination between lake and fumarolic gas (Galle et al., 2002). The measurements of fumarolic gas composition were conducted manually and in close proximity to the source, allowing the operator to accurately target gas originating from the fumaroles. Mobile Multi-GAS measurements of the fumarolic gas are conducted on top of the dome to the south of the lake (Fig. 1, Movies S1 and S2) and the high gas concentration and poor visibility (a full face gas mask required) in the fumarolic plume make it exceedingly obvious to the operator when the Multi-GAS is measuring true fumarolic gas.

In the field it is also visually obvious that the fumarolic gas plume does not usually travel close to the site of Lake Multi-GAS station, however it is less trivial to quantitatively prove that our fixed Multi-GAS station lake gas compositions are not affected by mixing with fumarolic gas. We can however track the fumarolic plume direction using webcam footage, as is shown in Fig. S6 and Movie S1. The wind direction at Poás is almost always towards the west and based on assessment of all of the available webcam footage recorded during the deployment of the lake Multi-GAS instrument, we find that the fumarolic plume was **traveling** towards the west or southwest (away from the Multi-GAS station) for 97.6% of the time. The plume **traveled** towards the camera (north) for 1.7% of the time, and east (towards the Multi-GAS) for 0.7% of the time. When the fumarolic plume does travel towards the east, it is when there is very little wind and the plume rises buoyantly above the Lake Multi-GAS station. In none of the webcam footage have we observed a time period during a Multi-GAS analysis (i.e. the 2 h per 24 h that the instrument was actually acquiring data) where the fumarolic plume was **traveling** towards the lake Multi-GAS station.

Our second test for fumarole gas contamination is based on the difference in $\text{H}_2\text{S}/\text{SO}_2$ between the fumarolic and lake gas compositions. No H_2S has been detected in the lake emissions during mobile Multi-GAS measurements or by the fixed lake station during the period of study, whereas the fumarolic gas contains H_2S well above the detection limit of the Multi-GAS. This distinction between fumarolic and ultra-acidic lake gas emissions has also been observed at Aso volcano, Japan (Galle et al., 2002), and at Copahue volcano, Argentina (Aiuppa et al., 2009). However, the H_2S detection limit for Multi-GAS instruments is not trivial to determine precisely, and is dependent on the cross sensitivity of the H_2S electrochemical sensor to SO_2 gas. We determined the cross sensitivity during Multi-GAS calibrations by introducing H_2S -free gas with known SO_2 concentration to the instrument and measuring the apparent concentration of H_2S . For both of our instruments,

the cross sensitivity of the H_2S sensor to SO_2 is $20 \pm 2\%$, thus at 50 ppm SO_2 (roughly the highest concentrations observed in the Lake Multi-GAS emission data), the apparent H_2S concentration is 10 ± 1 ppm. After correction for SO_2 cross sensitivity (H_2S concentration = H_2S apparent concentration - $20 \pm 2\%$ SO_2 concentration), we observe no discernable H_2S , with an estimated detection limit of 1 ppm.

Based on SO_2 concentrations in lake Multi-GAS measurements and the $\text{H}_2\text{S}/\text{SO}_2$ values measured in the fumarolic gas plume (average of 0.12) at Poás, we can assess the expected H_2S concentrations if the gas measured by the lake Multi-GAS station originating from the fumaroles. This analysis shows that for 21 lake Multi-GAS measurements where high SO_2 concentrations and high SO_2/CO_2 values are observed, H_2S concentrations up to 5.9 ppm would be expected if the measured gas were fumarolic. The maximum H_2S concentrations observed in the lake Multi-GAS data are ≤ 1 ppm, which is below the detection limit. Therefore, we find no compositional evidence that the high observed SO_2/CO_2 values measured in the lake gas emissions are due to fumarolic gas from the dome **traveling** to the fixed Lake Multi-GAS site.

3. Results

The results of the lake Multi-GAS experiment show that significant short-period changes occur in lake gas emission composition. These variations are correlated with eruptive activity. Fig. 2 shows the SO_2/CO_2 (molar) as a function of time in Poás crater lake emissions as measured by the lake Multi-GAS station between April and June 2014 (Table S1). Even considering our conservative estimations of error, the time series trends in SO_2/CO_2 measured by the lake Multi-GAS are statistically significant (Spearman's rho tests on monotonic data segments indicate r values >0.9) and the variations are well beyond reasonable estimations of total error. Phreatic eruptions are plotted with their associated radiated seismic energy (SE, in units of kJ), which is used throughout as a relative measure of eruptive magnitude only (Table S4; see Supplementary Information). Our dataset, the first high-frequency time series of gas compositions acquired during phreatic eruptions, demonstrates that explosions occur after measurable short-term increases in SO_2/CO_2 . Importantly, the composition of lake gas emissions approaches, but does not surpass the SO_2/CO_2 value of the high temperature fumaroles (450°C – 750°C) and all of the largest eruptions ($\text{SE} > 30$) are preceded by trends of increasing SO_2/CO_2 starting two to six days beforehand. Our values of SO_2/CO_2 for the fumaroles are similar to the ratios measured in the highest temperature direct gas samples from Poás (Pering et al., 2014) and are consistent with regional trends in SO_2/CO_2 (Shinohara et al., 2015).

In order to statistically characterize the relationship between eruptions and lake gas composition, we initially consider the proportion of eruptions associated with SO_2/CO_2 values above the median SO_2/CO_2 value (1.64) of the lake Multi-GAS dataset to those associated with SO_2/CO_2 below the median. Considering the SO_2/CO_2 measurement closest in time to the occurrence of each eruption (Table S5), all but one of the fourteen eruptions have $\text{SO}_2/\text{CO}_2 \geq 1.64$. If we define the SO_2/CO_2 of each eruption as the mean value of gas compositions recorded within 12 h of the eruption time, thirteen of the fourteen eruptions have $\text{SO}_2/\text{CO}_2 > 1.64$, and for a 24 h time window, eleven eruptions have associated $\text{SO}_2/\text{CO}_2 > 1.64$ (Fig. 3). Thus, phreatic eruptions are at least 3 times more likely to be associated with high SO_2/CO_2 than not within a meaningful time window for monitoring purposes. The discordant phreatic events are of low energy, with $\text{SE} < 10$ (Fig. 3, Table S5), which do not pose a serious risk even to scientists working in the crater. The eruptions naturally fall into four groups based on SE (Fig. 3). Those with $\text{SE} < 10$ display high variability in terms




Fig. 3. Relationship between SE and SO_2/CO_2 . Diamonds are the average SO_2/CO_2 recorded within 24 h of eruption and circles indicate the closest SO_2/CO_2 value measured before each eruption. The individual eruptions (open symbols) naturally fall into four categories (closed symbols) based on seismic energy: $\text{SE} < 10$; $10 < \text{SE} < 30$; $30 < \text{SE} < 150$; $\text{SE} > 150$. Small eruptions on average are associated with lower SO_2/CO_2 and display high variance, whereas large eruptions show high associated SO_2/CO_2 , closest to fumarolic gas. This relationship is maintained considering only precursory gas measurements (green filled circles) and the generally lower SO_2/CO_2 values observed before eruptions reflect the fact that eruptions tend to occur before the peak in SO_2/CO_2 (Fig. 2). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

of associated SO_2/CO_2 , whereas all of those with $\text{SE} > 10$ have associated $\text{SO}_2/\text{CO}_2 > 1.9$ and their SO_2/CO_2 ratio displays a positive correlation with SE (diamonds in Fig. 3). If we consider the SO_2/CO_2 recorded prior to each eruption (Table S5; circles in Fig. 3) all but one eruption with $\text{SE} > 10$ show a precursory SO_2/CO_2 value greater than the median value (1.64) and the positive relationship between SO_2/CO_2 and SE is preserved.

We further explore the association between gas composition and the occurrence and energy of phreatic blasts by categorizing each lake gas data point based on diagnostic statistical tests to determine whether an eruption occurred within a given time window of the measurement (Fig. 4). In this analysis we follow the method of Tamburello et al. (2015), considering time windows of 24 h, 36 h, and 48 h, and the occurrence of eruptions with $\text{SE} > 1$ (i.e. all seismically registered eruptions), $\text{SE} > 10$ (8 eruptions), and $\text{SE} > 30$ (combining the 4 largest eruptions into one group; Fig. 3). Of the 40 measured gas compositions with $\text{SO}_2/\text{CO}_2 > 1.64$ (positive values; Fig. 4), 26 are associated with a phreatic blast with $\text{SE} > 10$ within 36 h (true positive values) and 14 are not (false positive values). The ratio of true positives to total number of positives gives a statistical measure of the association between high SO_2/CO_2

and eruption, which is 0.65 for eruptions of $\text{SE} > 10$ and a time window of 36 h. This ratio is known as “true predicted value”, and increases with the number of eruptions considered and larger time window (Fig. 4). Importantly, the ratio of true negatives (low SO_2/CO_2 and no eruption) to total number of negatives (Fig. 4; no eruption “predicted” based on low SO_2/CO_2) is high for the most dangerous eruptions, and approaches 1 for eruptions with $\text{SE} > 30$ and short time window (Table S5). The results show that both true predicted values (successful “prediction” of eruption based on high SO_2/CO_2) and false predicted values (successful “prediction” of quiescence based on low SO_2/CO_2) are high, indicating strong evidential worth for the association between gas composition and eruptions.

It is important to note that “predictive” in the sense of the above statistical analysis demonstrates trait associations between gas composition and the occurrence of eruptions, but does not prove that changes in SO_2/CO_2 occur before eruptions in a predictive sense useful for eruption forecasting. However, the transition from low to high SO_2/CO_2 prior to eruption occurs gradually over a period of days (Fig. 2), providing the possibility of developing a practically viable short-term precursory indicator of imminent eruption. We envisage a hypothetical tool for forecasting phreatic eruptions in which the slope of the SO_2/CO_2 trend (Fig. 2, Fig. 5) is evaluated in real-time considering lake Multi-GAS data collected during a defined time window (Fig. 5). The choice of time window depends firstly on a practical factor, which is data completeness and existence of data gaps due to low gas concentration (i.e. unfavorable meteorological conditions) or technical issues (usually low battery charge). The second consideration is timely response of the slope parameter to changes in gas composition. Choice of a shorter time window results in more variable slope values, more periods where the slope cannot be calculated due to too few data, but more rapid response of the parameter to changes in gas composition. Choice of a longer time window results in smoother but slower response, with fewer periods where the slope cannot be calculated (Fig. 5). For our dataset, 48 h and 90 h seem to be useful time windows to consider, with the shorter time window capturing changes in gas composition rapidly enough to show changes in slope sign prior to some eruptions that are not captured by the longer time window (e.g. 19 May eruption, Fig. 5). However, the shorter time window also produces a noisier response of the slope parameter that can cause false negative scenarios (e.g. gap and negative slope on 12–13 May during a series of eruptions; Fig. 5).

The slope analysis uses the data recorded prior to the last recorded data point and thus simulates the hypothetical situation of data arriving in real time, using the most recent data to assess the potential for eruption. The analysis using 48 h and 90 h




Fig. 4. Example of diagnostic test for categorizing lake gas SO_2/CO_2 data points and association with eruption (left). A true positive result is if an eruption of $\text{SE} > X$ (10 in this case) occurs within Y h (36 h here) of a measured gas composition with $\text{SO}_2/\text{CO}_2 > Z$ (1.64; the mean lake gas composition in this case). A false positive is if no eruption of $\text{SE} > X$ occurs within Y h of a gas measurement with $\text{SO}_2/\text{CO}_2 > Z$. A true negative result is if no eruption of $\text{SE} > X$ occurs within Y h of a gas measurement with $\text{SO}_2/\text{CO}_2 < Z$. A false negative result is if an eruption of $\text{SE} > X$ occurs within Y h of a gas measurement with $\text{SO}_2/\text{CO}_2 < Z$. This last category (eruptions associated with low SO_2/CO_2) is the least desirable scenario for hazards assessment. Eruptions with $\text{SE} < 10$ present little danger even to scientists working within the crater. The table on the right shows ratios of true positive results to total positive results and true negative results to total negative results (i.e. positive predicted values and negative predicted values; top and bottom, respectively) considering different SE thresholds (see Figs. 2 and 3) and time windows. Full results and errors are given in Table S6.




Fig. 5. We envisage a hypothetical scenario in which the slope of the SO_2/CO_2 trend (Fig. 2) is evaluated in real-time considering lake Multi-GAS data collected during a defined time window (e.g. past 48 h or 90 h) as a practically viable short-term eruption forecasting tool for phreatic eruptions. The size of the blue diamonds indicates eruptions by size as in Fig. 2. $\Delta(\text{SO}_2/\text{CO}_2)/\Delta t$ is calculated considering data collected in the past X h in a hypothetical scenario considering data arriving in real-time. Five of the eight eruptions with $\text{SE} > 10$ and three of the four eruptions with $\text{SE} > 30$ are predictable based on the 48 h slope. See Table S7 for summary of results. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Table 1

Excerpt of diagnostic test results on the statistical relationship between SO_2/CO_2 and the occurrence of phreatic eruptions within a given time window of the measurement (see Table S6 for full results. The number of cases is given for true positives, false positives, false negatives and true negatives for a SO_2/CO_2 gas ratio threshold of 1.64. The number of cases where the measured SO_2/CO_2 value falls within analytical error of the defined gas ratio threshold is given for each set of conditions. See text and Fig. 4 for details.

| | Eruption threshold | | | | | |
|---------------------------------|--------------------|-----------------|--------------------|-----------------|--------------------|-----------------|
| | SE > 30 (n = 4) | Within error | SE > 10 (n = 8) | Within error | SE > 1 (n = 14) | Within error |
| 12 h window | | | | | | |
| True positives | 8 | 4 | 13 | 4 | 20 | 6 |
| False positives | 32 | 8 | 27 | 8 | 20 | 6 |
| False negatives | 1 | 1 | 1 | 1 | 5 | 2 |
| True negatives | 39 | 9 | 39 | 9 | 35 | 8 |
| True positives/total positive s | 0.20 | | 0.33 | | 0.50 | |
| True negatives/total negative s | 0.98 | | 0.98 | | 0.88 | |
| 24 h window | | | | | | |
| True positives | 11 | 4 | 21 | 5 | 28 | 7 |
| False positives | 29 | 8 | 19 | 7 | 12 | 5 |
| False negatives | 2 | 1 | 2 | 1 | 12 | 4 |
| True negatives | 38 | 9 | 38 | 9 | 28 | 6 |
| True positives/total positives | 0.28 | | 0.53 | | 0.70 | |
| True negatives/total negatives | 0.95 | | 0.95 | | 0.70 | |
| 36 h window | | | | | | |
| True positives | 14 | 4 | 26 | 7 | 33 | 8 |
| False positives | 26 | 8 | 14 | 5 | 7 | 4 |
| False negatives | 3 | 1 | 9 | 3 | 15 | 5 |
| True negatives | 37 | 9 | 31 | 7 | 25 | 5 |
| True positives/total positives | 0.35 | | 0.65 | | 0.83 | |
| True negative/total negatives | 0.93 | | 0.78 | | 0.63 | |
| 48 h window | | | | | | |
| True positives | 18 | 5 | 31 | 9 | 38 | 11 |
| False positives | 22 | 7 | 9 | 3 | 2 | 1 |
| False negatives | 4 | 1 | 15 | 3 | 21 | 6 |
| True negatives | 36 | 9 | 25 | 7 | 19 | 4 |
| True positives/total positives | 0.45 | | 0.78 | | 0.95 | |
| True negatives/total negatives | 0.90 | | 0.63 | | 0.48 | |

time windows suggests that based on our actual data it would be possible to accurately forecast five out of the eight phreatic eruptions with $\text{SE} > 10$ (Table S7), demonstrating the high potential of real-time gas measurements to improve short-term forecasting of phreatic eruptions at volcanic lakes. The slope parameter is also successful at forecasting periods of quiescence. Negative slope values are associated with lack of eruptive events, and the ratio of true negative scenarios to total negatives (as assessed for Fig. 4


and Table 1) is greater than 0.8 for eruptions with $\text{SE} > 10$ for both 48 and 90 h time windows (Table S7).

The relationship between gas composition and eruptions also holds for the medium term at Poás, as no eruptions have occurred since October 2014 and campaign Multi-GAS measurements show the SO_2/CO_2 of lake gas emissions were consistently < 0.5 in 2015 and at the beginning of 2016. More importantly perhaps, our data show statistically significant relationships between gas composi-

tions and the occurrence of phreatic eruptions (Fig. 4) that yield a conceptual model for the short-term mechanisms that generate phreatic eruptions, which may be more widely applicable to monitoring wet volcanoes.

4. Discussion

The variations in SO₂/CO₂ in emissions from the Poás crater lake can be understood in terms of the relative susceptibility of SO₂ and CO₂ to reaction with hydrothermal fluids. The crater lake is the surface expression of an extensive hydrothermal system, contains a layer of liquid native sulfur at its base (Rowe et al., 1992; Aiuppa et al., 2014), is rich in dissolved sulfate, and precipitates a diverse array of sulfur bearing minerals from its waters (Biggs et al., 2014; Bennett and Raccichini, 1978). These secondary sulfur phases are derived from hydrolysis of magmatic gases injected into the sub-limnic hydrothermal system (Bennett and Raccichini, 1978). The dominant reaction in stripping SO₂ from magmatic gases in the crater lake is (Bennett and Raccichini, 1978; Oppenheimer and Stevenson, 1989):



as evidenced by sulfate-rich lake waters (Biggs et al., 2014) and abundant floating sulfur spherules (Movie S2). The degree of S isotope fractionation between HSO₄⁻ and elemental S ($\Delta^{34}\text{S}_{\text{HSO}_4-\text{S}_0} + 23.3\%$) in the lake strongly supports the dominance of equation (1) (Oppenheimer and Stevenson, 1989; Martínez et al., 2000). Sulfur isotope fractionation between native sulfur and bisulfate, and oxygen isotope fractionation between water and bisulfate in both indicate chemical and isotopic equilibration temperatures (>250 °C) far in excess of measured lake temperatures (~50 °C). High SO₂/CO₂ and high SO₂/H₂S in crater lake emissions also indicate gas equilibration temperatures (~300 °C) far higher than lake temperatures (Aiuppa et al., 2009). Therefore, the dynamics of the higher temperature sublimnic hydrothermal system are primarily responsible for the dominant features in the gas chemistry. Thermodynamics of reaction 1 dictate that SO₂ gas is less reactive at higher temperature and lower pH. Significantly, no H₂S was observed in crater lake gas emissions during the period of this study (above the Multi-GAS detection limit of ~1 ppm in SO₂-rich gas), whereas the H₂S/SO₂ measured in the fumaroles was 0.07–0.26. Importantly, H₂S injected into the sublimnic hydrothermal system is therefore completely stripped from gas entering the lake by partitioning into liquid native S, precipitation of sulfides, and reaction with SO₂ to form polythionates and elemental S (Oppenheimer and Stevenson, 1989). Disproportionation reactions producing H₂S from SO₂, commonly thought to be the “smoking gun” for gas interaction with hydrothermal fluids (Delmelle et al., 2000), are not significant in the Poás lake, calling for re-examination of gas-water interaction processes (Aiuppa et al., 2009). Whereas S is partially stripped from the magmatic gas input to the lake, CO₂ is effectively inert in acidic fluids and efficiently passes through the hydrothermal-lake system (Christenson et al., 2010).

We suggest that the short-term variations in SO₂/CO₂ in the lake gas emissions are due to varying influx of magmatic gas, which also determines whether phreatic eruptions occur or not. Comparing all DOAS lake SO₂ flux measurements (19 traverses conducted between March and November 2014 in which the lake and fumarolic plumes are distinguishable; Fig. 1, Table 2), the mean SO₂ flux associated with periods of phreatic activity is 81.4 ± 23.9 T/day ($n = 11$), whereas during non-eruptive episodes the lake SO₂ flux is 26.5 ± 8.9 T/day ($n = 8$). Increased gas flux reduces gas residence time and the interaction between gas species and lake water, kinetically inhibiting SO₂ hydrolysis (Werner et al., 2008; Aiuppa et al., 2009; Delmelle et al., 2000). Increased input of




Fig. 6. Box model for gas inputs and outputs for the Poás hydrothermal system (see Table 2). The SO₂ flux from the system (FSO_{2,out}; all fluxes in units of T/day) and SO₂/CO_{2,out} are constrained by DOAS and Multi-GAS measurements, respectively, allowing calculation of the CO₂ flux from the lake (FCO_{2,out}). The CO₂ flux from the lake is equal to the CO₂ flux into the lake (FCO_{2,in}) because CO₂ is inert in acidic fluids. The composition of gas feeding the lake is constrained by gas ratios measured at the high temperature fumaroles (X/CO_{2,fum}; where X = H₂O, SO₂, H₂S, and H₂), which allows calculation of the flux of gases into the crater lake (F_{X,in}) and the flux of S lost to the hydrothermal system (F_{s,hydrothermal}; expressed in T/d SO₂). As the flux and composition of magmatic gas into the lake are constrained, the heat transfer to the crater lake (dQ/dt) can be quantified (see supplementary online text for calculation details).

magmatic volatiles, high in SO₂, HCl, and HF (Pering et al., 2014; Kusakabe et al., 2000), also increases the acidity and temperature of sub-limnic (Oppenheimer and Stevenson, 1989) hydrothermal fluids, both of which shift equation (1) to the left and decrease the efficacy of the hydrothermal system in removing SO₂ from gas (Aiuppa et al., 2009). Fig. 6 presents a box model for the Poás lake system allowing calculation of the gas and heat flux into the base of the hydrothermal system (see supplementary online text and Table 2) and the mass of S lost from the gas input during eruptive and non-eruptive periods.

The DOAS results show that during eruptive episodes the SO₂ flux from the lake is ~3.1 times higher than during non-eruptive episodes, whereas the CO₂ flux is only a factor of ~1.8 higher. The mass of S lost from the gas to the hydrothermal system is similar during high and low gas flux, indicating that equation (1) has a reduced capacity for removing S from the magmatic gases during times of high volatile input. Thus, kinetics of S hydrolysis, the thermodynamics of the SO₂ disproportionation reaction occurring in the sub-limnic hydrothermal system, and the causative relationship between the injection of hot magmatic gas and phreatic explosions conspire to make SO₂/CO₂ in crater lake gas emissions a sensitive monitoring parameter for detecting short-term gas and heat flux variations leading to phreatic eruptions.

Magmatic gas is the principal source of volatiles (S, Cl, F) in volcanic lakes and also the main source of heat input (Rowe, 1994), which drives phreatic eruptions. As we can constrain the flux of gas to the lake (Fig. 6) for eruptive and non-eruptive episodes, we can calculate the heat flux for both cases. The heat flux to the lake is determined by integrating the temperature dependent heat capacity functions for H₂O, SO₂, CO₂, H₂, and H₂S from measured fumarolic temperature to the condensation temperature of water (100 °C), and summing the total enthalpy from gas cooling with the enthalpy of evaporation of water (Rowe, 1994) (see Supplementary Materials and Table 2). The average heat flux calculated for eruptive episodes is 58.9 MW compared to 27.2 MW for non-eruptive episodes (Fig. 5). Based on our seismic record of phreatic eruptions (Table S8) and the correlation between seismicity and column height (Fig. S7) as observed by webcam for a subset of eruptions (27 of the 60 seismically registered eruptions in 2014 were recorded by webcam; Movie S1), we can estimate the column height, as well as the energy released for all events following Symonds et al. (2001). The total eruptive power output for 2014 is estimated to be ~2.5 MW (Table S8). The difference in heat flux

Table 2
Flux calculations for Poás cráter lake for non-eruptive and eruptive periods.

| Date (dd-mm) | Non-eruptive fluxes | | | | Eruptive fluxes | | | | | | |
|----------------------------------|---------------------|--------|-------|-------|-----------------|--------------|--------|--------|--------------|-------|--------------|
| | 29-04 | 16-10 | 29-10 | 26-11 | 19-03 Av. | 16-05 | 28-08 | 30-09 | Eruptive Av. | | |
| Lake output | | | | | | | | | | | |
| SO ₂ flux | T/day | 18.1 | 29.6 | 10.4 | 47.8 | 26.5 | 51.6 | 76.6 | 94.1 | 103.4 | 81.4 |
| | ± | 3.4 | 10.4 | 7.4 | 14.4 | 8.9 | 11.6 | 38.3 | 18.8 | 26.7 | 23.9 |
| | n | 1 | 4 | 1 | 2 | 8 | 2 | 1 | 2 | 6 | 11 |
| SO ₂ /CO ₂ | molar | 0.61 | 0.95 | 1.08 | 0.89 | 0.88 | 0.88 | 2.07 | 1.64 | 1.31 | 1.47 |
| | ± | 0.04 | 0.16 | 0.06 | 0.21 | 0.12 | 0.05 | 0.10 | 0.50 | 0.20 | 0.21 |
| CO ₂ flux | T/day | 20.3 | 22.1 | 6.6 | 39.2 | 20.6 | 40.4 | 26.0 | 42.4 | 55.3 | 38.0 |
| | ± | 5.8 | 13.1 | 6.9 | 22.7 | 11.6 | 13.7 | 19.0 | 24.0 | 24.0 | 20.3 |
| Fumarole gas | | | | | | | | | | | |
| H ₂ O/CO ₂ | molar | 140.43 | 91.63 | 91.63 | 33.74 | 89.36 | 130.89 | 125.25 | 59.28 | 80.92 | 99.08 |
| SO ₂ /CO ₂ | molar | 2.33 | 2.65 | 2.65 | 1.46 | 2.28 | 2.17 | 2.44 | 2.88 | 2.00 | 2.37 |
| H ₂ /CO ₂ | molar | 0.36 | 0.32 | 0.32 | na | 0.33 | na | 0.38 | na | 0.45 | 0.41 |
| H ₂ S/CO ₂ | molar | 0.21 | 0.70 | 0.70 | 0.17 | 0.44 | 0.17 | 0.25 | 0.32 | 0.18 | 0.23 |
| H ₂ O | mol% | 97.29 | 95.15 | 95.15 | 92.77 | 95.1 | 97.51 | 96.85 | 93.39 | 95.71 | 95.9 |
| SO ₂ | mol% | 1.62 | 2.76 | 2.76 | 4.02 | 2.8 | 1.62 | 1.89 | 4.53 | 2.37 | 2.6 |
| CO ₂ | mol% | 0.69 | 1.04 | 1.04 | 2.75 | 1.4 | 0.75 | 0.77 | 1.58 | 1.18 | 1.1 |
| H ₂ | mol% | 0.25 | 0.33 | 0.33 | na | 0.3 | na | 0.29 | na | 0.53 | 0.4 |
| H ₂ | mol% | 0.15 | 0.73 | 0.73 | 0.46 | 0.5 | 0.12 | 0.19 | 0.50 | 0.21 | 0.3 |
| FLIR Temp | °C | 383 | 616 | 405 | 334 | 434.5 | 591 | 337 | 644 | 640 | 553.0 |
| Total Q | MJ/kg | 2.97 | 3.41 | 3.01 | 2.86 | 3.1 | 3.38 | 2.89 | 3.45 | 3.46 | 3.3 |
| Lake input | | | | | | | | | | | |
| H ₂ O flux | T/day | 1039 | 736 | 221 | 481 | 670 | 1923 | 1186 | 914 | 1627 | 1369 |
| SO ₂ flux | T/day | 69.0 | 85.3 | 25.6 | 83.5 | 68.3 | 127.8 | 92.4 | 177.5 | 160.9 | 131.1 |
| H ₂ flux | T/day | 0.7 | 0.6 | 0.2 | na | 0.6 | na | 0.9 | na | 2.2 | 1.4 |
| H ₂ flux | T/day | 3.5 | 12.6 | 3.8 | 5.4 | 7.5 | 5.5 | 5.4 | 11.0 | 8.1 | 7.1 |
| Total gas flux | kg/s | 13.1 | 9.9 | 3.0 | 7.0 | 8.9 | 24.3 | 15.2 | 13.2 | 21.4 | 17.9 |
| Heat flux | MW | 39.0 | 33.8 | 8.9 | 20.2 | 27.2 | 81.9 | 43.8 | 45.7 | 74.2 | 58.9 |

to the lake between eruptive and non-eruptive periods is about an order of magnitude higher than the eruptive power, indicating that more than enough energy is available through pulses of increased volatile input from the deeper magmatic system to generate eruptions (Fig. 5). It is worth noting that the amount of energy released during short-lived phreatic eruptions is very high, with instantaneous power of >200 GW for the largest eruptions, suggesting that energy is stored and catastrophically released below a low permeability seal (Sano et al., 2015; Kaneshima et al., 1996; Christenson et al., 2007; Fischer et al., 2015). Many of the eruptions occur slightly before the peak in SO₂/CO₂ (Fig. 2), perhaps suggesting that during increasing gas flux, an upward propagating front between vapor and liquid dominated hydrothermal regimes (Pering et al., 2014) encounters a confining zone (such as a sealed hydrothermal carapace Christenson et al., 2007, 2010; Giggenbach et al., 1990) saturated in lake water infiltrated during quiescent episodes. This water is vaporized and pressure builds until failure of the seal and phreatic eruption occur.

Poás is not an isolated case where changes in gas chemistry precede phreatic eruptions. Rincón de la Vieja is a remote volcano in northern Costa Rica that also hosts a hyper-acid lake and produced seismically registered phreatic eruptions in September and October of 2014. Multi-GAS surveys were conducted in March of 2013 (Shinohara et al., 2015), April of 2014, and October of 2014 and a six-fold increase in SO₂/CO₂ (Fig. 7) as well as a very large increase in SO₂/H₂S are observed in association with the transition from non-eruptive to eruptive behavior. Rare clear weather conditions on 14 April 2014 allowed us to measure SO₂ flux for the first time at Rincón de la Vieja (66 ± 17 T/d), which constrains the CO₂ flux to 312 ± 193 T/day. Although these most recent eruptions at Rincón de la Vieja were small, larger phreatic to phreato-magmatic events (VEI 2–3) at this volcano have produced eruptive columns up to ~4 km asl and generated destructive lahars.

Factors other than changes in gas flux and variations in the delivery of heat magmatic heat from depth must also play a role in the generation of phreatic eruptions and their size. The observa-

tion that the CO₂ flux at Rincón de la Vieja in April 2014 (before eruption) was an order of magnitude higher than at Poás during eruptive behavior (Fig. 5) suggests that differences in magmatic-hydrothermal plumbing systems dictate the heat flux threshold needed to generate phreatic eruptions, which probably plays a fundamental role in the potential for large phreatic explosions. Changes in the physical characteristics of the Poás lake are small (less than 10% from webcam images and field lake level measurements) compared to changes in gas fluxes (Fig. 6) and occur on longer timescales than the highly transient pulses of high temperature gas that generate individual phreatic blasts. Thus, changes in lake volume at Poás were probably not large enough to be the primary control on phreatic activity on short timescales. The composition of the gas emitted by the lake and the sulfur isotope systematics indicate that the sublimic hydrothermal systems are more influential on gas chemistry than superficial acid lakes. However, surficial physical changes may superimpose an additional control on the occurrence of phreatic eruptions. At the time of writing no phreatic eruptions had occurred at Poás since October of 2014, and since then the lake level has risen by ~4 m and H₂S has appeared in lake gas emissions. An outstanding question is whether a decrease in heat flux led to both an increase in lake level and the cessation of eruptions, or whether increased meteoric water input increased the hydraulic pressure in the sublimic hydrothermal system, which may have caused magmatic gases and heat to be channeled through the dome fumaroles, resulting in the cessation of phreatic eruptions. This outstanding issue is the subject of ongoing research.

5. Conclusions

Phreatic eruptions occurring through crater lakes represent the wet endmember of a spectrum of “non-magmatic” eruption types (Rouwet et al., 2014a). Here, we show that high frequency gas monitoring may provide a vital short- to medium-term precursor heralding phreatic blasts and that these eruptions are the direct




Fig. 7. Results of Multi-GAS campaigns conducted at Rincón de la Vieja volcano (Northern Costa Rica), which also hosts an ultra-acid crater lake. Data for April 2013 from Galle et al. (2002). Seismically registered phreatic eruptions occurred in September and October of 2014 and were associated with significant increase in SO₂/CO₂ and SO₂/H₂S.

result of fluctuations in magmatic gas flux from depth. Sano et al. (2015) recently showed that the 2014 Mt. Ontake eruption was preceded by long-term increase in ³He/⁴He, which was similarly interpreted to represent increased magmatic volatiles to the hydrothermal system. Together, these promising results call for enhanced gas monitoring at volcanoes prone to phreatic eruptions because magmatic heat, transferred by gases, is the driving force of these eruptions. We show that the transition from non-eruptive to eruptive state can occur due to relatively small changes in gas input (a factor of two; Fig. 6), which may not be detectable with standard geophysical methods. Traditional geochemical monitoring of volcanic lake water compositions has in many cases shown long term geochemical changes associated with increased magmatic volatile input, but monthly sampling campaigns and laboratory-based approaches will not provide key short-term phreatic precursors (Rouwet et al., 2014b). Furthermore, temporal changes in crater lake water chemistry are likely buffered by fluid residence time in the sublimic hydrothermal system (Oppenheimer and Stevenson, 1989). Gases, on the other hand, can pass rapidly through this reaction zone and dynamic changes in the sublimic system determine the extent to which reactive species are stripped from the magmatic gas. Short-term precursors are the indicators needed in order to meaningfully forecast eruptions and to evacuate proximal areas. Real-time, high frequency, *in situ* gas monitoring may provide the missing link between long term signals of volcanic reactivation (Sano et al., 2015) and sudden onset of phreatic eruption.

Explosive volcanic eruptions involving interaction with water are a diverse, poorly characterized, and particularly hazardous category of eruptions that remain the most challenging to forecast by traditional volcano monitoring techniques. Here we show that individual phreatic eruptions at Poás are preceded by changes in lake gas composition due to increasing input of magmatic volatiles. This finding suggests an intricate tie between shallow-sourced eruptions and short-period fluctuations in magma degassing intensity. These processes and associated precursory signals likely operate at many volcanoes where magmatic gases sustain hydrothermal systems prone to phreatic eruptions. Our results call for high frequency real-time monitoring of gas emissions at volcanoes to advance understanding of short-term changes in magma degassing, shallow energy storage, and associated volcanic hazards.

Acknowledgements

This project was supported through funding from the Deep Carbon Observatory Deep Earth Carbon Degassing program, as well as from the European Research Council (FP7/ERC grant agreement no. 305377), the Comisión Nacional de Prevención de Riesgos y Atención de Emergencias (CNE Costa Rica), and the United States Agency for International Development (USAID). Guillermo Alvarado and Cyril Müller are thanked for insightful discussions. Cindy Werner, Matthew Kirk, and Simon Robson are thanked for external reviews of earlier versions of this manuscript.

Author contributions: JMdm, AA, and TPF conceived of the study. JMdm, JP, GA, ML, MM, and GG collected the data. ML and GG built the Multi-GAS instruments and provided technical expertise with instrument installation, maintenance, and data processing. JMdm, JP, CK, GA, and MM processed the data. JMdm, AA, JP, CK, TPF and GA wrote the paper, with important contributions and comments from ML and MM.

Appendix A. Supplementary material

Supplementary material related to this article can be found online at <http://dx.doi.org/10.1016/j.epsl.2016.02.056>.

Uncited references

(Brown et al., 1989) (Pasternack and Varekamp, 1997)

References

- Aiuppa, A., et al., 2007. Forecasting Etna eruptions by real-time observation of volcanic gas composition. *Geology* 35, 1115–1118.
- Aiuppa, A., et al., 2009. The 2007 eruption of Stromboli volcano: insights from the real-time measurement of the volcanic gas plume CO₂/SO₂ ratio. *J. Volcanol. Geotherm. Res.* 182, 221–230.
- Aiuppa, A., et al., 2013. First observations of the fumarolic gas output from a restless caldera: implications for the current period of unrest (2005–2013) at Campi Flegrei. *Geochem. Geophys. Geosyst.* 14, 4153–4169.
- Aiuppa, A., et al., 2014. Gas measurements from the Costa Rica–Nicaragua volcanic segment suggest possible along-arc variations in volcanic gas chemistry. *Earth Planet. Sci. Lett.* 407, 134–147.
- Barberi, F., Bertagnini, A., Landi, P., Principe, C., 1992. A review on phreatic eruptions and their precursors. *J. Volcanol. Geotherm. Res.* 52, 131–246.
- Bennett, F., Raccichini, S., 1978. Subaqueous sulphur lake in Volcan Poas. *Nature* 271, 342–344.
- Biggs, J., et al., 2014. Global link between deformation and volcanic eruption quantified by satellite imagery. *Nature* 5. <http://dx.doi.org/10.1038/ncomms4471>.

- 1 Brown, G., et al., 1989. Energy budget analysis for Poás crater lake: implications for
2 predicting volcanic activity. *Nature* 339, 370–373.
- 3 Christenson, B.W., et al., 2007. Hazards from hydrothermally sealed volcanic con-
4 ducts. *Eos* 88, 53–55.
- 5 Christenson, B.W., et al., 2010. Cyclic processes and factors leading to phreatic erup-
6 tion events: insights from the 25 September eruption through Ruapehu Crater
7 Lake, New Zealand. *J. Volcanol. Geotherm. Res.* 191, 15–32.
- 8 Delmelle, P., Bernard, A., Kusakabe, M., Fischer, T.P., Takano, B., 2000. Geochemistry
9 of the magmatic-hydrothermal system of Kawah Ijen volcano, East Java, Indone-
10 sia. *J. Volcanol. Geotherm. Res.* 97, 31–53.
- 11 Fischer, T.P., et al., 2015. Temporal variations in fumarole gas chemistry at Poás vol-
12 cano, Costa Rica. *J. Volcanol. Geotherm. Res.* 294, 56–70.
- 13 Galle, B., et al., 2002. A miniaturized ultraviolet spectrometer for remote sensing of
14 SO₂ fluxes: a new tool for volcano surveillance. *J. Volcanol. Geotherm. Res.* 119,
15 241.
- 16 Giggenbach, W.F., Garcia, N., Londoño, A., Rodriguez, L., Rojas, N., Calvache, M.L.,
17 1990. The chemistry of fumarolic vapor and thermal-spring discharges from the
18 Nevado del Ruiz volcanic-magmatic-hydrothermal system, Colombia. *J. Volcanol.*
19 *Geotherm. Res.* 42, 13–39.
- 20 Global Volcanism Program, 2013. *Volcanoes of the World*, v.4.4.1. Ven-
21 zke, E. (Ed). Smithsonian Institution. Downloaded 15 Jan 2015.
22 <http://dx.doi.org/10.5479/si.GVP.VOTW4-2013>.
- 23 Jolly, A.D., Sherburn, S., Jousset, P., Kilgour, G., 2010. Eruption source processes from
24 seismic and acoustic observations of the 25 September 2007 Ruapehu – North
25 Island, New Zealand. *J. Volcanol. Geotherm. Res.* 191, 35–43.
- 26 Kaneshima, S., et al., 1996. Mechanism of phreatic eruption at Aso volcano inferred
27 from near-field broadband seismic observations. *Science* 273, 642–645.
- 28 Kusakabe, M., Komoda, Y., Takano, B., Abiko, T., 2000. Sulfur isotopic effects in the
29 disproportionation reaction of sulfur dioxide in hydrothermal fluids: implica-
30 tions for the $\delta^{34}\text{S}$ variations of dissolved bisulfate and elemental sulfur from
31 active crater lakes. *J. Volcanol. Geotherm. Res.* 97, 287–307.
- 32 Martínez, M., et al., 2000. Chemical evolution and volcanic activity of the ac-
33 tive crater lake of Poás volcano, Costa Rica, 1993–1997. *J. Volcanol. Geotherm.*
34 *Res.* 97, 127.
- 35 Mastin, L.G., Witter, J.B., 2000. The hazards of eruptions through lakes and seawater.
36 *J. Volcanol. Geotherm. Res.* 97, 195–214.
- 37 Oppenheimer, C., Stevenson, D., 1989. Liquid sulphur lakes at Poás volcano. *Nature*
38 342, 790–793.
- 39 Pasternack, G.B., Varekamp, J.C., 1997. Volcanic lake systematic I. Physical con-
40 straints. *Bull. Volcanol.* 58, 528–538.
- 41 Pering, et al., 2014. High time resolution fluctuations in volcanic carbon dioxide
42 degassing from Mount Etna. *J. Volcanol. Geotherm. Res.* 270, 115–121.
- 43 Rouwet, D., et al., 2014a. Recognizing and tracking volcanic hazards related to non-
44 magmatic unrest: a review. *J. Appl. Volcanol.* 3, 17.
- 45 Rouwet, D., Tassi, F., Mora-Amador, R.A., Sandri, L., Chiarini, V., 2014b. Past,
46 present and future of volcanic lake monitoring. *J. Volcanol. Geotherm. Res.* 272,
47 78–97.
- 48 Rowe, G.L., 1994. Oxygen, hydrogen, and sulfur isotope systematic of the crater lake
49 system of Poás Volcano, Costa Rica. *Geochem. J.* 28, 263–287.
- 50 Rowe, G.L., et al., 1992. Fluid-volcano interaction in an active stratovolcano: the
51 crater lake system of Poás volcano, Costa Rica. *J. Geophys. Res.* 49, 23–51.
- 52 Sano, Y., et al., 2015. Ten-year helium anomaly prior to the 2014 Mt Ontake erup-
53 tion. *Sci. Rep.* 5, 13069. <http://dx.doi.org/10.1038/srep13069>.
- 54 Shinohara, H., Yoshikawa, S., Miyabuchi, Y., 2015. In: Rouwet, D., Christenson,
55 B.W., Tassi, F., Vandemeulebrouck, J. (Eds.), *Volcanic Lakes*. Springer, Heidelberg,
56 pp. 201–217.
- 57 Symonds, R.B., Gerlach, T.M., Reed, M.H., 2001. Magmatic gas scrubbing: implica-
58 tions for volcano monitoring. *J. Volcanol. Geotherm. Res.* 108, 303–341.
- 59 Tamburello, G., et al., 2015. Intense magmatic degassing through the lake of Co-
60 pahue volcano, 2013–2014. *J. Geophys. Res., Solid Earth* 120. <http://dx.doi.org/10.1002/2015JB012160>.
- 61 Werner, C., et al., 2008. Variability of passive gas emissions, seismicity, and de-
62 formation during crater lake growth at White Island Volcano, New Zealand,
63 2002–2006. *J. Geophys. Res.* 113, B01204.
- 64 Williams, S.N., Stoiber, R.E., Garcia, N.P., Adela Londoño, C., Gemmell, Bruce, Lowe,
65 Donald R., Connor, C.B., 1986. Eruption of the Nevado del Ruiz Volcano, Colom-
66 bia, on 13 November 1985: gas flux and fluid geochemistry. *Science* 233,
964–967.

XMLVIEW: extended**Appendix A. Supplementary material**

The following is the Supplementary material related to this article.

Label: MMC 1

caption: The Supplementary Material file contains text describing the data processing methods used for analysis of the Multi-GAS, DOAS, and seismic data, details of the uncertainty assessments for Multi-GAS measurements, and calculations used in estimates of gas and energy budgets for the Poás crater lake. Data are presented in Tables S1–S8 and Figs. S1–S7.

link: **APPLICATION : mmc1**

Label: MMC 2

caption: Movie showing examples of phreatic eruptions at Poás volcano, Costa Rica (see separate file).

link: **VIDEO : mmc2**

Label: MMC 3

caption: Movie showing the behavior of gas plumes from the fumaroles and lake at Poás volcano, Costa Rica (see separate file).

link: **VIDEO : mmc3**

Sponsor names

Do not correct this page. Please mark corrections to sponsor names and grant numbers in the main text.

5 **Q8** **European Research Council**, country=Belgium, grants=3 05377

6 **USAID**, country=United States, grants=

UNCORRECTED PROOF

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66

67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132

Highlights

- Changes in volcanic lake gas emissions precede phreatic eruptions.
- First identified short-term precursor to phreatic eruptions.
- Sulfur scrubbing can be negligible in hyper-acid volcanic lake environment.
- Gas flux measurements allow quantification of hydrothermal system energy budget.
- Transient pulses of magmatic gas cause phreatic eruptions.

UNCORRECTED PROOF

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
6667
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132