



Annual Review of CyberTherapy and Telemedicine

Virtual Reality in Healthcare:
Medical Simulation and Experiential Interface

Editors:

Brenda K. Wiederhold, Ph.D., MBA, BCB, BCN

Giuseppe Riva, Ph.D., M.S., M.A.

Mark D. Wiederhold, M.D., Ph.D., CPE, FACP, FACPE



Annual Review of Cybertherapy And Telemedicine 2015

Virtual Reality in Health Care:
Medical Simulation and Experiential Interface

Edited by

Brenda K. Wiederhold

*Interactive Media Institute, San Diego, CA, USA
Virtual Reality Medical Institute, Brussels, Belgium*

Giuseppe Riva

*Catholic University of Milano, Milan, Italy
Istituto Auxologico Italiano, Milan, Italy*

and

Mark D. Wiederhold

Virtual Reality Medical Center, San Diego, CA, USA

Annual Review of CyberTherapy and Telemedicine, Volume 13

Annual Review of CyberTherapy and Telemedicine

Copyright © 2015 Interactive Media Institute
9565 Waples Street, Suite 200
San Diego, CA 92121, USA

ISBN: 1554-8716

All rights reserved.
Printed in the United States of America

Journal Web site: <http://www.arctt.info>
Interactive media Institute Website: <http://www.interactivemediainstitute.com>

LEGAL NOTICE

The publisher is not responsible for the use which might be made of the following information

General Information

Annual Review of CyberTherapy and Telemedicine (ARCTT – ISSN: 1554-8716) is published annually (once per year) by the Interactive Media Institute (IMI), a 501c3 non profit organization, dedicated to incorporating interdisciplinary researchers from around the world to create, test, and develop clinical protocols for the medical and psychological community. IMI realizes that the mind and body work in concert to affect quality of life in individuals and works to develop technology that can be effectively used to improve the standards and reduce the cost of healthcare delivery.

Interactive Media Institute
9565 Waples Street, Suite 200
San Diego, CA, 92121 USA
Telephone: (858) 642-0267
Fax: (858) 642-0285
E-mail: cybertherapy@vrphobia.com IMI
Web site: <http://www.interactivemediainstitute.com>
Journal Web site: <http://www.arctt.info>

Copyright © 2015 by Interactive Media Institute. Printed in the United States of America.

About the Journal

ARCTT is a peer-reviewed all-purpose journal covering a wide variety of topics of interest to the mental health, neuroscience, and rehabilitation communities. The mission of ARCTT is to provide systematic, periodic examinations of scholarly advances in the field of CyberTherapy and Telemedicine through original investigations in the Telemedicine and CyberTherapy areas, novel experimental clinical studies, and critical authoritative reviews. It is directed to healthcare providers and researchers who are interested in the applications of advanced media for improving the delivery and efficacy of mental healthcare and rehabilitative services.

Manuscript Proposal and Submission

Because Annual Review papers examine either novel therapeutic methods and trials or a specific clinical application in depth, they are written by experienced researchers upon invitation from our Editorial Board. The editors nevertheless welcome suggestions from our readers. Questions or comments about editorial content or policies should be directed to the editors only.

Manuscript Preparation

Manuscripts should be submitted in electronic format on CD-Rom or floppy disks as well as on 8 1/2 x 11-in. paper (three copies), double-spaced format. Authors should prepare manuscripts according to the Publication Manual of the American Psychological Association (5th Ed.). Original, camera-ready artwork for figures is required. Original color figures can be printed in color at the editors' discretion and provided the author agrees to pay in full the associated production costs; an estimate of these costs is available from the ARCTT production office on request. ARCTT policy prohibits an author from submitting the same manuscript for concurrent consideration by two or more publications. Authors have an obligation to consult journal editors concerning prior publication of any data upon which their article depends. As this journal is a primary journal that publishes original material only, ARCTT policy prohibits as well publication of any manuscript that has already been published in whole or substantial part elsewhere, unless authorized by the journal editors.

Disclaimer

All the published papers, editorial news and comments, opinions, findings, conclusions, or recommendations in ARCTT are those of the author(s), and do not necessarily reflect or constitute the opinions of the Journal, its Publisher, and its editorial staff.

Decision Making and Cognitive Behavioral Flexibility in a OCD Sample: a Study in a Virtual Environment

Filippo LA PAGLIA^{a,1}, Caterina LA CASCIA^a, Rosalinda RIZZO^a, Giuseppe RIVA^b,
and Daniele LA BARBERA^a

^a *Department of Experimental Biomedicine and Clinical Neuroscience, University of Palermo, Italy*

^b *Applied Technology for Neuro-Psychology Lab, Istituto Auxologico Italiano, Milan, Italy*

Abstract. Neuropsychological disorders are common in Obsessive-Compulsive Disorder (OCD) patients. Executive functions, verbal fluency and verbal memory, shifting attention from one aspect of stimuli to others, mental flexibility, engaging in executive planning and decision making, are the most involved cognitive domains. We focus on two aspects of neuropsychological function: decision making and cognitive behavioral flexibility, assessed through a virtual version of the Multiple Errand Test (V-MET), developed using the NeuroVR software. Thirty OCD patients were compared with thirty matched control subjects. The results showed the presence of difficulties in OCD patients with tasks where the goal is not clear, the information is incomplete or the parameters are ill-defined.

Keywords. Obsessive-Compulsive Disorder, decision making, cognitive behavioral flexibility, Virtual Multiple Errands Test, neuropsychological assessment

Introduction

Obsessive-Compulsive Disorder (OCD) is a psychiatric disorder involving distressing intrusive thoughts and related compulsive behaviours [1]. Neuropsychological studies have also shown that patients with OCD show deficits in cognitive abilities that are strictly linked to the functioning of the frontal lobe and its related fronto-subcortical structures, such as executive functioning deficits and insufficient cognitive-behavioral flexibility [2]. Individuals with OCD have been observed to experience difficulties in shifting attention from one aspect of stimuli to others, engaging in executive planning [3]. A relevant cognitive skill linked to the executive functions is decision-making. It is a cognitive mechanism that can be related to different executive processes and involves the ability to evaluate environmental information of various options that ensures actions are taken in the light of positives and negatives of each option. The effectiveness of decision-making requires some degree of predicting outcome certainty

¹ Corresponding Author. Filippo La Paglia, Dipartimento di Biomedicina Sperimentale e Neuroscienze Cliniche, Università di Palermo, Italy. E-mail: filippolapaglia@gmail.com; filippo.lapaglia@unipa.it

of each option in relation to particular situations. Decision making appears to be dysfunctional in the OCD, in the context of doubting and uncertainty as well as in dual and sequencing tasks.

The role of uncertainty in decision-making has yet to be fully investigated systematically in OCD [4]. However, clinical decision-making performance reveals that OCD participants request more information about, and spend more time deliberating over low-risk scenarios and OCD-relevant decisions compared to non-anxious controls [5]. OCD patients seem to require more information before making a decision and excessive worry and doubt, core features of OCD, could mediate difficulties in decisions-making [5].

The present study is aimed to explore and assess the decision-making and the mental and behavioral flexibility in OCD patients through complex tasks of virtual version (VR) of the Multiple Errands Test (MET) [6, 7, 8] developed based on the NeuroVR software. The virtual environment is a supermarket, performed in a shopping setting and involves the completion of various tasks of different complexity levels (e.g. buy a sponge or two products from the refrigerated products aisle as going to the beverage aisle and asking about). To complete the task, participants must follow several rules: *“you are not allowed to enter any aisle unless it is a part of a task, you are not allowed to go into the same aisle more than once, you are not allowed to buy more than two items for item category”*. The time is stopped when the participant says: *“I finished”*. After the tasks and the rules have been explained, patients are able to plan and choose the sequence of actions needed to complete the tasks. In this way, many different executive functions are stimulated: ability to plan a sequence of actions, problem solving and cognitive and behavioral flexibility.

Based on this, the virtual version of the Multiple Errands Test (VMET) has been recently developed and tested in different clinical populations [9, 10, 11].

Methods

Participants

We evaluated 30 OCD patients (M=15, F=15; mean age=33,07 years, std.dev.= 9,906) and 30 healthy controls (M=16, F=14; mean age=34,00 years, std.dev.=10,841) with no history of psychiatric disorder. Patients, selected from the Outpatient Unit of Psychiatry of Palermo University Hospital, were excluded from the study if they had a severe cognitive impairment (MMSE<19), a severe motor impairment which did not allow subjects to perform the procedure, auditory language comprehension difficulties (Token Test<26,5), object recognition impairments (Street Completion Test<2,25), excessive state and trait anxiety (STAI>40) and excessive depression state (Beck Depression Inventory>16).

Instrument and procedure

Decision making and cognitive behavioral flexibility were assessed by a 1) validated classic neuropsychological battery and by 2) Virtual Multiple Errands Test (V-MET), with uncertain goal, lacking information and undefined parameters.

The neuropsychological battery including: Trail Making Test (TMT, Forms A, B and B-A) to assess selective attention and task switching; Frontal Assessment Battery (FAB) to assess executive functions including ability to initiate and stop actions, to monitor and change behavior as needed, and to plan future behavior when faced with novel tasks and situations. Executive functions allow to anticipate outcomes and adapt to changing situations. Furthermore, to evaluate multi-tasking and cognitive flexibility, Phonological verbal fluency and Semantic verbal fluency were employed. Second assessment stimulates many different executive functions such as planning a sequence of actions, problem solving and cognitive and behavioral flexibility.

Results and conclusions

One-way analyses of variance (AnOVAs) were used to compare patients and controls performance.

Results showed a statistical significant difference between two groups. Both evaluation revealed a significantly poorer performance of OCD patients.

Regarding the neuropsychological evaluation, OCD patients show deficits in the attention function TMT A ($F= 10,806$, $p= .002$) and in task switching TMT A-B ($F= 6,004$, $p= .018$); difficulties in executive function, FAB ($F= 13,857$, $p= .001$) and in mental flexibility, Phonemic Fluencies ($F= 11,212$, $p= .002$), Semantic Fluencies ($F= 18,288$, $p= .000$).

In order to investigate differences in VMET, analyzed variables were the execution time for the entire task; the errors in executing the tasks; partial tasks failures, inefficiencies. Specific items of task failures regarding divided attention, self-correction, and no evidence of perseveration. After virtual reality assessment the clinical sample show a performance significantly lower (table 1).

First of all, in regard the time used by participant to complete the task, analysis showed significant differences between groups and indicated that the healthy control group took significantly less time compared with OCD patients ($F=4,069$, $p= 0,040$). Indeed the execution time for the whole task was higher in patients with OCD compared to controls, suggesting that patients with OCD need more time in planning than controls.

Concerning the task failure, results showed significant differences between groups, too. Patients showed errors in tasks requiring a change in the primary goal and the ability to respond simultaneously to multiple task demands (sub-tests 6, buying two products from the refrigerated products aisle), (sub-tests 7, going to the beverage aisle and asking about what to buy). Regarding inefficiencies (less capacity of planning) and errors of perseveration (perseverating in errors is a clear sign of reduced flexibility) data revealed significant differences between groups. Patients showed poorer ability in using effective strategies that facilitate the carrying out of the tasks; for example, accurate planning before starting a specific subtask or using the map to move into the virtual supermarket. These executive deficits may reflect a specific deficit in cognitive flexibility; namely, ability to change selectively to effectively respond to external/internal stimulation. More, scores for divided attention are lower in controls (lower is the score, better is the performance).

On the basis of these results OCD patients showed difficulties in mental flexibility and in the task requiring divided attention. Consequently, patients are not able to recognize their own errors and autocorrect (reduced cognitive flexibility typical in OCD patients), modifying their behavior and their strategies based on the goal to reach. In conclusion,

Table 1. Virtual reality assessment

	OCD patients	Healthy control	
	<i>Mean ± SD</i>	<i>Mean ± SD</i>	<i>ANOVAs</i>
Time	10,77 ± 5,345	8,230 ± 4,284	F= 4,069, p= 0,040*
Sub test 6 <i>Buy two products from the refrigerated products</i>	9,66 ± 1,951	8,27 ± ,691	F= 13,457, p= 0,001†
Sub test 7 <i>Go to the beverage aisle and ask what to buy</i>	10,52 ± 2,181	8,77 ± 1,431	F= 13,373, p= 0,001†
Inefficiencies range: 8 (great ineff.) to 32 (no ineff.)	22,55 ± 4,733	25,57 ± 4,415	F= 6,407, p= 0,014*
Divided attention range: 7 (no errors) to 14 (great errors)	10,448 ± 2,667	8,333 ± 1,647	F= 13,531, p= 0,001†
Self-correction range: 7 (no errors) to 14 (great errors)	9,241 ± 1,902	7,833 ± ,833	F= 13,718, p= 0,000†
No perseveration range: 7 (no errors) to 14 (great errors)	8,724 ± 1,830	7,733 ± ,944	F= 6,896, p= 0,011*

†. Correlation is significant at the 0.01 level (2-tailed).*. Correlation is significant at the 0.05 level (2-tailed)

our results showed that VMET appears sensitive to evaluate the functional status of OCD with normal cognition, as manifested in terms of executive deficits. In particular, VMET allows the possibility to evaluate some subcomponents of executive functions in ecological settings, giving an accurate analysis of patients' deficits as well as traditional tests. Our data, confirming previous study, provide support for the feasibility of using the Virtual Multiple Errands Test (VMET) as an assessment tool of executive functions.

References

- [1] American Psychiatric Association, Diagnostic and Statistical Manual of Mental Disorders. (Fourth ed.) American Psychiatric Pub, Washington, DC, 1994.
- [2] M. Abbruzzese, S. Ferri, S. Scarone. The selective breakdown of frontal functions in patients with obsessive- compulsive disorder and in patients with schizophrenia: a double dissociation experimental finding. *Neuropsychologia* 35 (1997), 907–912.
- [3] M. M. Fenger, A. Gade, KH. Adams, ES. Hansen, TG. Bolwig, GM. Knudsen. Cognitive deficits in obsessive- compulsive disorder on tests of frontal lobe functions. *Nordic Journal of Psychiatry* 59 (2005). 39-44.
- [4] P. Cavadini, A. Gorini, L. Bellodi. Understanding obsessive compulsive disorder: focus on decision making. *Neuropsychol Rev* 16 (1) (2006), 3–15.

- [5] E. B. Foa, A. Mathews, J.S. Abramowitz, N. Amir, A. Przeworski, J. C. Filip, A. Alley. Do patients with obsessive-compulsive disorder have deficits in decision-making? *Cognitive Therapy and Research* 27(4) (2003), 431–445.
- [6] S. Raspelli, L. Carelli, F. Morganti, B. Poletti, B. Corra, V. Silani, G. Riva, Implementation of the Multiple Errands Test in a NeuroVR-supermarket: a Possible Approach. *Studies in Health Technology and Informatics* 154 (2010), 115–119.
- [7] F. La Paglia, C. La Cascia, R. Rizzo, G. Riva, D. La Barbera, Assessment of Executive Functions in Patients with Obsessive Compulsive Disorder by Neuro VR. *Studies in Health Technology and Informatics* 181 (2012), 98–102.
- [8] F. La Paglia, C. La Cascia, R. Rizzo, F. Cangialosi, M. Sanna, G. Riva, D. La Barbera, Cognitive Assessment of OCD Patients: NeuroVR vs Neuropsychological test. *Studies in Health Technology and Informatics* 199 (2014), 40–44.
- [9] P. Cipresso, F. La Paglia, C. La Cascia, G. Riva, G. Albani, D. La Barbera, Break in volition: A virtual reality study in patients with obsessive-compulsive disorder. *Experimental Brain Research* 229 (2013), 443–449.
- [10] F. La Paglia, C. La Cascia, R. Rizzo, L. Sideli, A. Francomano, D. La Barbera, Cognitive Rehabilitation of Schizophrenia Through NeuroVR Training. *Studies in Health Technology and Informatics* 191 (2013), 158–162.
- [11] F. La Paglia C. La Cascia, P. Cipresso, R. Rizzo, A. Francomano, G. Riva, D. La Barbera. Psychometric assessment using classic neuropsychological and virtual reality based test: A study in Obsessive-Compulsive Disorder (OCD) and Schizophrenic patients. *Lecture Notes of the Institute for Computer Sciences, Social Informatics and Telecommunications Engineering* 100 (2014), 23–32.