21st Meeting of the Group of European Charophytologists 18-22 September 2017, València, Spain



PROGRAM & ABSTRACTS



Organizing committee

Dr. Maria A. Rodrigo (Chair)

Dr. CARMEN ROJO

http://www.uv.es/~rodrigoa/

http://www.uv.es/~rojo/

Associate Professors of Ecology (Department of Microbiology and Ecology, University of València) Senior Researchers

SARA CALERO ERIC PUCHE

MATI SEGURA

PhD Students

Freelance biologist Integrative Ecology lab collaborator

Integrative Ecology Laboratory
Ecology, Ethology and Evolution Group (e³)
Cavanilles Institute for Biodiversity and Evolutionary Biology
University of Valencia
www.uv.es/cavanilles/



Cavanilles Institute for Biodiversity and Evolutionary Biology



ICBIBE UV

Sponsored by:





The organizers acknowledge Dr. María José Carmona, Dr. Ximo Baixeras, Dr. Javier Armengol and Dr. Francesc Mesquita for the loan of binocular microscopes for the determination session, and the help of the Institute Cavanilles staff (Alejandra Valderrama, Pilar Ruíz) in the organization tasks. We also thank Dr. Allan Chivas for revising the English.

Printed in Valencia, July 2017



Oral presentation 18

CHAROPHYTES FOR DESCRIPTION AND MONITORING OF INLAND WATERS IN SICILY

Angelo TROIA and Rossella BARONE

Università degli Studi di Palermo, Dip. STEBICEF, sez. Botanica ed Ecologia Vegetale, via Archirafi 38, 90123 Palermo, Italy. Presenter e-mail: angelo.troia@unipa.it

The island of Sicily, in the centre of the Mediterranean basin, is a well-known biodiversity hotspot (Médail and Quézel, 1997). Unfortunately, the knowledge of its charophyte flora is absolutely unsatisfactory. The few available data derive from some scattered phytosociological works and some articles published more than 100 years ago. According to the last national synthesis (Bazzichelli and Abdelahad, 2009) a total of 19 Charophyte species should occur in the island, distributed in 4 genera: Chara L. (11), Nitella Agardh (5), Tolypella (A. Braun) A. Braun (2) and Lamprothamnium J. Groves (1). However an updated list of the species occurring in the island, together with their distributions, is currently lacking. It is noteworthy that, owing to this lack of local knowledge about charophytes, information currently available on these habitats occurring in the sites of the Natura 2000 network in Sicily is absolutely insufficient. We think that the group deserves a better knowledge, both for taxonomic, biogeographic and "pure" science, and for "applied" aspects connected with water quality and biodiversity conservation, also according to important European Directives (2000/60/CE and 92/43/EEC, respectively). So our work started, gathering data and making preliminary surveys. During the preliminary investigations, charophytes have been found in both freshwater and brackish waters, from 0 to 1400 m a.s.l. In particular, Chara vulgaris L. and Lamprothamnium papulosum (Wallr.) J. Groves have been identified in different coastal seasonally flooded habitats, referred to the priority habitat "1150* coastal lagoons" according to the mentioned 92/43 Directive. Chara contraria A. Braun ex Kütz. has been identified in springs and streams, referred to the habitat "3140 Hard oligo-mesotrophic waters with benthic vegetation of Chara spp.". Other species have been found in other habitat types, such as the priority habitat "3170* Mediterranean temporary ponds".

Médail and Quézel (1997). Annals of the Missouri Botanical Garden 84: 112–127.

Bazzichelli and Abdelahad (2009). Alghe d'acqua dolce d'Italia. Flora analitica delle Caroficee.

Sapienza Università di Roma - Ministero dell'Ambiente e della Tutela del Territorio e del

Mare.