

Microbial methane oxidation leading to extreme isotopic fractionation in thermal springs of central Greece

D'Alessandro W.*¹, Daskalopoulou K.²⁻³, Gagliano A.L.¹, Calabrese S.², Fiebig J.⁴, Tassi F.⁵, Kyriakopoulos K.³
& Li Vigni L.²

¹ Istituto Nazionale di Geofisica e Vulcanologia, sezione di Palermo

² Dipartimento di Scienze della Terra e del Mare, Università di Palermo

³ National and Kapodistrian University, Athens, Greece

⁴ Institut für Geowissenschaften, Goethe-Universität, Frankfurt-am-Main, Germany

⁵ Dipartimento di Scienze della Terra, Università di Firenze

* Corresponding email: walter.dalessandro@ingv.it

Keywords: geothermal gases, methanotrophy, stable isotopes.

The Greek territory belongs to the geodynamically active Alpine-Himalayan orogenic belt. As such, it shows intense seismic activity, active volcanic systems and areas of enhanced geothermal fluxes. One of these areas is the Sperchios basin and the northern part of Evia island in central Greece, which present widespread thermal manifestations (D'Alessandro et al., 2014). Five of them with temperatures from 33 to 80°C present bubbling gases whose dominating species are either CO₂ or N₂. All gases contain from 27 to 4000 ppm of CH₄. The isotopic composition of CH₄ in these gases covers a wide range with δ¹³C values ranging from -21.7 to +16.9 ‰ and δ²H values ranging from -124 to +301 ‰.

The hottest manifestation displays the lowest isotopic values within the typical range of volcanic and geothermal systems. All the remaining samples fit a methane oxidation trend reaching very positive values. If we consider the lowest values as the deep hydrothermal marker the obtained ΔH/ΔC values range between 5 and 13 which are close to those typical of microbially driven oxidation (Kinnaman et al., 2007).

Although the outlet temperature of the hottest manifestations is at the upper limit for methanotrophic microorganisms (Sharp et al., 2014), we can hypothesize that environmental conditions are not favorable for their survival at this site while they can thrive in the other, strongly consuming methane and producing very positive isotopic values.

D'Alessandro, W., Brusca, L., Kyriakopoulos, K., Bellomo, S., Calabrese S. (2014): A geochemical traverse along the "Sperchios Basin-Evoikos Gulf" Graben (Central Greece): origin and evolution of the emitted fluids. *Mar. Petrol. Geol.*, 55, 295-308.

Kinnaman, F.S., Valentine, D.L., Tyler, P.A. (2007): Carbon and hydrogen isotope fractionation associated with the aerobic microbial oxidation of methane, ethane, propane and butane. *Geochim. Cosmochim. Acta*, 71, 271-283.

Sharp, C.E., Smirnova, A.V., Graham, J.M., Stott, M.B., Khadka, R., Moore, T.R., Grasby, S.E., Strack, M., Dunfield, P.F. (2014): Distribution and diversity of Verrucomicrobia methanotrophs in geothermal and acidic environments. *Environ. Microbiol.*, 16, 1867-1878.