

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/329176585>

A new contribution to the alien macroalgal flora of the Ustica Island Marine Protected Area (Tyrrhenian Sea, Italy)

Article in *BioInvasions Records* · January 2018

DOI: 10.3391/bir.2018.7.4.03

CITATIONS

0

READS

6

4 authors, including:



Anna Maria Mannino
Università degli Studi di Palermo

41 PUBLICATIONS 287 CITATIONS

[SEE PROFILE](#)



Maria Cristina Gambi
Stazione Zoologica Anton Dohrn di Napoli

312 PUBLICATIONS 6,444 CITATIONS

[SEE PROFILE](#)



Paola Gianguzza
Università degli Studi di Palermo

79 PUBLICATIONS 892 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Project Effects of elevated CO₂ from volcanic vents on the trophic structure of seagrass meadows [View project](#)



Project Physiological adaptation and plasticity to acidification in marine invertebrates living around shallow water CO₂ vent systems [View project](#)

Rapid Communication

A new contribution to the alien macroalgal flora of the Ustica Island Marine Protected Area (Tyrrhenian Sea, Italy)

Anna Maria Mannino¹, Maria Cristina Gambi², Tiziana Dieli³ and Paola Gianguzza^{4,*}

¹Department of Biological, Chemical and Pharmaceutical Sciences and Technologies, Section of Botany and Plant Ecology, University of Palermo, Via Archirafi 38, 90123 Palermo, Italy

²Stazione Zoologica Anton Dohrn, Dept. of Integrative Marine Ecology, Villa Dohrn-Benthic Ecology Center, 80070 Ischia (Napoli), Italy

³AQUAPRO S.a.s., Via A. Di Giovanni 14, 90144 Palermo, Italy

⁴Department of Earth and Marine Science, University of Palermo, ConISMa, Via Archirafi 18, 90123 Palermo, Italy

Author e-mails: annamaria.mannino@unipa.it (AMM), gambimc@gmail.com (MCG), tdiel@hotmail.com (TD), paola.gianguzza@unipa.it (PG)

*Corresponding author

Received: 16 January 2018 / Accepted: 18 July 2018 / Published online: 1 October 2018

Handling editor: Carla Lambertini

Abstract

A second record of the non-indigenous species *Botryocladia madagascariensis* G. Feldmann (Rhodophyta) along the coasts of the “Ustica Island” Marine Protected Area (Tyrrhenian Sea, Italy) is reported. With this additional record for the Sicilian coasts, we provide a description of collected specimens of *B. madagascariensis* and habitat details. Numerous fertile female gametophytes with cystocarps immersed in vesicles were also observed, for the second time in the Mediterranean Sea, and described.

Key words: non-indigenous species, *Botryocladia madagascariensis*, Mediterranean Sea

Introduction

The spread of non-indigenous species (NIS) is an ongoing phenomenon which is widely recognized as a major threat to biodiversity and natural ecosystem function (Wallentinus and Nyberg 2007). In the Mediterranean Sea, the annual number of recorded NIS has increased exponentially over the last 100 years (Occhipinti-Ambrogi et al. 2011a, b; Katsanevakis et al. 2013). Multiple human-related stressors (i.e. habitat fragmentation and destruction, pollution, maritime transport, tourism, aquaculture, exploitation of resources, global warming, and opening of the Suez Canal) have led to the introduction of ~ 1000 NIS, of which 134 are macrophytes (24 Chlorophyta, 79 Rhodophyta, 30 Ochrophyta, and 1 Tracheophyta) (Verlaque et al. 2015; Alós et al. 2016; Galil et al. 2017).

Identifying future NIS and taking effective steps to prevent their dispersal and establishment constitutes an enormous challenge to marine biologists. Sicily

and its surrounding islands, located at the crossroads between the Western and Eastern Mediterranean, are particularly vulnerable to the introduction of NIS. In this region, the exposure to multiple transport pathways, including fisheries and recreational fleets, has promoted the introduction either accidentally or intentionally of several NIS, despite the high number of Marine Protected Areas (MPAs) already established. This area plays an important role as a receiver, transit and donor zone for NIS within the Mediterranean Sea, and monitoring and surveillance is required to reduce the risk of future NIS introductions. The creation of permanent habitat “alarm systems” might be an effective tool in the management of NIS introductions. In this regard, Sicilian MPAs can play an important role as “sentinel sites” where the effects of NIS invasion can be studied and management strategies can be developed to counter and contrast such negative effects. In order to develop effective managing plans for the prevention and control of NIS,




Figure 1. Map showing the localities in the Mediterranean Sea where *Botryocladia madagascariensis* has been recorded. The records are chronologically arranged in the following order: 1 and 2 (Lampedusa Island and Castelluccio (Syracuse), respectively; Cormaci et al. 1992), 3 (Malta; Cormaci et al. 1997), 4 (Comino; Cormaci et al. 1997), 5 (Riserva Ciclopi; Pizzuto 1999), 6 (Gargano promontory; Cecere et al. 2000), 7 (Tremiti Islands; Cormaci et al. 2000), 8 (Seki Beach, Gulf of Antalya; Turna et al. 2000), 9 (Side region, Gulf of Antalya; Turna et al. 2000), 10 (Gulf of Taranto; Cecere and Petrocelli 2004), 11 (Porto Cesareo; Cecere et al. 2005), 12 (Ustica Island; Catra et al. 2005), 13 (Linosa Island; Serio et al. 2006), 14 (Favignana Island; Catra et al. 2006), 15 (Karpathos Island; Catra and Giardina 2009), 16 (Porto Germeno, Korinthiakos Gulf; Tsiamis and Verlaque 2011), 17 (Baleari Islands; Joher et al. 2012), 18 (Messenian Gulf; Nicolaïdou et al. 2012), 19 (Lakonic Gulf; Nicolaïdou et al. 2012), 20 (Cyprus; Taşkin et al. 2013), 21 (Limni, Evia Island; Tsiamis and Panayotidis 2015), 22 (Ustica Island; present study).

Table 1. Comparison of taxonomic characters of *Botryocladia botryoidea* and *B. madagascariensis* specimens described in Wilkes et al. (2006) and examined in the present study.

Species	Vesicle morphology			Secretory cell		
	Shape	Cortex	Cell layers	Shape	No. secretory cells/group	Position within vesicle
<i>Botryocladia botryoidea</i> (Wulfen) Feldmann 1941	pyriform	continuous	2-3	~ spherical	1/2	borne directly on inner medullary cells
<i>Botryocladia madagascariensis</i> G. Feldmann 1944	ovoid	continuous	6?	ovoid/subspherical	2-4 (rarely 1)	borne on support cells and directly on medullary cells
<i>Botryocladia madagascariensis</i> (present study)	obovate/ovoid	continuous	3-4	ovoid/spherical	1 (rarely 2)	borne on support cells and directly on medullary cells

data on the distribution and spread dynamics of current NIS are essential.

Botryocladia madagascariensis G. Feldmann, a Rhodophyta macroalgal, is widely distributed along the Indian Ocean coast of South Africa (e.g., along the coasts of Madagascar, Natal and the Canary Islands) and has already been reported within the Mediterranean Sea (for details on localities and chronology see Figure 1). *B. madagascariensis* may be easily confused with the native species *Botryocladia botryoidea* (Wulfen) Feldmann, but they differ in morphology at

the cellular level (Table 1). This species was first recorded in the Mediterranean Sea at Lampedusa Island and Castelluccio (Syracuse) (Sicily; Cormaci et al. 1992), and later at Malta (170 and 210 kilometres, respectively, from the first recorded site). Afterwards, it was reported from southern Italy, several Sicilian localities, Turkey, Greece, Balearic Islands, and Cyprus (Figure 1). Both shipping or Suez Canal have been speculated as pathways of introduction for *B. madagascariensis* into the Mediterranean Sea (Cormaci et al. 2004; Verlaque et al. 2015). The goal

of this study was to provide a new record of *B. madagascariensis* along the coasts of the Ustica Island MPA, a description of recorded specimens, and habitat details.

Material and methods

Study area

The study was carried out at Ustica Island, which is the protruding part of an extinct volcano located in the southern Tyrrhenian Sea (Western Mediterranean; 38.7°N; 13.18333°E), 60 km north of the Sicilian coast. Its base lies at about 2000 m depth and is composed mainly of alkaline basalts and sedimentary rocks. The Marine Protected Area (MPA), created in 1986, encompasses a total area of 16.000 ha and contains three zones with different degrees of protection. The no-take zone (zone A) covers 65 ha along the western part of the Island, while the general reserve (zone B) and the take zone (zone C) share the remaining area equally (Figure 2). According to the constitutive decree of the MPA (D.I. 12/11/1986), only research activities are authorized in the no-take zone, whereas recreational activities, such as SCUBA diving, boat anchoring, swimming and angling, are strictly prohibited. There are no restrictions on recreational activities in either the B or C zone (Gianguzza et al. 2006).

During surveys carried out on August 17 and October 12, 2017 at Colombara Bank off Ustica Island (38.7256°N; 13.1883°E), specimens of *B. madagascariensis* were observed and collected. Colombara Bank is a large basaltic rocky bank located approx. 1 km offshore the northern coast of Ustica Island in the B zone of the MPA. The summit of the bank, with a surface of approx. 500 square meters, is shallow (3–8 m depth) and almost horizontally orientated. The bank summit is almost entirely covered by a dense and rich photophilous macroalgal assemblage, mainly composed of brown algae including habitat-formers *Cystoseira* spp. (Giaccone et al. 1985). The perimeter of the summit abruptly breaks off in steep cliffs with a columnar basaltic structure, which descends almost vertically down to 40–45 m depth.

Sampling and processing

Specimens of *B. madagascariensis* were sampled by SCUBA diving and hand collection since the thalli were easily detachable from the substrate. Since *B. madagascariensis* is similar in habit and morphology to the




Figure 2. Map showing the zonation of the Ustica Island MPA. The site of Colombara Bank is indicated by the black arrow.

native *B. botryoides*, visual identification in the field can be challenging. In the laboratory, fresh, whole specimens were photographed, manually sectioned with a razor blade, and observed under a light microscope for the analyses of diagnostic characters. Photos were taken with a Stereo Microscope (Stemi SV 11 Apo Zeiss) and photomicrographs with an Inverted Microscope (Axio Observer Zeiss). The main distinctive character between the native and invasive species is the disposition of the gland cells, which in *B. botryoides* are usually solitary and only occur on the inner layer of medullary cells (Table 1).

Results

Habitat description

In August some specimens ($N = 30$) were found at 7 m depth on the summit of Colombara Bank, a few meters from the margin of the break off the southern oriented cliff. In October all the observed thalli ($N = 35$) were collected at 7.5 m depth on the top of the vertical cliff of the same relief, exposed to the south-west/west. The collected thalli (Figure 1) clumped together and were attached directly to the rocky substrate via holdfasts. They settled in spots among or below other algal species, such as *Halopteris scoparia* (Linnaeus) Sauvageau and *Cystoseira* spp., but were not overgrown by them and visible by a naked eye. In October, the specimens were smaller and settled relatively close to specimens of the Mediterranean endemic scleractinia *Astroites calicularis* (Pallas, 1766).

Specimen description

All of the analyzed specimens ($N = 35$), 2.5–3.5 cm high, presented erect and cartilaginous red thalli that

were pseudodichotomously branched with numerous elliptical/obovate hollow and mucilage-filled vesicles (up to 5 mm long and 4 mm wide) all around the axis (Figure 3a). Vesicles showed a continuous layer of small cortical cells (4–5 µm in diameter) and 3–4 layers of medullary cells gradually increasing in size inward (up to 80 µm long and 60 µm wide) (Figure 3b, c). Secretory cells, single or rarely in pairs, were produced both on innermost medullary cells and special bearing cells that protruded into the vesicular cavity (Figure 3d, e). Female gametophytes with immersed cystocarps scattered irregularly over the vesicles (4–5 per vesicle) were present (Figure 3a, f). Cystocarps reached up to 950 µm in diameter and 550 µm in height (including the pericarp). The mature gonia was sub-spherical and composed of carposporangia (14 to 23 µm in diameter).

Discussion

This finding represents the second record of *Botryocladia madagascariensis* from the “Ustica Island” MPA (the first one reported in Catra et al. 2005). Since the species has been recorded at least twice and spread over time and space (see Zenetos et al. 2005), we can consider it as established at Ustica Island MPA. While Catra et al. (2005), reported the species in a list with no description or images, this study provides the first description of specimens collected in this MPA with detailed observations on location and habitat. Our finding is also the first record of female gametophytes with cystocarps from Sicily, and the second one for the Mediterranean Sea (Cormaci et al. 1997). In general, reproductive structures were rarely reported in the Mediterranean Sea (but see sporophytes by Cormaci et al. 1992 and female gametophytes by Cormaci et al. 1997). Reproductive structures have only been observed on *B. madagascariensis* in the Western basin of the Mediterranean Sea (i.e., Lampedusa and/or Castelluccio, Ustica Island and Maltese Islands). It is possible that *B. madagascariensis*, a warm-temperate species, spreads mainly vegetatively in the Mediterranean, and sexual reproduction occurs only in areas where environmental conditions are suitable (Norris 1989).

Examining chronological records (Figure 1), the alga was reported first from the eastern coast (Lampedusa Island and Castelluccio, Syracuse), then from the western (Egadi Islands) and northern (Ustica Island MPA) coasts (see Figure 1), confirming the warm-temperate affinity hypothesis (Norris 1989). Giaccone et al. (1985) highlighted, through an analysis of the marine flora of Ustica, that this Island is subtropical. The subtropical character has also been highlighted by all the other NIS species

occurring in the Island, namely the macroalgae *Acrothamnion preissii* (Sonder) E. M. Wollaston, *Asparagopsis armata* Harvey, *Caulerpa cylindracea* Sonder and *Womersleyella setacea* (Hollenberg) R.E. Norris (Catra et al. 2007; Di Trapani et al. 2016; Gianguzza P. personal observation), the heterobranch mollusc *Aplysia dactylomela* (Rang, 1828) (Gambi et al. 2017) and the decapod crustacean *Percnon gibbesi* (H. Milne Edwards, 1853) (Noè et al. 2018).

Few previously published reports of *B. madagascariensis* in the Mediterranean provide descriptions or images of the species. Since confusion with *B. botryoides* is possible without the aid of microscopes, it is natural to wonder how many of these records are really records of *B. madagascariensis* and how many are the result of misidentification. Therefore, we currently do not know the exact distribution of *B. madagascariensis* in the Mediterranean Sea. In addition, *B. madagascariensis* could not be a recently introduced species in the Mediterranean Sea (Ribera and Boudouresque 1995). Seagrief (1984) highlighted a misidentification between *B. madagascariensis* and *B. botryoides* in South Africa in the 1980s; therefore, it is possible that the distribution areas of the two species also overlap in the Mediterranean Sea (Turna et al. 2000). This study confirms that each new NIS record must be accompanied by either illustrations or descriptions of diagnostic traits that are difficult to check in the field, and which are often necessary for taxonomic certainty. Moreover, when morphological diagnostic characters of a NIS are not easy to check, as for the genus *Botryocladia* within which the species delineation remains complex (see Afonso-Carrillo and Sobrino 2003; Wilkes et al. 2006), additional molecular techniques should be used. Therefore, further molecular research is needed to verify all of the records of *B. madagascariensis* reported for the Mediterranean Sea.

The presence of NIS in “Ustica Island” MPA, but also in other MPAs (Cecere et al. 2005; Mannino and Balistreri 2017; Mannino et al. 2014, 2017), confirms that MPAs, despite their recognized role as a management tool for the protection of local biodiversity, are not immune from NIS invasions. However, NIS have been largely disregarded in marine conservation plans (Giakoumi et al. 2016). For this reason, it is crucial to establish a long-term monitoring programme to limit the introduction of NIS and control the spread of existing ones. The creation of permanent observatories involving all Sicilian MPAs will enable scientists and managers to detect new introductions and follow the spread of species already present to help manage NIS in the MPAs and in neighboring areas.




Figure 3. *Botryocladia madagascariensis* collected from the “Ustica Island” Marine Protected Area (Tyrrhenian Sea, Italy). Female gametophytes showing cystocarps (arrow) on vesicles (a), surface view of cortical cells of vesicles (b), transverse section of vesicle wall showing cortical and medullary cell layers (c), a secretory cell (arrow) on a special bearing cell (d), inner surface of the vesicle wall showing a secretory cell (arrow) on a special bearing cell (e), transverse section of a cystocarp showing the pericarp of smaller cells and carposporangia (f). Photographs by A.M.

Acknowledgements

The authors are grateful to Mauro Maniscalco (Alta Marea Diving Center Ustica) and Bruno Zava for their essential help during field collection and transport of samples. We also wish to thank Enric Ballesteros and Donatella Serio, for initial advice in species identification, Marc Verlaque for providing bibliographic assistance and two anonymous reviewers for their helpful comments and criticism.

References

- Afonso-Carrillo J, Sobrino C (2003) Vegetative and reproductive morphology of *Botryocladia botryoides*, *B. occidentalis* and *B. canariensis* sp. nov. (Rhodymeniaceae, Rhodophyta) from the Canary Islands. *Phycologia* 42: 138–150, <https://doi.org/10.2216/i0031-8884-42-2-138.1>
- Alós J, Tomas F, Terrados J, Verbruggen H, Ballesteros E (2016) Fast-spreading green beds of recently introduced *Halimeda incrassata* invade Mallorca Island (NW Mediterranean Sea). *Marine Ecology Progress Series* 558: 153–158, <https://doi.org/10.3354/meps11869>
- Catra M, Giardina S (2009) A survey of the marine macroalgae of Karpathos Island (the Aegean Sea, Greece). *Plant Biosystems* 143: 509–515, <https://doi.org/10.1080/11263500903172185>
- Catra M, Drago D, Giaccone T, Giardina S, Nicastro A (2005) Biodiversità vegetale marina dell'Isola di Ustica (Palermo). *Bullettino Accademia Gioenia Scienze Naturali Catania* 38(365): 193–211
- Catra M, Alongi G, Serio D, Cormaci M, Furnari G (2006) The benthic algal flora on rocky substrata of the Egadi Islands, a marine protected archipelago off the western coast of Sicily (Italy, Mediterranean Sea). *Nova Hedwigia* 82: 489–538, <https://doi.org/10.1127/0029-5035/2006/0082-0489>
- Catra M, Giardina S, Giaccone T, Basso D, Giaccone G (2007) Il Coralligeno del Secchitello di Ustica (Palermo). Coralligenous assemblages of Secchitello (Ustica - Palermo). *Biologia Marina Mediterranea* 14(2): 174–175
- Cecere E, Petrocelli A (2004) Floristic and biogeographic considerations about the benthic macroalgal flora in the Gulf of Taranto. *Biogeographia* 25: 7–18, <https://doi.org/10.21426/B6110108>
- Cecere E, Petrocelli A, Alongi G, Saracino OD, Cormaci M, Furnari G (2000) Marine benthic flora of the Gargano promontory (Adriatic Sea, southern Italy). *Flora Mediterranea* 10: 325–347
- Cecere E, Petrocelli A, Saracino OD (2005) Biodiversity of phytobenthic communities in the marine reserve of Porto Cesareo. *Biologia Marina Mediterranea* 12(1): 78–87
- Cormaci M, Furnari G, Alongi G, Dinaro R, Pizzuto F (1992) On the occurrence in Sicily of three Florideophyceae new to the Mediterranean Sea. *Botanica Marina* 35: 447–450, <https://doi.org/10.1515/botm.1992.35.5.447>
- Cormaci M, Lanfranco E, Borg JA, Buttigieg S, Furnari G, Micallef SA, Mifsud C, Pizzuto F, Scammacca B, Serio D (1997) Contribution to the knowledge of benthic marine algae on rocky substrata of the Maltese Islands (Mediterranean Sea). *Botanica Marina* 40: 203–215, <https://doi.org/10.1515/botm.1997.40.1-6.203>
- Cormaci M, Furnari G, Alongi G, Catra M, Serio D (2000) The benthic algal flora on rocky substrata of the Tremiti Islands (Adriatic Sea). *Plant Biosystems* 134: 133–152, <https://doi.org/10.1080/11263500012331358404>
- Cormaci M, Furnari G, Giaccone G, Serio D (2004) Alien macrophytes in the Mediterranean Sea: a review. In: Pandalai SG (ed), Recent Research Developments in Environment Biology, Vol. 1. Part I, Chapter 9. Research Signpost, India, pp 153–202
- Di Trapani F, Noè S, Bonaviri C, Gianguzza P (2016) First record of the alien alga *Caulerpa cylindracea* Sonders in the Mediterranean marine protected area of Ustica Island (Tyrrhenian Sea, Italy). In: Proceedings of the International Congress GeoSub. Ustica, Italy, September 13–17, 2016, p 39
- Galil B, Marchini A, Occhipinti-Ambrogi A, Ojaveer H (2017) The enlargement of the Suez Canal - Erythraean introductions and management challenges. *Management of Biological Invasions* 8: 141–152, <https://doi.org/10.3391/mbi.2017.8.2.02>
- Gambi MC, Carugati F, Crocetta F, Gianguzza P (2017) Aplysie in ...piscina! Una popolazione di *Aplysia dactylomela* (Heterobranchia, Aplysiidae) nell'Isola di Ustica. *Nottuario S.I.B.M.* 72: 99–102
- Giaccone G, Alessi MC, Toccaceli M (1985) Flora e vegetazione marina dell'isola di Ustica. *Bollettino Accademia Gioenia Scienze Naturali Catania* 18(326): 505–536
- Giakoumi S, Guilhaumon F, Kark S, Terlizzi A, Claudet J, Felline S, Cerrano C, Coll M, Danovaro R, Fraschetti S, Koutsoubas D, Ledoux JB, Mazor T, Mérigot B, Micheli F, Katsanevakis S (2016) Space invaders; biological invasions in marine conservation planning. *Diversity and Distribution* 22: 1220–1231, <https://doi.org/10.1111/ddi.12491>
- Gianguzza P, Chiantore M, Bonaviri C, Cattaneo-Vietti R, Vielmini I, Riggio S (2006) The effects of recreational *Paracentrotus lividus* fishing on distribution patterns of sea urchins at Ustica Island MPA (Western Mediterranean, Italy). *Fisheries Research* 81: 37–44, <https://doi.org/10.1016/j.fishres.2006.06.002>
- Joher S, Ballasteros E, Cebrán E, Sánchez N, Rodríguez-Prieto C (2012) Deep-water macroalgal-dominated coastal detritic assemblages on the continental shelf off Mallorca and Menorca (Balearic Islands, Western Mediterranean). *Botanica Marina* 55: 485–497, <https://doi.org/10.1515/bot-2012-0113>
- Katsanevakis S, Zenetos A, Belchior C, Cardoso AC (2013) Invading European Seas: assessing pathway of introduction of marine aliens. *Ocean Coastal Management* 76: 64–74, <https://doi.org/10.1016/j.ocecoaman.2013.02.024>
- Mannino AM, Balistreri P (2017) An updated overview of invasive *Caulerpa* taxa in Sicily and circum-Sicilian Islands, strategic zones within the NW Mediterranean Sea. *Flora Mediterranea* 27: 221–240, <https://doi.org/10.7320/FIMedit27.221>
- Mannino AM, Balistreri P, Yókés MB (2014) First record of *Aplysia dactylomela* (Opisthobranchia, Aplysiidae) from the Egadi Islands (western Sicily). *Marine Biodiversity Record* 7: e22, <https://doi.org/10.1017/S1755267214000190>
- Mannino AM, Paraspoto M, Crocetta F, Balistreri P (2017) An updated overview of the marine alien and cryptogenic species from the Egadi Islands Marine Protected Area (Italy). *Marine Biodiversity* 47: 469–480, <https://doi.org/10.1007/s12526-016-0496-z>
- Nicolaidou A, Alongi G, Aydogan O, Catra M, Cavas L, Cevik C, Dosi A, Circosta V, Giakoumi S, Giménez-Casalduero F, Filiz H, Izquierdo-Muñoz A, Kalogirou S, Konstantinidis E, Kousteni V, Kout J, Legaki A, Megalofonou P, Ovalis P, Paolillo G, Paschos I, Perdikaris C, Poursanidis D, Ramos-Esplá AA, Reizopoulou S, Sperone E, Taskin E, Tripepi S, Vázquez Luis M (2012) New Mediterranean Biodiversity Records (June 2012). *Mediterranean Marine Science* 13: 162–174, <https://doi.org/10.12681/mms.33>
- Norris RE (1989) Natalian Botryocladia (Rhodymeniales, Rhodophyceae) including description of a new, long axis-forming species. *Botanica Marina* 32: 131–148, <https://doi.org/10.1515/botm.1989.32.2.131>
- Occhipinti-Ambrogi A, Marchini A, Cantone G, Castelli A, Chimenz-Gusso C, Cormaci M, Froglio C, Furnari G, Gambi MC, Giaccone G, Giangrande A, Gravili C, Mastrototaro F, Mazzotti C, Orsi-Rolini L, Piraino S (2011a) Alien species along the Italian coasts: an overview. *Biological Invasions* 13: 215–237, <https://doi.org/10.1007/s10530-010-9803-y>
- Occhipinti-Ambrogi A, Marchini A, Cantone G, Castelli A, Chimenz-Gusso C, Cormaci M, Froglio C, Furnari G, Gambi MC, Giaccone G, Giangrande A, Gravili C, Mastrototaro F, Mazzotti C, Orsi-Rolini L, Piraino S (2011b) Erratum to: alien

- species along the Italian coasts: an overview. *Biological Invasions* 13: 531–532, <https://doi.org/10.1007/s10530-010-9856-y>
- Pizzuto F (1999) On the structure, typology and periodism of a *Cystoseira brachycarpa* J. Agardh emend. Giaccone community and of a *Cystoseira crinita* Duby community from the eastern coast of Sicily (Mediterranean Sea). *Plant Biosystems* 133: 15–35, <https://doi.org/10.1080/11263509909381529>
- Ribera MA, Boudouresque CF (1995) Introduced marine plants, with special reference to macroalgae: mechanisms and impact. In: Round FE, Chapman DJ (eds), *Progress in Phycological Research*, Vol. 11. Biopress Ltd., Bristol, UK, pp 187–268
- Seagrief SC (1984) A Catalogue of South African Green, Brown and Red Algae. *Memoires Botanical Surveys, South Africa* 47: 1–72
- Serio D, Alongi G, Catra M, Cormaci M, Furnari G (2006) Changes in the benthic algal flora of Linosa Island (Straits of Sicily, Mediterranean Sea). *Botanica Marina* 49: 135–144, <https://doi.org/10.1515/BOT.2006.018>
- Taşkın E, ÖzTÜRK M, Kurt O, Ulçay S (2013) Benthic marine algae in Northern Cyprus (Eastern Mediterranean Sea). *Journal of Black Sea/Mediterranean Environment* 19(2): 143–161
- Tsiamis K, Verlaque M (2011) A new contribution to the alien red macroalgal flora of Greece (Eastern Mediterranean) with emphasis on *Hypnea* species. *Cryptogamie Algologie* 32: 393–410, <https://doi.org/10.7872/crya.v32.iss4.2011.393>
- Tsiamis K, Panayotidis P (2015) New record and further expansion of alien macroalgae in Greece. In: Proceedings of the 11th Pan-Hellenic Symposium on Oceanography and Fisheries. Mytilene, Lesvos, Greece, May 13–17, pp 601–604
- Turna I, Cormaci M, Furnari G (2000) First record of *Botryocladia madagascariensis* G. Feldmann (Rhodymeniaceae, Rhodophyceae) from the Gulf of Antalya, Mediterranean coast of Turkey. *Plant Biosystems* 15: 111–115, <https://doi.org/10.1080/11263500012331350415>
- Verlaque M, Ruitton S, Mineur F, Boudouresque CF (2015) CIESM Atlas of exotic species in the Mediterranean, Vol. 4. Macrophytes. CIESM Publishers, Monaco, 362 pp
- Wallentinus I, Nyberg CD (2007) Introduced marine organisms as habitat modifiers. *Marine Pollution Bulletin* 55: 323–332, <https://doi.org/10.1016/j.marpolbul.2006.11.010>
- Wilkes RJ, McIvor L, Guiry MD (2006) Vegetative morphology and rbcL phylogeny of some members of the genera *Botryocladia* and *Irvinea* (Rhodymeniaceae, Rhodophyta). *Phycologia* 45: 481–494, <https://doi.org/10.2216/05-11.1>
- Zenetas A, Cinar M, Pancucci-Papadopoulou M, Harmelin J, Furnari G, Andaloro F, Bellou N, Streftaris N, Zibrowius H (2005) Annotated list of marine alien species in the Mediterranean with records of the worst invasive species. *Mediterranean Marine Science* 6: 63–118, <https://doi.org/10.12681/mms.186>