

# Antibiotic Resistance Profiling, Analysis of Virulence Aspects and Molecular Genotyping of *Staphylococcus aureus* Isolated in Sicily, Italy

Maria Vitale,<sup>1</sup> Salvatore Gaglio,<sup>1,\*</sup> Paola Galluzzo,<sup>1</sup> Giuseppe Cascone,<sup>1</sup> Chiara Piraino,<sup>1</sup> Vincenzo Di Marco Lo Presti,<sup>1</sup> and Rosa Alduina<sup>2</sup>

## Abstract

*Staphylococcus aureus* is the major cause of foodborne diseases worldwide. In this retrospective study, 84 *S. aureus* strains were characterized. The collection comprises 78 strains isolated during 1998 and 2014 from dairy products and tissue samples from livestock bred for dairy production in Sicily. One isolate was obtained from a pet (dog), one from an exotic animal (a circus elephant), and four human isolates were obtained during a severe food poisoning outbreak that occurred in Sicily in 2015. All the strains were characterized by pulsed-field gel electrophoresis (PFGE), for antibiotic resistance and presence of toxin genes. PFGE results showed 10 different pulsotypes, with three relatively frequent and three unique. The antibiotic resistance profiling showed that penicillin G (35.7%) and tetracycline (20.2%) resistance is largely spread. Most isolates contained at least one toxin gene making them a potential threat for public health. Enterotoxin *sec* gene was observed in 28.6% and *seg* in 23.8% of the strains, respectively; the human isolates were the only ones to concurrently harbor both *seg* and *sei* genes. In addition, 24 isolates were randomly selected and analyzed by multilocus sequence typing. Interestingly, the analysis showed the presence of 12 sequence types (STs), of which 6 were novel. One of them, ST700, was detected in 29% of the isolates and was found to be spread throughout Sicily. ST700 has been present in the island for almost 16 years (1998–2014) and it shows no host preference since it was isolated from different ruminant species. Four human isolates shared both the pulsotype (PT10) and the sequence type (ST9), as well as the virulence genes (*seg-sei*); this observation suggests that the isolates originated from a single clone, although they were obtained from two different individuals.

**Keywords:** antibiotic resistance, MLST, PFGE, *Staphylococcus aureus*, MRSA, toxin genes

## Introduction

*STAPHYLOCOCCUS AUREUS* IS a major resident or transient colonizer of the skin and the mucosa of humans and primates. *S. aureus* can cause a variety of infections, from superficial skin infections to severe, and potentially fatal, invasive diseases (Wang *et al.*, 2014; Aires-de-Sousa, 2017; Sergelidis and Angelidis, 2017). *S. aureus* is also a common pathogen of ruminants such as cattle, goats, and sheep that may lead to clinical and subclinical mastitis. The pathogen

can spread from the udder of the infected animal into raw milk and dairy products, affecting the quality and quantity of the products; therefore, the pathogen can become a significant economic burden for farmers and a serious problem for the dairy industry (Seegers *et al.*, 2003).

In the last years, different multidrug-resistant strains have emerged making *S. aureus* a major concern for public health. The multidrug-resistant phenotype is a particular characteristic of the methicillin-resistant *S. aureus* (MRSA) strains (Gould *et al.*, 2012; Rodvold and McConeghy, 2014). The

<sup>1</sup>Istituto Zooprofilattico Sperimentale della Sicilia, Palermo, Italy.

<sup>2</sup>Dipartimento Scienze e Tecnologie Biologiche, Chimiche e Farmaceutiche, Viale delle Scienze, University of Palermo, Palermo, Italy.

\*Current address: Ge.Me.S. s.r.l., Zona Industriale, Calatafimi Segesta, Trapani, Italy.

*mecA* gene, present in the Staphylococcal cassette chromosome *mec* (SCC*mec*), is a biomarker gene responsible for resistance to methicillin and other  $\beta$ -lactam antibiotics (Peacock and Paterson, 2015; Liu *et al.*, 2016). Methicillin-resistant *S. aureus* isolates have been widely observed not only in hospitals but also in human communities with no risk factors for MRSA acquisition (Aires-De-Sousa, 2017). In recent years, several cases of MRSA transmission from pets or animals of the food chain to humans have been reported. The adaptation of MRSA clones of human origin to animal hosts has also been observed (Pomba *et al.*, 2016).

Some *S. aureus* strains produce toxins, such as superantigen staphylococcal toxic shock syndrome toxin (TSST-1), staphylococcal enterotoxins (SEs), or enterotoxin-like proteins (SEI).

Up to now, more than 20 SEs or enterotoxin-like proteins have been identified (Mehrotra *et al.*, 2000; De Buyser *et al.*, 2009; Argudín *et al.*, 2010).

The presence of SEs in food can lead to staphylococcal food poisoning, one of the most common in the world. In Sicily, food poisoning cases occur sporadically (Kadariya *et al.*, 2014). A previous screening for enterotoxigenic *S. aureus* strains isolated from food samples showed that some pathogenic *S. aureus* strains were circulating in farms with apparently healthy animals. A high percentage of the isolates (46%) carried a toxin gene, creating significant concern that pathogenic *S. aureus* strains can be transmitted through food (Vitale *et al.*, 2015).

The molecular characterization of bacterial strains is important for the detection of transmission routes and infection sources and for the monitoring of bacterial strain circulation among animal populations (Lange *et al.*, 1999; Rodriguez *et al.*, 2015; Macori *et al.*, 2017). Pulsed-field gel electrophoresis (PFGE) and multilocus sequence typing (MLST) techniques are excellent methods for epidemiological studies and for the identification of sources and transmission routes for control improvement (Golding *et al.*, 2015).

The aim of this work was the molecular characterization and the analysis of antibiotic resistance and the detection of toxin genes in *S. aureus* isolates derived from food and livestock. Four human isolates, one isolate from a dog, and one from a circus elephant were also characterized.

## Materials and Methods

### Origin and biochemical analysis of the bacterial isolates

Eighty-four *S. aureus* strains collected between 1998 and 2015 and isolated from food, animals, and humans were analyzed. Seventy-eight strains were isolated from dairy products and from animal tissue samples such as cow milk and cheese, sheep milk and cheese, goat milk, sheep skin flakes, and sheep udder. Two isolates obtained from skin flakes of other animals (elephant and dog) were used for comparison. Four human isolates were obtained from clinical samples of two individuals deceased after a food poisoning episode, which also affected four other patients (who fully recovered after severe gastroenteric symptoms). Single hemolytic colonies were inoculated in the brain/heart infusion broth (BHI) agar at 37°C. The strains were subjected to Gram staining and biochemical analysis, including coagulase, catalase, and Voges-Proskauer (VP) tests (BioMérieux), oxidase test (Oxoid), glucose and mannitol acidification in red phenol broth (Difco). The colonies had been identified as *S. aureus*

by the API STAPH test (BioMérieux). Bacteria were maintained as frozen cell glycerol stocks as described elsewhere (Giardina *et al.*, 2010; Lo Grasso *et al.*, 2015).

### PFGE analysis

Plug preparation, genomic restriction, and PFGE analysis of isolates were carried out as described in Alduina and Pisciotta (2015). In short, a single colony was inoculated into 5 mL of BHI broth and incubated at 37°C for 24 h. Cells were harvested and suspended in TE buffer (10 mM Tris HCl, 1 mM EDTA, pH 8). Three microliter lysostaphin solution (Sigma-Aldrich) (1 mg/mL in 20 mM sodium acetate) and 2% (wt/vol) SeaKem Gold agarose (Cambrex, Rockland, Maine) in TE buffer were added. The mixtures were dispensed into the wells of a small mold. Once solidified, the plugs were incubated in EC lysis buffer (6 mM Tris HCl, 1 M NaCl, 100 mM EDTA, 0.2% sodium deoxycholate, 0.5% sodium lauroyl sarcosine) at 37°C for 4 h. Plugs were washed with TE buffer three times and stored at 4°C.

The DNA was digested with 20 U *Sma*I (New England Biolabs) at room temperature for 4 h. Macrorestriction fragments were separated using a BioRad CHEF System (30" 12 h, 15" 6 h, 1% gel in Tris-borate-EDTA [TBE] 0.5 $\times$ , 200 V) and a PFGE size standard (CHEF DNA Size Standard, Lambda [ $\lambda$ ] ladder; BioRad) was added.

After run, the gels were stained with ethidium bromide (0.5  $\mu$ g/mL) and viewed under UV light. Gel images were captured by Molecular Imager Gel Doc XR (BioRad) and the banding patterns were used to establish isolate relatedness. Identical PFGE profiles (100% similarity) were defined as a pulsotype. The pulsotypes identified were given customized names PT1-10. PFGE pulsotypes were classified on the basis of the number of isolates sharing the same PT as major (more than six isolates/PFGE types), intermediate (between two and six isolates/PFGE types), or unique pulsotypes.

### Multilocus sequence typing

MLST was carried out using the protocol described in Enright *et al.* (2000) on 24 isolates of the collection. The selection was performed in such a way that at least one isolate for each pulsotype and different year, if available, could be analyzed. Polymerase chain reaction (PCR) was performed in a 30  $\mu$ L volume reaction containing 1.5 U of recombinant Taq DNA polymerase (Invitrogen, Life Technologies) as described in Randazzo *et al.* (2015).

PCR products derived from the seven housekeeping genes (*arcc*, *aroe*, *glpf*, *gmk*, *pta*, *tpi*, *yqil*) were treated with HT ExoSAP-IT (Affymetrix) following the manufacturer's instruction. The purified samples were used for sequencing using the BigDye Terminator v1.1 Cycle Sequencing Kit (Applied Biosystems) followed by capillary electrophoresis on the ABI Prism 310 Genetic Analyzer (Applied Biosystems) as described in D'Andrea *et al.* (2012). The sequences were then analyzed using the ABI3130 Genetic Analyzer (Applied Biosystems). The allelic profile for these genes was obtained from the MLST website (www.mlst.net). The combination of the seven allele numbers defines the allelic profile of the strain and each different allelic profile was assigned as a sequence type (ST), which is used to describe the strain (Aanensen and Spratt, 2005). All STs described in the study were compared with the major international *S. aureus* STs published in the MLST

TABLE 1. *STAPHYLOCOCCUS AUREUS* STRAINS ISOLATED IN SICILY ANALYZED BY MULTILOCUS SEQUENCE TYPING AND PULSED-FIELD GEL ELECTROPHORESIS

Isolate	Year	Sample	Typing by PFGE and MLST	
			PT	ST
1	1998	Elephant skin	9	ST1614
2	1998	Sheep milk	4	ST700
3	1999	Dog skin	5	ST522
4	2001	Sheep skin	7	ST700
5	2001	Sheep skin	6	ST425
6	2002	Sheep skin	5	ST1616
7	2003	Sheep milk	2	ST1626
8	2005	Sheep cheese	3	ST700
9	2006	Sheep skin	3	ST700
10	2007	Goat milk	1	ST1
11	2008	Goat milk	5	ST130
12	2008	Sheep milk	4	ST700
13	2008	Goat cheese	4	ST130
14	2008	Cow cheese	4	ST97
15	2009	Sheep milk	4	ST130
16	2010	Sheep udder	4	ST700
17	2010	Cow milk	7	ST1615
18	2012	Cow cheese	3	ST1627
19	2012	Sheep udder	8	ST522
20	2014	Goat milk	4	ST700
A	2015	Human	10	ST9
B	2015	Human	10	ST9
C	2015	Human	10	ST9
D	2015	Human	10	ST9
21	2008	Cow milk	3	NA
22	2008	Cow milk	3	NA
23	2008	Cow milk	3	NA
24	2008	Cow milk	3	NA
25	2008	Cow milk	3	NA
26	2008	Cow milk	3	NA
27	2008	Cow milk	7	NA
28	2008	Goat cheese	3	NA
29	2008	Cow milk	3	NA
30	2009	Cow milk	7	NA
31	2009	Sheep milk	4	NA
32	2010	Goat milk	3	NA
33	2010	Sheep skin	7	NA
34	2014	Sheep milk	7	NA
35	2010	Sheep milk	3	NA
36	2010	Sheep skin	5	NA
37	2010	Sheep milk	5	NA
38	2010	Sheep milk	3	NA
39	2010	Sheep skin	5	NA
40	2010	Sheep milk	5	NA
41	2010	Sheep milk	5	NA
42	2010	Sheep milk	5	NA
43	2010	Sheep milk	5	NA
44	2010	Sheep milk	5	NA
45	2010	Sheep milk	5	NA
46	2010	Sheep milk	4	NA
47	2010	Sheep milk	3	NA
48	2010	Sheep milk	8	NA
49	2010	Goat udder	4	NA
50	2008	Sheep cheese	3	NA
51	1998	Sheep udder	4	NA
52	2004	Goat cheese	4	NA
53	2007	Goat skin	8	NA

(continued)

TABLE 1. (CONTINUED)

Isolate	Year	Sample	Typing by PFGE and MLST	
			PT	ST
54	2008	Goat cheese	4	NA
55	2005	Sheep milk	8	NA
56	2005	Sheep skin	8	NA
57	2007	Sheep milk	4	NA
58	2005	Sheep milk	4	NA
59	2005	Sheep milk	3	NA
60	2005	Sheep milk	5	NA
61	2005	Sheep milk	4	NA
62	2005	Sheep milk	5	NA
63	2005	Sheep milk	4	NA
64	2005	Sheep udder	4	NA
65	2006	Sheep milk	5	NA
66	2006	Sheep milk	5	NA
67	2010	Sheep milk	3	NA
68	2002	Goat cheese	3	NA
69	2002	Goat cheese	3	NA
70	2012	Sheep milk	4	NA
71	2012	Sheep milk	4	NA
72	2012	Sheep udder	4	NA
73	2012	Sheep milk	4	NA
74	2012	Goat skin	4	NA
75	2012	Sheep skin	4	NA
76	2012	Sheep milk	4	NA
77	2012	Sheep milk	3	NA
78	2012	Sheep milk	4	NA
79	2012	Sheep milk	4	NA
80	2012	Goat milk	4	NA

Numbers 1–80 indicate isolates from animals or dairy products, letters A–D indicate the human isolates. All the strains were typed by PFGE, the first 20 randomly chosen isolates and the 4 human isolates were additionally typed using MLST. The choice of the 20 isolates was done using at least an isolate per year.

MLST, multilocus sequence typing; NA, not analyzed; PFGE, pulsed-field gel electrophoresis; PT, pulsotype; ST, sequence type.

website [www.mlst.net/databases/default.asp](http://www.mlst.net/databases/default.asp). The sequence of the new alleles was deposited in the MLST website.

*Antimicrobial susceptibility tests*

The antimicrobial susceptibility profiles to the main classes of antibiotics were determined by using the Kirby–Bauer method using Mueller–Hinton agar (MHA) medium, as described by the NCCLS (CLSI, 2015). Bacterial suspensions in BHI broth with a turbidity equivalent to a 0.5 McFarland standard were prepared and spread on the surface of MHA plates. Antibiotic disks containing the aminoglycosides gentamycin (CN, 10 µg) and kanamycin (K, 30 µg), the lincosamide lincomycin (MY, 2 µg), the macrolide erythromycin (E, 15 µg), tetracycline (TE, 30 µg), and the β-lactam antibiotics cefoperazone (CFP, 75 µg) and penicillin G (P, 10 U.I) were used. Antimicrobial disks were obtained from Oxoid (United Kingdom). The results were interpreted in accordance with the standards for inhibition zone diameters for *Staphylococcus* spp. (CLSI, 2015). *S. aureus* ATCC 25923 was used as a reference strain for antimicrobial susceptibility testing.

Detection of *SE* (*sea-see*, *seg-sei*, *sej*, *sep*), *tsst-1*, *eta*, *etb*, and *mecA* genes

Total DNA was extracted from each isolate by boiling the samples for 20 min in 1 mL of TE buffer. Two multiplex PCR assays described in Mehrotra *et al.* (2000) were used to amplify *sea-see* and *tsst-1*, *eta*, *etb*, *mecA* genes, respectively. For detection of *seg*, *seh*, *sei*, *sej*, and *sep*, a multiplex PCR assay described by De Buysse *et al.* (2009) was followed. Detection of *femA* was used as an internal positive control and *S. aureus* ATCC 25923 as quality control. For the multiplex reactions, the 25  $\mu$ L reaction mixture contained 1 U of AmpliTaq Gold 360 (Thermo Fisher Scientific), 2.5 mM MgCl<sub>2</sub>, 0.2 mM dNTPs, 1 $\times$ PCR buffer, 0.2  $\mu$ M of each primer, and 1  $\mu$ L of total DNA. PCR was performed on a 9700 Thermo cycler (Applied Biosystems). The thermal cycle for the amplification of *sea-see*, *tsst-1*, *eta*, *etb*, and *mecA* genes included the following: an initial denaturation at 94°C for 5 min, followed by 35 cycles of denaturation at 94°C for 1 min, annealing at 57°C for 1 min, extension at 72°C for 1 min 30s, and a final extension at 72°C for 7 min. The conditions for the multiplex PCR of *seg*, *seh*, *sei*, *sej*, and *sep* genes were as described above, except that the annealing step was performed at 52°C for 30 s. Positive strains carrying enterotoxin genes, kindly provided by the Italian reference laboratory for *Staphylococcus* spp., were used as controls. The following reference strains with relative genes in parenthesis were used: FRIS6 (*sea*, *seb*); FRI137 (*sec*, *seg*, *seh*, *sei*); HMPL280 (*sed*, *seg*, *sei*, *sej*, *sep*, *ser*); and FRI326 (*see*). The amplicons were detected using a 2.5% agarose gel containing ethidium bromide and visualized under ultraviolet light. The presence of a band of the expected size was considered as positivity to the presence of the corresponding gene.

## Results

### Molecular typing of *S. aureus* isolates

Eighty *S. aureus* strains isolated in Sicily from 1998 to 2014 from a collection of milk, cheese, and animal tissue

were analyzed; in addition, four human samples recovered from a food poisoning episode were added (Table 1). PFGE analysis resulted in the detection of 10 pulsotypes (PT1-PT10, Fig. 1A). Sixty-five of the isolates (PT3, -4, and -5) grouped into 3 major pulsotypes (77.4%), 15 isolates (PT7, -8, and -10) grouped into 3 intermediate pulsotypes (17.9%), and 4 isolates (PT1, -2, -6, and -9) showed unique restriction profiles.


The dendrogram (Fig. 1B) clearly shows that the human isolates (PT10) are more distant from the other isolates that share provenience and that appear to be evolved from the same clone.

MLST analysis was carried out on 18 of the 78 isolates obtained from dairy products and livestock randomly selected (including at least one for each pulsotype and each year, if available), together with the four human isolates, the isolate from the pet, and that from the exotic animal (Table 1). The analysis revealed that 12 *S. aureus* isolates belonged to 5 known allelic profiles: ST9 (four human isolates), ST130 (three isolates), ST522 (two), ST1 (one), ST97 (one), and ST425 (one). The remaining 12 isolates showed 6 new MLST combinations: ST700, ST1614, ST1615, ST1616, ST1626, and ST1627. In particular, 29% of the isolates belonged to the ST700 allelic profile. The isolate collected in 1998 from the skin sample of a circus elephant showed the new pulsotype ST1614. The isolate from dog skin belonged to the ST522 allelic profile (Table 1).

In addition, PFGE and MLST, carried out on human isolates after a severe food poisoning episode occurred in 2015, revealed that they belonged to the same pulsotype PT10 and to the same sequence type ST9 (Table 1, marked as A-D).

### Antimicrobial susceptibility

Thirty isolates (35.7%) were found to be resistant to penicillin G, 17 (20.2%) to tetracycline, 4 (5%) to gentamycin, 3 (3.75%) to lincomycin, 3 (3.75%) to cefoperazone, 2 (2.5%) to erythromycin, and 1 (1.25%) to kanamycin (Table 2). All isolates belonging to the ST700 profile resulted sensitive to



**FIG. 1.** Pulsotypes of *Staphylococcus aureus* isolates. (A) Example of the PFGE profiles after *Sma*I digestion of genomic DNA, detected in this study. (B) Dendrogram of the ten PFGE profiles.

TABLE 2. ANTIBIOTIC PROFILE AND DETECTION OF VIRULENCE GENES OF 84 *STAPHYLOCOCCUS AUREUS* ISOLATES

Isolate	Antibiotic profile							Presence of virulence genes		
	CN <i>S</i> ≥15 <i>I</i> 13–14 <i>R</i> ≤12	K <i>S</i> ≥18 <i>I</i> 14–17 <i>R</i> ≤13	MY <i>S</i> ≥21 <i>I</i> 15–20 <i>R</i> ≤14	E <i>S</i> ≥23 <i>I</i> 14–22 <i>R</i> ≤13	TE <i>S</i> ≥19 <i>I</i> 15–18 <i>R</i> ≤14	CFP <i>S</i> ≥21 <i>I</i> 16–20 <i>R</i> ≤15	P <i>S</i> ≥29 <i>I</i> — <i>R</i> ≤28	sea-see	seg-i, seg, sep	tsst, eta, etb, mecA
	1	S	I	S	I	S	I	R	ND	seg
2	S	I	S	S	S	S	S	sec	seg	tsst1
3	S	I	R	I	R	I	R	ND	seg	ND
4	S	I	S	I	S	I	S	sec	ND	tsst1
5	S	S	S	I	S	S	S	ND	seg	ND
6	S	S	S	S	R	I	R	ND	seg	ND
7	S	I	S	I	R	I	R	ND	ND	ND
8	S	I	I	I	I	S	S	ND	ND	ND
9	S	I	S	S	S	S	S	sec	ND	tsst1
10	S	I	S	I	I	S	S	ND	seg	ND
11	S	S	S	I	S	S	S	ND	seg, sep	ND
12	S	I	I	S	S	S	S	ND	seg	ND
13	S	S	S	I	S	S	S	sec	seg	tsst1
14	S	I	S	S	S	S	R	ND	ND	ND
15	S	S	S	I	S	S	S	ND	seg	ND
16	S	I	S	I	S	S	R	ND	seg	ND
17	I	I	S	I	S	S	S	ND	seg	ND
18	S	I	S	I	R	I	R	sec	ND	tsst1
19	S	S	S	I	S	I	R	ND	seg	ND
20	S	I	S	I	S	S	S	sec	seg	ND
A	S	S	S	I	I	S	I	ND	seg, sei	ND
B	S	S	S	I	I	S	I	ND	seg, sei	ND
C	S	S	S	I	I	S	I	ND	seg, sei	ND
D	S	S	S	I	I	S	I	ND	seg, sei	ND
21	S	S	S	S	S	S	S	sea	ND	ND
22	S	I	S	S	S	S	S	ND	ND	ND
23	S	S	S	S	S	S	S	ND	ND	ND
24	S	S	S	I	S	I	R	ND	ND	ND
25	S	S	S	I	S	S	S	ND	ND	ND
26	S	S	S	S	S	S	S	see	ND	ND
27	S	S	S	I	S	S	S	ND	seg	ND
28	S	S	S	S	S	S	R	ND	seg	ND
29	S	I	S	I	S	S	S	ND	seg, seg	tsst1
30	S	I	I	I	R	I	R	sec, sed	ND	eta, tsst1
31	S	S	S	S	S	S	S	sea	ND	ND
32	S	S	S	S	S	S	S	ND	seh	ND
33	I	I	R	I	R	I	R	ND	seh	eta
34	S	S	S	S	S	S	S	ND	ND	ND
35	S	S	S	S	S	S	S	ND	ND	ND
36	S	S	S	S	S	S	S	ND	ND	ND
37	S	S	S	I	S	S	S	ND	seg	ND
38	S	S	S	S	S	S	S	ND	seh	ND
39	S	S	S	I	S	I	S	ND	seh	ND
40	S	S	S	S	S	S	S	ND	ND	ND
41	S	S	S	S	S	S	S	ND	ND	ND
42	S	I	S	S	S	S	R	ND	seg	ND
43	S	S	S	S	S	S	S	ND	seg	ND
44	S	I	S	S	S	S	S	ND	ND	ND
45	S	I	S	R	R	S	S	sea	ND	ND
46	S	S	S	S	S	S	S	ND	ND	ND
47	S	S	S	S	R	S	R	ND	seg, seg	ND
48	S	S	S	S	S	S	S	ND	seg, seg	ND
49	S	S	S	S	S	S	S	ND	ND	ND
50	S	I	S	I	I	R	R	ND	ND	ND
51	S	S	S	S	S	S	S	sea, see	ND	ND
52	S	S	S	S	S	S	S	ND	ND	ND
53	S	S	S	S	S	S	S	ND	ND	ND

(continued)

TABLE 2. (CONTINUED)

Isolate	Antibiotic profile							Presence of virulence genes		
	CN	K	MY	E	TE	CFP	P	sea-see	seg-i, sej, sep	tsst, eta, etb, mecA
	<i>S</i> ≥15 <i>I</i> 13–14 <i>R</i> ≤12	<i>S</i> ≥18 <i>I</i> 14–17 <i>R</i> ≤13	<i>S</i> ≥21 <i>I</i> 15–20 <i>R</i> ≤14	<i>S</i> ≥23 <i>I</i> 14–22 <i>R</i> ≤13	<i>S</i> ≥19 <i>I</i> 15–18 <i>R</i> ≤14	<i>S</i> ≥21 <i>I</i> 16–20 <i>R</i> ≤15	<i>S</i> ≥29 <i>I</i> — <i>R</i> ≤28			
54	R	I	I	I	R	I	R	ND	ND	<i>tsst1</i>
55	S	I	S	S	S	S	S	<i>sec</i>	ND	<i>tsst1</i>
56	S	S	S	S	S	S	S	ND	ND	<i>eta</i> , <i>tsst1</i>
57	S	S	S	I	R	S	R	ND	ND	<i>mecA</i>
58	S	I	S	S	S	S	S	<i>sec</i>	ND	<i>tsst1</i>
59	S	I	I	I	I	I	R	ND	<i>seh</i>	<i>mecA</i>
60	S	I	S	R	R	S	R	<i>sec</i>	ND	<i>tsst1</i>
61	S	I	I	I	R	I	R	<i>sec</i>	ND	<i>tsst1</i>
62	S	S	S	I	S	S	S	<i>sec</i>	ND	ND
63	I	S	S	S	S	I	R	<i>sec</i>	ND	<i>tsst1</i>
64	S	S	S	S	S	S	S	<i>sec</i>	ND	<i>tsst1</i>
65	S	S	S	S	S	S	R	ND	<i>sej</i>	<i>mecA</i>
66	S	S	S	S	R	S	R	ND	<i>seh</i>	<i>mecA</i>
67	R	I	R	I	S	I	R	<i>sec</i>	ND	ND
68	S	S	S	S	R	S	R	<i>sec</i>	ND	<i>tsst1</i>
69	S	S	S	S	S	S	S	<i>sec</i>	<i>sej</i>	ND
70	I	I	I	I	R	I	R	<i>sec</i>	ND	<i>tsst1</i>
71	S	S	S	S	S	S	S	<i>sec</i>	ND	<i>tsst1</i>
72	S	I	S	S	S	S	S	<i>sec</i>	ND	<i>tsst1</i>
73	S	S	S	S	S	S	S	<i>sec</i>	ND	<i>tsst1</i>
74	S	I	S	I	R	S	R	<i>see</i>	ND	ND
75	S	S	S	S	S	S	S	<i>sec</i> , <i>see</i>	ND	<i>tsst1</i>
76	S	I	S	I	S	S	S	<i>see</i>	ND	ND
77	I	I	I	I	I	R	R	<i>sec</i> , <i>see</i>	ND	<i>tsst1</i>
78	R	S	S	S	S	S	R	<i>see</i>	ND	ND
79	R	R	S	I	S	S	R	<i>sec</i> , <i>see</i>	ND	<i>tsst1</i>
80	S	I	S	S	R	R	S	ND	ND	<i>etb</i>
A	S	S	S	I	I	S	I	ND	<i>seg</i> , <i>sei</i>	ND
B	S	S	S	I	I	S	I	ND	<i>seg</i> , <i>sei</i>	ND
C	S	S	S	I	I	S	I	ND	<i>seg</i> , <i>sei</i>	ND
D	S	S	S	I	I	S	I	ND	<i>seg</i> , <i>sei</i>	ND

Numbers 1–80 indicate isolates from animals or dairy products, letters A–D indicate the human isolates. For antibiotic profile, gentamycin (CN), kanamycin (K), lincomycin (MY), erythromycin (E), tetracycline (TE), cefoperazone (CFP), penicillin G (P) were tested. Antibiotic disk diffusion ranges (mm) for susceptible (S), intermediate (I), and resistant (R) phenotypes are given under the name of the antibiotic. For the presence of virulence genes, three different multiplex polymerase chain reactions were used to detect the genes indicated. ND indicates the analyzed genes were not detected. *se*: staphylococcal enterotoxins, *tsst1*: toxic shock syndrome toxin, *eta* and *etb*: exfoliative toxins, *mecA*: encodes the low-affinity penicillin-binding protein 2A (PBP 2A) and it determines resistance to methicillin.

gentamycin and intermediate to kanamycin, one was resistant to CFP, one to tetracycline, and one to penicillin G. Six isolates (7.1%) were classified as potentially dangerous, in that they display resistance to three different classes of antibiotics (Table 2). The human isolates showed an intermediate resistance to erythromycin, tetracycline, and penicillin G.

#### Detection of virulence genes

The presence of enterotoxin, *tsst-1*, exfoliative toxins (*eta* and *etb*), and *mecA* genes was investigated in the 84 isolates by using multiplex PCRs (Table 2). The *sec* gene was the most frequently detected ( $n=24$ , 28.6%), followed by *tsst-1* ( $n=23$ , 27.4%), *seg* ( $n=20$ , 23.8%), *sej* ( $n=8$ , 9.5%), *see* ( $n=7$ , 8.3%), *seh* ( $n=6$ , 7.1%), *sea*, *sei*, *sep*, and *mecA* ( $n=4$ , 4.8%), *eta* ( $n=3$ , 3.5%), *sed* and *etb* ( $n=1$ , 1.2%). The simultaneous presence of several toxin genes was detected in

27 isolates (Table 2). Interestingly, the four human isolates carried both *seg* and *sei* toxin genes.

#### Discussion

This is the first report of molecular genotyping, evaluation of resistance profiles, and analysis of toxin genes of *S. aureus* in bacterial isolates from dairy animals and dairy food in Sicily. Our study was carried out on a collection of *S. aureus* isolates obtained during the years 1998–2014, and it demonstrates the existence of 9 pulsotypes (Fig. 1) and 11 sequence types with high heterogeneity. MLST analysis demonstrated the presence of six new sequence types. Seven ST700 isolates were found in sheep, cow, and goat milk and udder, from different areas in Sicily. This allelic profile has been observed in Sicily for the last 16 years; in 1998 in sheep udder and in 2014 in cow milk (Table 1). The detection

of a new major clone among all isolates evidenced no host preference for animal species (sheep, cattle, and goat) and its distribution was spread all over Sicily. Among the novel profiles, ST1614 contains a new *aroe* allele that had never been previously detected in Sicily; however, it was isolated from an Indian elephant present in 1998 in an Italian circus, and so, the actual origin is unknown and we could not have any further information on the animal at this time. The ST425 and ST522 types had never been isolated in Sicily before; ST522 was found in both dog skin and in sheep udder, but with different pulsotypes, antimicrobial susceptibilities, and enterotoxin genes. Another study carried out in Spain suggested that ST522 is the most common *S. aureus* clone associated with small ruminants (Porrero *et al.*, 2012). Allelic profiles ST1 and ST97 had already been isolated as hospital-associated methicillin-resistant strains in Catania (Campanile *et al.*, 2009) and in Italian pig finishing holdings (Battisti *et al.*, 2009).

Antibiotic profiling showed a high level of penicillin (35.7%) and tetracycline (20.2%) resistance (Table 2). Resistance to penicillin remains the most common, as observed in other studies (Spanu *et al.*, 2014; Jamali *et al.*, 2015; Ferreira *et al.*, 2016). The prevalence of resistance to  $\beta$ -lactam antibiotics is frequent in *S. aureus* strains obtained from milk and related products worldwide, as reported by Daka *et al.* (2012), Hu *et al.* (2013), and Xu *et al.* (2014).

The tetracycline resistance observed in this study is more significant than that found in Italy by Spanu *et al.* (2014) in strains isolated from cheese (10.6%) or that found by Ferreira *et al.* (2016) in strains isolated from artisanal cheese (10.3%). In Sicilian farms, tetracyclines or a mix of clavulanic acid and amoxicillin is frequently used to fight infections, often without veterinarian prescriptions, thus, antibiotic resistance is likely to have increased over the years.

Molecular analysis (Table 2) showed that only four isolates contained the *mecA* gene, hinting at the circulation of methicillin resistance in dairy products. The *mecA* gene is related to methicillin resistance (Liu *et al.*, 2016).

The ST522 isolates were resistant to penicillin G, but the isolate from dog skin flakes also resulted resistant to lincosamin and tetracycline. The fact that the strain isolated from a pet showed a multiresistant phenotype (i.e., resistance to three different classes of antibiotics) confirms that multidrug resistance is easily spread among pets (Davis *et al.*, 2014). This could be due to close physical contact between pets and humans, which may allow strain transmission, or to the fact that many pets are often treated with antibiotics used in human medicine (Boost *et al.*, 2008; van Duijkeren *et al.*, 2008; Knox *et al.*, 2015).

In this study, we found that 78.5% of the isolates contained at least a toxin gene. The highest frequency was observed for the *sec* gene among classical SEs, and for the *seg* gene among the new SEs. Toxigenic strains of *S. aureus* were isolated in Sicilian healthy farms (Vitale *et al.*, 2015) and from sheep and goat cheese in Southern Italy (Basanisi *et al.*, 2016).

PFGE and MLST analyses showed the same pulsotype (PT10) and sequence type (ST9) for the four human isolates hinting that they probably belong to a single clone although they were obtained from two different individuals. ST9 was shown to be frequently spread among animals, whereas it appears to be rare among *S. aureus* isolates from human infections (Kehrenberg *et al.*, 2009). In our study, the leftover

food that probably caused the case of food poisoning resulted negative to the isolation procedures and it was not possible to identify any food source. The human isolates did not show any novel profile in the genetic analysis, however, they all showed the concurrent presence of *sei* and *seg*, not found in the other isolates. The human isolates likely carry the operon *ecg* containing *seg* and *sei* (Smyth *et al.*, 2005). The SEI toxin was shown to have a high pathogenicity, in that only 10 ng of SEI is sufficient for a lethal effect in rabbits (Roetzer *et al.*, 2016).

## Conclusions

This study showed for the first time a high heterogeneity and novelty of sequence types of *S. aureus* isolates collected in Sicily from tissues and/or dairy products from different animals between the years 1998 and 2014. Moreover, our analysis showed which *S. aureus* strains circulate in Sicily as well as a high diffusion of penicillin G and tetracycline resistance and toxin genes among the isolates. In addition, we showed that isolates obtained from patients involved in a food poisoning episode in 2015 belonged to the same allelic type ST9, and contained *sei* and *seg* toxin genes.

## Acknowledgments

This study was supported by research grants provided by the Italian Ministry of Health to M.V. (IZS SI 13/15 RC) and by the University of Palermo, Fondo Finalizzato alla Ricerca to R.A. We thank the Italian Reference Laboratory for Staphylococci for the enterotoxin-positive strains. We thank Luca Dolce for critical reading of the article.

## Disclosure Statement

No competing financial interests exist.

## References

- Aanensen DM, Spratt BG. The multilocus sequence typing network: mlst.net. *Nucleic Acids Res* 2005;33(Web Server issue):W728.
- Aires-de-Sousa M. MRSA among animals: Current overview. *Clin Microbiol Infect* 2017;23:373–380.
- Alduina R, Pisciotto A. Pulsed field gel electrophoresis and genome size estimates. *Methods Mol Biol* 2015;1231:1.
- Argudín MÁ, Mendoza MC, Rodicio MR. Food poisoning and *Staphylococcus aureus* enterotoxins. *Toxins (Basel)* 2010;2:1751–1773.
- Basanisi MG, Nobilia G, La Bella G, Russo R, Spano G, Normanno G, La Salandra G. Molecular characterization of *Staphylococcus aureus* isolated from sheep and goat cheeses in southern Italy. *Small Rumin Res* 2016;135:17.
- Battisti A, Franco A, Meriardi G, Hasman H, Iurescia M, Lorenzetti R, Feltrin F, Zini M, Aarestrup FM. Heterogeneity among methicillin-resistant *Staphylococcus aureus* from Italian pig finishing holdings. *Vet Microbiol* 2009;142:361.
- Boost MV, O'Donoghue MM, James A. Prevalence of *Staphylococcus aureus* carriage among dogs and their owners. *Epidemiol Infect* 2008;136:953.
- Campanile F, Bongiorno D, Borbone S, Stefani S. Hospital-associated methicillin-resistant *Staphylococcus aureus* (HAMRSA) in Italy. *Ann Clin Microbiol Antimicrob* 2009;8:22.

- Clinical and Laboratory Standards Institute (CLSI). *Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Fifth Informational Supplement*. CLSI document M100-S25. Wayne, PA: CLSI, 2015.
- Daka D, Solomon GS, Yihdego D. Antibiotic-resistance *Staphylococcus aureus* isolated from cow's milk in the Hawassa area, South Ethiopia. *Ann Clin Microbiol Antimicrob* 2012; 11:26.
- D'Andrea A, Martinez YZ, Alduina R, Monteverde V, Molina CF, Vitale M. Comparison of two PCR methods for detection of *Leptospira interrogans* in formalin-fixed and paraffin-embedded tissues. *Mem Inst Oswaldo Cruz* 2012;107: 85–88.
- Davis JA, Jackson CR, Fedorka-Cray PJ, Barrett JB, Brousse JH, Gustafson J, Kucher M. Carriage of methicillin-resistant staphylococci by healthy companion animals in the US. *Lett Appl Microbiol* 2014;59:1.
- De Buyser ML, Grout J, Brisabois A, Assere A, Lombard B. *Detection of Genes Encoding Staphylococcal Enterotoxins. Multiplex PCR for sea to see and ser. Method of the crl for Coagulase Positive Staphylococci Including Staphylococcus aureus*, 1<sup>st</sup> ed. Maisons-Alfort, France: CRL CPS, AFSSA, 2009; pp. 1–5.
- Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J Clin Microbiol* 2000;38:1008.
- Ferreira MA, Bernardo LG, Neves LS, Campos MR, Lamaro-Cardoso J, André MC. Virulence profile and genetic variability of *Staphylococcus aureus* isolated from artisanal cheese. *J Dairy Sci* 2016;99:8589.
- Giardina A, Alduina R, Gottardi E, Di Caro V, Süßmuth RD, Puglia AM. Two heterologously expressed *Planobispora rosea* proteins cooperatively induce *Streptomyces lividans* thiostrepton uptake and storage from the extracellular medium. *Microb Cell Fact* 2010;9:9–44.
- Golding GR, Campbell J, Spreitzer D, Chui L. Pulsed-field gel electrophoresis of *Staphylococcus aureus*. *Methods Mol Biol* 2015;1301:85.
- Gould IM, David MZ, Esposito S, Garau J, Lina G, Mazzei T, Peters G. New insights into methicillin-resistant *Staphylococcus aureus* (MRSA) pathogenesis, treatment and resistance. *Int J Antimicrob Agents* 2012;39:96–104.
- Hu SK, Liu SY, Hu WF, Zheng TL, Xu JG. Molecular biological characteristics of *Staphylococcus aureus* isolated from food. *Eur Food Res Technol* 2013;236:285.
- Jamali H, Paydar M, Radmehr B, Ismail S, Dadrasnia A. Prevalence and antimicrobial resistance of *Staphylococcus aureus* isolated from raw milk and dairy products. *Food Control* 2015;54:383.
- Kadariya J, Smith TC, Thapaliya D. *Staphylococcus aureus* and staphylococcal food-borne disease: An ongoing challenge in public health. *Biomed Res Int* 2014;2014:827965.
- Kehrenberg C, Cuny C, Strommenger B, Schwarz S, Witte W. Methicillin-resistant and -susceptible *Staphylococcus aureus* strains of clonal lineages ST398 and ST9 from swine carry the multidrug resistance gene *cfr*. *Antimicrob Agents Chemother* 2009;2:779–781.
- Knox J, Uhlemann AC, Lowy FD. *Staphylococcus aureus* infections: Transmission within households and the community. *Trends Microbiol* 2015;23:437.
- Lange C, Cardoso M, Senczek D, Schwarz S. Molecular subtyping of *Staphylococcus aureus* isolates from cases of bovine mastitis in Brazil. *Vet Microbiol* 1999;67:127.
- Liu J, Chen D, Peters BM, Li L, Li B, Xu Z, Shirliff ME. Staphylococcal chromosomal cassettes mec (SCCmec): A mobile genetic element in methicillin-resistant *Staphylococcus aureus*. *Microb Pathog* 2016;101:56–67.
- Lo Grasso L, Maffioli S, Sosio M, Bibb M, Puglia AM, Alduina R. Two master switch regulators trigger A40926 biosynthesis in *Nonomuraea sp. Strain ATCC 39727*. *J Bacteriol* 2015;197:2536.
- Macori G, Giacinti G, Bellio A, Gallina S, Bianchi DM, Sagrafoli D, Marri N, Giangolini G, Amatiste S, Decastelli L. Molecular epidemiology of methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* in the ovine dairy chain and in farm-related humans. *Toxins (Basel)* 2017;9:161.
- Mehrotra M, Wang G, Johnson WM. Multiplex PCR for detection of genes for *Staphylococcus aureus* enterotoxins, exfoliative toxins, toxic shock syndrome toxin 1, and methicillin resistance. *J Clin Microbiol* 2000;38:1032–1035.
- Peacock SJ, Paterson GK. Mechanisms of methicillin resistance in *Staphylococcus aureus*. *Annu Rev Biochem* 2015;84:577–601.
- Pomba C, Rantala M, Greko C, Baptiste KE, Catry B, van Duijkeren E, Mateus A, Moreno MA, Pyörälä S, Ružauskas M, Sanders P, Teale C, Threlfall EJ, Kunsagi Z, Torren-Edo J, Jukes H, Törneke K. Public health risk of antimicrobial resistance transfer from companion animals. *J Antimicrob Chemother* 2016;72:957–968.
- Porrero MC, Hasman H, Vela AI, Fernández-Garayzábal JF, Domínguez L, Aarestrup FM. Clonal diversity of *Staphylococcus aureus* originating from the small ruminants goats and sheep. *Vet Microbiol* 2012;156:157–161.
- Randazzo L, Montana G, Alduina R, Quatrini P, Tsantini, E, Salemi B. Flos Tectorii degradation of mortars: An example of synergistic action between soluble salts and biodeteriogens. *J Cult Herit* 2015;16:838–847.
- Rodríguez M, Hogan PG, Satola SW, Crispell E, Wylie T, Gao H, Sodergren E, Weinstock GM, Burnham CA, Fritz SA. Discriminatory indices of typing methods for epidemiologic analysis of contemporary *Staphylococcus aureus* strains. *Medicine (Baltimore)* 2015;94:e1534.
- Rodvold KA, McConeghy KW. Methicillin-resistant *Staphylococcus aureus* therapy: Past, present, and future. *Clin Infect Dis* 2014;58 Suppl 1:S20–S27.
- Roetzer A, Gruener CS, Haller G, Beyerly J, Model N, Eibl MM. Enterotoxin gene cluster-encoded SEI and SEIN from *Staphylococcus aureus* isolates are crucial for the induction of human blood cell proliferation and pathogenicity in rabbits. *Toxins (Basel)* 2016;pii:E314.
- Seegers H, Fourichon C, Beaudeau F. Production effects related to mastitis and mastitis economics in dairy cattle herds. *Vet Res* 2003;34:475.
- Sergelidis D, Angelidis AS. Methicillin-resistant *Staphylococcus aureus*: A controversial food-borne pathogen. *Lett Appl Microbiol* 2017;64:409–418.
- Smyth DS, Hartigan PJ, Meaney WJ, Fitzgerald JR, Deobald CF, Bohach GA, Smyth CJ. Superantigen genes encoded by the *egc* cluster and SaPI<sub>bov</sub> are predominant among *Staphylococcus aureus* isolates from cows, goats, sheep, rabbits and poultry. *J Med Microbiol* 2005;54:401–411.
- Spanu V, Scarano C, Cossu F, Pala C, Spanu C, De Santis EP. Antibiotic resistance traits and molecular subtyping of *Staphylococcus aureus* isolated from raw sheep milk cheese. *J Food Sci* 2014;79:M2066–M2071.
- van Duijkeren E, Houwers DJ, Schoormans A, Broekhuizen-Stins MJ, Ikawaty R, Fluit AC, Wagenaar JA. Transmission of methicillin-resistant *Staphylococcus intermedius* between humans and animals. *Vet Microbiol* 2008;128:213.



- Vitale M, Scatassa ML, Cardamone C, Oliveri G, Piraino C, Alduina R, Napoli C. Staphylococcal food poisoning case and molecular analysis of toxin genes in *Staphylococcus aureus* strains isolated from food in Sicily, Italy. *Foodborne Pathog Dis* 2015;12:21.
- Wang X, Li G, Xia X, Yang B, Xi M, Meng J. Antimicrobial susceptibility and molecular typing of methicillin-resistant *Staphylococcus aureus* in retail foods in Shaanxi, China. *Foodborne Pathog Dis* 2014;11:281–286.
- Xu J, Shi C, Song M, Xu X, Yang P, Paoli G, Shi X. Phenotypic and genotypic antimicrobial resistance traits of foodborne *Staphylococcus aureus* isolates from Shanghai. *J Food Sci* 2014;79:M635.

Address correspondence to:

*Rosa Alduina, PhD*  
*Dipartimento Scienze e Tecnologie Biologiche*  
*Chimiche e Farmaceutiche*  
*Viale delle Scienze*  
*University of Palermo*  
*Parco d'Orleans II*  
*Palermo 90128*  
*Italy*

*E-mail: valeria.alduina@unipa.it*