

Article

An Appraisal of Calcium Cyanamide as Alternative N Source for Spring-Summer and Fall Season Curly Endive Crops: Effects on Crop Performance, NUE and Functional Quality Components

Leo Sabatino ^{1,*}, Giovanni Iapichino ^{1,*}, Salvatore La Bella ¹, Teresa Tuttolomondo ¹, Fabio D'Anna ¹, Mariateresa Cardarelli ², Beppe Benedetto Consentino ¹ and Youssef Rouphael ³

¹ Department of Agricultural, Food and Forest Sciences, University of Palermo, 90128 Palermo, Italy; salvatore.labella@unipa.it (S.L.B.); teresa.tuttolomondo@unipa.it (T.T.); fabio.danna@unipa.it (F.D.); beppebenedetto.consentino@unipa.it (B.B.C.)

² Consiglio per la Ricerca in Agricoltura e L'analisi Dell'economia Agraria, Centro di Ricerca Orticoltura e Florovivaismo, 84098 Pontecagnano Faiano, Italy; mteresa.cardarelli@crea.gov.it

³ Department of Agricultural Sciences, University of Naples Federico II, 80055 Portici, Italy; youssef.rouphael@unina.it

* Correspondence: leo.sabatino@unipa.it (L.S.); giovanni.iapichino@unipa.it (G.I.); Tel.: +39-091-23862252 (L.S.); +39-091-23862215 (G.I.)

Received: 28 July 2020; Accepted: 7 September 2020; Published: 10 September 2020



Abstract: A two-year study was conducted in both spring-summer and fall seasons to evaluate calcium cyanamide (CaCN₂) as an alternative nitrogen (N) source for curly endive (*Cichorium endivia* L. var. *crispum*) grown in a Mediterranean environment. Four types of N applications were administered: (i) pre-transplanting base application of 100 kg N ha⁻¹ corresponding to 100% of the supplied N (100_{CC}), (ii) pre-transplanting base application of 50 kg N ha⁻¹ corresponding to 50% of the supplied N (50_{CC}) complemented with 50 kg N ha⁻¹ as ammonium nitrate (50_{AN}) supplied through fertigation, (iii) standard application of 100 kg N ha⁻¹ as ammonium nitrate (100_{AN}) supplied entirely through fertigation, and (iv) a N-deprived control (0 kg N ha⁻¹) used as base reference to calculate the N use efficiency indices (NUE). Fall season increased head fresh weight, head height, stem diameter and plant visual quality, compared with the spring-summer season. The CaCN₂ and standard fertigation N applications were equally effective in increasing head fresh weight and other physical parameters such as, head height, stem diameter, visual quality, number of leaves and head dry matter when compared to the unfertilized control. However, in spring-summer season, CaCN₂, especially when applied straight at 100 kg N ha⁻¹, effectively increased ascorbic acid and total phenolic content, whereas, in fall season, an increase in TSS and ascorbic acid was recorded. In both, spring-summer and fall seasons, CaCN₂ significantly decreased N content and nitrogen accumulation (N_{acc}). Furthermore, CaCN₂ pre-transplant application improved NUE indices both in terms of N fertilizer recovery efficiency and in terms of physiological efficiency of applied N. Our results finally demonstrated that NUE indices increased in the fall season as compared to the spring-summer season.

Keywords: *Cichorium endivia* L. var. *crispum*; leafy green vegetables; cultivation season; nitrogen fertilizer; CaCN₂; functional properties

1. Introduction

Curly endive (*Cichorium endivia* L. var. *crispum*) is a leafy green vegetable belonging to the *Asteraceae* family, native to the eastern Mediterranean region and probably originated from the cross between *C. intybus* and *C. pumilum* [1,2]. Curly endive is an extensively grown vegetable in Europe and is praised for its typical crunchy texture and slightly bitter taste, rendering it suitable for direct consumption or as an ingredient of mixed ready-to-eat salads. Besides, curly endive is attractive due to its significant amount of nutritional and nutraceutical compounds such as minerals, ascorbic acid, phenolics, glucosinolates and sesquiterpene lactones [3–5]. Nitrogen above-optimal fertilization of vegetable crops represents a potential risk for human health. In particular, due to its accumulation in the edible part of leafy vegetables, NO_3^- is considered an anti-nutrient [6,7] and therefore a cut back of its dietary consumption is suggested as a precautionary health care action. We assume that curly endive, although less cultivated than other leafy vegetables, is equally prone to NO_3^- accumulation in the leaves when subjected to above-optimal nitrogen fertilization. Various authors reported that calcium cyanamide (CaCN_2) might be used as a soil fumigant for managing soil-borne diseases [8], pests and viruses in the soil [9], or as a tool to impede soil acidification, and expand yield and the quality of fruits [10,11]. However, CaCN_2 is also a well-recognized nitrogen fertilizer, particularly in the vegetable crop sector, and it may represent a valuable alternative to traditional soluble fertilizers. It may also improve the crop NUE [12] through a better modulation of N supply [13]. Several researches have demonstrated that nitrogen use efficiency (NUE) and NO_3^- accumulation are affected by numerous interrelated factors such as fertilization rate and source, irrigation type, and growing season [14–20]. CaCN_2 application represents an easy, efficient and practical technique for leafy green vegetables fertilization. However, to our knowledge, there are no reports concerning the relationship between growing season and CaCN_2 fertilization doses for curly endive. Thus, the scope of our research was to assess the effects of various CaCN_2 pre-transplanting applications on yield, NUE, nutritional and functional traits of curly endive grown in spring-summer and fall seasons.

2. Materials and Methods

2.1. Plant Material, Growing Conditions and Treatments

The experiments were conducted in open field in two successive cropping seasons, spring-summer and autumn 2018 and repeated in 2019 at the Experimental Farm of the Department of Agricultural, Food and Forest Sciences of Palermo (SAAF), Sicily (Italy). Meteorological data were collected by the meteorological station of SAAF. Maximal and minimal daily temperature and rainfall in the course of the plant growth cycles were registered (Figures 1 and 2).

On 3 May and on 24 September 2018, curly endive (*Cichorium endivia* L., var. *crispum* Hegi) (var. Trusty, HM Clause, France) plug plants at the four-five true leaves stage were transplanted at a density of 10 plants m^{-2} . The soil of the experimental field was a Typic Rhodoxeralf soil made of sand, silt and clay with a percentage of 46.5, 22.3 and 31.2, respectively, and a pH of 7.2, highly rich in exchangeable K_2O , phosphorous, total nitrogen and organic matter with the following concentrations of 660 ppm, 68 ppm, 2% and 3.1%, respectively. Two doses of CaCN_2 at 100 kg N ha^{-1} (100_{CC}) or at 50 kg N ha^{-1} (50_{CC}) were applied by broadcasting on the whole plot surface and subsequently incorporating into the soil (10 cm depth) 505.0 kg ha^{-1} or 252.5 kg ha^{-1} , respectively of granular CaCN_2 (19.8% of N) (Perlka; AlzChem, Trostberg, Germany). CaCN_2 was applied 15 days before transplant. The pre-transplant implementation treatment of 100 kg N ha^{-1} (100_{CC}) (100% of the supplied nitrogen) was compared with the treatment including a pre-transplant implementation of 50 kg N ha^{-1} (50_{CC}) (50% of the supplied nitrogen) by adding to the latter an implementation of 50 kg N ha^{-1} (50_{AN}) supplied in the form of NH_4NO_3 (26% of N) by fertigation (192.3 kg ha^{-1} NH_4NO_3). These two treatments were compared with a treatment including a conventional implementation of 100 kg N ha^{-1} (100_{AN}) supplied in the form of NH_4NO_3 (26% of N) by fertigation (384.6 kg ha^{-1} NH_4NO_3). Drip irrigation was applied to all plots as per standard grower practice and in order to avoid plant

water stress. Two growing seasons (spring-summer vs. autumn) were combined with the four fertilization rates (Table 1) in a two factorial experimental design accounting for a total of eight treatments. Each treatment was made of three replicates with 10 plants per replicate and 240 plants overall. The experiment was repeated in 2019 in the same growing period.




Figure 1. Daily minimum and maximum air temperature and rainfall at Marsala, Trapani province (longitude 12°26' E, latitude 37°47' N, altitude 37 m) during the spring-summer growing season (2018 and 2019).




Figure 2. Daily minimum and maximum air temperature and rainfall at Marsala, Trapani province (longitude 12°26' E, latitude 37°47' N, altitude 37 m) during the fall growing season (2018 and 2019).

Table 1. Fertilization rates adopted for curly endive grown in spring-summer and fall seasons in a Mediterranean environment.

Nitrogen Rate (kg ha ⁻¹)	Nitrogen Source (kg ha ⁻¹)	
	CaCN ₂ (19.8% of N)	NH ₄ NO ₃ (26% of N)
0	0.0	0.0
100 _{AN}	0.0	384.6
50 _{AN} -50 _{CC}	252.2	192.3
100 _{CC}	505.0	0.0

2.2. Growth, Yield and Plant Visual Quality

Seventy-four days after transplanting curly endive plants were harvested. Afterwards, the following measurements were carried out for all the plants: head fresh weight and height, stem diameter and leaf number. At harvest, the visual quality of plants was also assessed based on a continuous score ranging from one to nine. Where one indicated a seriously damaged plant, three indicated a condition where the plant produced a usable but not saleable head, five indicated a fair condition where the head is at a marketability limit, seven indicated a good condition and nine to an excellent appearance. Five randomly selected plants from each replicate were sampled and placed in a thermo-ventilated oven (Memmert, Serie standard, Venice, Italy) to determine dry matter content. The first two days the oven was set at 80 °C and then at 105 °C until the samples reached constant dry weight.

2.3. Nutritional and Functional Traits

The qualitative analysis samples were collected as stated by Sabatino et al. [5]. Thus, all samples were collected immediately after harvest. A sample of 200 g from each replicates was placed in a commercial juicer to extract its juice that was filtered later on. Then the content of soluble solids (SSC) was assessed by a digital refractometer (MTD-045nD, Three-In-161 One Enterprises Co. Ltd., New Taipei, Taiwan). Afterwards, through potentiometric titration with NaOH (0.1 M) up to a pH of 8.1, titratable acidity (TA) was evaluated. For this purpose, 15 mL of the extracted juice was used and the results were expressed in malic acid percentage equivalent [21]. A RQflex* 10 m reflectometer (Merck, Darmstadt, Germany) and Reflectoquant Ascorbic Acid Test Strips (Merck, Darmstadt, Germany) were used to measure ascorbic acid content. Distilled water was used to dissolve the leaf juice sample (1 g), where water was added and mixed until reaching a final volume of 10 mL. Then the test strips were immersed in the prepared samples and placed into the reflectometer. The obtained results were expressed as mg of ascorbic acid per kg fresh weight. Five grams of leaf sample extracted in methanol, were used to measure total phenolic content that was assayed quantitatively by A765 and expressed as mg of caffeic acid g⁻¹ fresh weight, according to the Folin–Ciocalteu method [22] with slight modifications [23]. Nitrogen (N) content was determined from the Kjeldahl method. Particularly, a sample rate was subjected to acid-catalyzed mineralization to convert the organic nitrogen into ammoniacal nitrogen. The ammoniacal nitrogen was then distilled in an alkaline pH. The ammonia created during this distillation was collected in a boric acid solution and determined via titrimetric dosage.

2.4. Nitrogen Accumulation and Nitrogen Efficiency Indices

Total N accumulation (N_{acc}) was determined by multiplying head dry matter, expressed as kg ha⁻¹, by its percentage content of N (N_{acc} = head dry matter × Ncontent). Nitrogen efficiency indices were calculated immediately after harvest, as reported by Greenwood et al. [24]. The partial factor productivity of applied N (PFP_N) represents the harvested product (kg) per the amount of applied N (kg), where the calculation was done as follows:

$$PFP_N = \frac{YF}{NF} \quad (1)$$

where NF is the used N fertilizer (kg ha^{-1}), and YF is the obtained crop yield (kg ha^{-1}) with the implementation of a noted NF rate (kg ha^{-1}).

The agronomic efficiency of applied N (AE_N) represents the increased yield (kg) per the amount of applied N (kg), where the calculation was done as follows:

$$AE_N = \frac{(YF - Y_0)}{NF} \quad (2)$$

The apparent N fertilizer recovery efficiency (REC_N) by the crop represents the increased N_{acc} (kg) per the amount of applied N (kg), where the calculation was done as follows:

$$REC_N = \frac{(AF - A_0)}{NF} \quad (3)$$

The physiological efficiency of applied N (PE_N) represents the increased yield (kg) per the increased N_{acc} (kg) from the applied fertilizer, where the calculation was done as follows:

$$PE_N = \frac{(YF - Y_0)}{(AF - A_0)} \quad (4)$$

where YF is the crop yield (kg ha^{-1}) acquired with the implementation of a noted NF rate (kg ha^{-1}), Y_0 is the crop yield acquired without the implementation of N fertilizer, AF is plant total N_{acc} (kg ha^{-1}) at maturity in the aerial biomass when N fertilization occur, and A_0 is the corresponding plant total N_{acc} (kg ha^{-1}) at maturity in the aerial biomass when no N fertilization occur.

2.5. Experimental Design and Statistical Analysis

The SPSS software package version 14.0 (StatSoft, Inc., Chicago, IL, USA) was used to analyze the data, which were subjected to one-way analysis of variance. Tukey Honestly Significant Difference (HSD) test ($p < 0.05$) was used to separate the means when the nitrogen source was significant, and to separate the means of the treatment in case a significance was demonstrated for a particular measured characteristic. In order to evaluate the effect of the year, a preliminary two-way analysis of variance (nitrogen source \times year) was performed.

3. Results

The experiment was repeated a second year using the same experimental scheme and obtaining similar results (Table S1). Therefore, data from 2018 are presented.

3.1. Crop Performance

In spring-summer season, nitrogen-unfertilized plants had the lowest head fresh weight (Table 2). The latter was not significantly affected by the type of supplied nitrogen, as it ranged from 363.87 g in plants grown with 100_{AN} to 375.40 g in plants supplied with 50_{AN}-50_{CC}. Data collected on head height, stem diameter and visual quality, number of leaves and head dry matter supported the trend established for the head fresh weight (Table 2). Although, plants cultivated in the fall season gave higher values in terms of head fresh weight, head height, stem diameter and visual quality, number of leaves and head dry matter, tendencies recorded in fall season were perfectly in accord with those recorded in the spring-summer season.

Table 2. Effect of nitrogen source on head fresh weight, head height, stem diameter and visual quality, number of leaves and head dry matter of curly endive grown in spring-summer and fall seasons.

Nitrogen Source	Spring-Summer Season										Fall Season													
	Head Fresh Weight (g)	Head Height (cm)	Stem Diameter (mm)	Visual Quality (1–9)	Number of Leaves (No.)	Head Dry Matter (g)	Head Fresh Weight (g)	Head Height (cm)	Stem Diameter (mm)	Visual Quality (1–9)	Number of Leaves (No.)	Head Dry Matter (g)												
0	270.70	b	19.73	a	19.00	b	4.33	b	46.33	b	17.13	b	330.8	b	23.23	b	21.7	b	5.00	b	59.00	b	25.13	b
100 _{AN}	363.87	a	22.27	a	22.33	ab	7.33	a	59.67	a	27.57	a	456.8	a	27.47	a	25.7	a	9.00	a	106.00	a	44.80	a
50 _{AN} -50 _{CC}	375.40	a	22.00	a	22.67	ab	7.00	a	63.33	a	29.13	a	456.8	a	26.57	a	25.3	a	9.33	a	103.00	a	45.23	a
100 _{CC}	371.00	a	21.90	a	23.33	a	7.33	a	67.00	a	27.80	a	466.2	a	26.90	a	25.7	a	9.33	a	106.67	a	46.30	a
Significance	***		NS		*		**		**		***		***		*		**		***		***		***	

Data within a column followed by the same letter are not significantly different at $p \leq 0.05$ according to Tukey Honestly Significant Difference (HSD) Test. The significance is designated by asterisks as follows: *** statistically significant differences at p -value below 0.001; ** statistically significant differences at p -value below 0.01; * statistically significant differences at p -value below 0.05; NS, not significant. 0, no nitrogen; 100_{AN}, fertilized with ammonium nitrate at 100 kg N ha⁻¹; 50_{AN}-50_{CC}, fertilized with ammonium nitrate at 50 kg N ha⁻¹ and with calcium cyanamide at 50 kg N ha⁻¹; 100_{CC} fertilized with calcium cyanamide at 100 kg N ha⁻¹.

3.2. Nutritional Traits, Functional Properties and N Accumulation

In spring-summer season, untreated plants and 100_{AN} treated plants showed the highest values in terms of TA (Table 3). Treating endive plants with 100_{CC} and with 50_{AN}-50_{CC} decreased TA by 32.0% and 16.0%, respectively, compared to plants supplied with 100_{AN}. In terms of TA, plants grown in the fall season showed higher values than plants cultivate in the spring-summer season, however, data recorded in the fall season supported the trend established in the spring-summer season (Table 3). Nitrogen source markedly affected ascorbic acid content. Treating endive plants with 100_{CC} or with 50_{AN}-50_{CC} increased ascorbic acid by 17.6% and 8.2%, respectively, compared to plants grown with 100_{AN}. The lowest ascorbic acid content was recorded in plants grown in absence of nitrogen. In respect to ascorbic acid, data recorded in the fall season supported the trend established in spring-summer season (Table 3).

Plants cultivated in the spring-summer season and supplied with 100_{CC}, 50_{AN}-50_{CC} or 100_{AN} had a significantly higher TSS than those unfertilized. Whereas, in fall season, plants treated with 100_{AN} showed no significant statistical differences compared to the untreated ones (Table 3). Moreover, in both seasons, unfertilized plants gave the lowest TSS value. Plants from the spring-summer season supplied with 100_{CC} gave the highest total phenolic content (0.90 mg of caffeic acid g⁻¹ fw) followed by those supplied with 50_{AN}-50_{CC}. Unfertilized plants and plant fertilized with 100_{AN} gave the lowest total phenolic values, whereas, in fall season, the lowest total phenolic levels were recorded in plants from control plots. Remarkably, in fall season, plants grown on plots treated with 100_{CC}, 50_{AN}-50_{CC} or 100_{AN} did not show significant differences in terms of total phenolic content.

As regards N content, plants grown in both seasons with 100_{AN} gave the highest N content followed by those supplied with 50_{AN}-50_{CC}. In fall season, significantly lower N values were detected in plants supplied with 100_{CC}. Whereas, in spring-summer season, ANOVA and mean separation did not highlight significant differences between plants treated with 50_{AN}-50_{CC} and those treated with 100_{CC}. Finally, data on Nacc supported the trend established for leaves N content (Table 4).

Table 3. Effect of nitrogen source on titratable acidity (TA) and ascorbic acid, total soluble solid (TSS) and total phenolic of curly endive grown in spring-summer and fall seasons.

Nitroge Source	Spring-Summer Season								Fall Season							
	TA (%)		Ascorbic Acid (mg kg ⁻¹ fw)		TSS (Brix°)		Total Phenolic (mg of Caffeic Acid g ⁻¹ fw)		TA (%)		Ascorbic Acid (mg kg ⁻¹ fw)		TSS (Brix°)		Total Phenolic (mg of Caffeic Acid g ⁻¹ fw)	
0	0.933	a	75.70	d	3.73	b	0.58	c	1.17	a	73.27	c	3.40	b	0.49	b
100 _{AN}	0.833	ab	87.13	c	4.03	a	0.63	c	1.03	ab	79.67	bc	3.63	b	0.69	a
50 _{AN} -50 _{CC}	0.700	bc	94.27	b	4.10	a	0.78	b	0.90	b	83.00	b	4.00	a	0.69	a
100 _{CC}	0.567	c	102.6	a	4.20	a	0.90	a	0.67	c	92.20	a	4.07	a	0.76	a
Significance	***		***		***		***		***		***		***			*

Data within a column followed by the same letter are not significantly different at $p \leq 0.05$ according to Tukey Honestly Significant Difference (HSD) Test. The significance is designated by asterisks as follows: *** statistically significant differences at p -value below 0.001; * statistically significant differences at p -value below 0.05; NS, not significant. 0, no nitrogen; 100_{AN}, fertilized with ammonium nitrate at 100 kg N ha⁻¹; 50_{AN}-50_{CC}, fertilized with ammonium nitrate at 50 kg N ha⁻¹ and with calcium cyanamide at 50 kg N ha⁻¹; 100_{CC} fertilized with calcium cyanamide at 100 kg N ha⁻¹.

Table 4. Effect of nitrogen source on N and N_{acc} of curly endive grown in spring-summer and fall seasons.

Nitrogen Source	Spring-Summer Season				Fall Season			
	N (mg g ⁻¹ DW)		N _{acc} (kg ha ⁻¹)		N (mg g ⁻¹ DW)		N _{acc} (kg ha ⁻¹)	
0	1.97	c	3.43	c	2.34	d	5.90	d
100 _{AN}	5.70	a	15.73	a	6.70	a	30.00	a
50 _{AN} -50 _{CC}	4.13	b	12.07	b	4.49	b	20.30	b
100 _{CC}	3.83	b	10.67	b	3.36	c	15.57	c
Significance	***		***		***		***	

Data within a column followed by the same letter are not significantly different at $p \leq 0.05$ according to Tukey Honestly Significant Difference (HSD) Test. The significance is designated by asterisks as follows: *** statistically significant differences at p -value below 0.001. 0, no nitrogen; 100_{AN}, fertilized with ammonium nitrate at 100 kg N ha⁻¹; 50_{AN}-50_{CC}, fertilized with ammonium nitrate at 50 kg N ha⁻¹ and with calcium cyanamide at 50 kg N ha⁻¹; 100_{CC} fertilized with calcium cyanamide at 100 kg N ha⁻¹.

3.3. Nitrogen Efficiency Indices

In both, spring-summer and fall seasons, nitrogen source did not significantly affect PFP_N and AE_N (Table 5).

Table 5. Effect of nitrogen source on PFP_N, AE_N, PE_N and REC_N of curly endive grown in spring-summer and fall seasons.

Nitrogen Source	Spring-Summer Season				Fall Season											
	PFP _N	AE _N	PE _N	REC _N	PFP _N	AE _N	PE _N	REC _N								
100 _{AN}	27.6	a	10.4	a	83.71	b	0.124	a	44.79	a	27.67	a	103.14	c	0.266	a
50 _{AN} -50 _{CC}	29.2	a	12.1	a	138.90	a	0.087	b	45.23	a	28.12	a	166.34	b	0.170	b
100 _{CC}	27.8	a	10.7	a	146.84	a	0.072	b	46.33	a	29.22	a	239.37	a	0.123	c
Significance	NS	NS	*	***	NS	NS	***	***								

Data within a column followed by the same letter are not significantly different at $p \leq 0.05$ according to Tukey Honestly Significant Difference (HSD) Test. The significance is designated by asterisks as follows: *** statistically significant differences at p -value below 0.001; * statistically significant differences at p -value below 0.05; NS, not significant. 0, no nitrogen; 100_{AN}, fertilized with ammonium nitrate at 100 kg N ha⁻¹; 50_{AN}-50_{CC}, fertilized with ammonium nitrate at 50 kg N ha⁻¹ and with calcium cyanamide at 50 kg N ha⁻¹; 100_{CC} fertilized with calcium cyanamide at 100 kg N ha⁻¹.

The effects of tested treatment on PE_N are presented in Table 5. Plants from the spring-summer season supplied with 100_{CC} and with 50_{AN}-50_{CC} increased PE_N by 75.4% and 65.9%, respectively compared to the plants supplied without 100_{AN}. Whereas, PE_N in plants from the fall season and supplied with 100_{CC} or 50_{AN}-50_{CC} increased by 132.1% and 61.3%, respectively compared to the plants supplied with 100_{AN}. Plants both from fall and spring-summer season and fertilized with 100_{AN} had the lowest PE_N values.

The effects of the nitrogen source on REC_N are presented in Table 5. In spring-summer season, REC_N was the highest in plants cultivated with 100_{AN} and the lowest in plants cultivated with 50_{AN}-50_{CC} or 100_{CC}. In fall season, the highest REC_N values were observed in 100_{AN} treatment, whereas, the lowest values were recorded from plants from 100_{CC} treatment. Data collected in fall season supported the trend established in spring-summer season. Overall, growing endive plants during the spring-summer season substantially reduced REC_N as compared to growing plant in the autumn season. Furthermore, increasing the amount of N provided by CaCN₂ from 0 to 100% resulted in a REC_N decrease, although, in respect to the spring-summer season, ANOVA analysis did not show significant differences between plants treated with 50_{AN}-50_{CC} and those treated with 100_{CC}.

4. Discussion

In this work, we evaluated the potential use of CaCN_2 as source of nitrogen to enhance production and quality of curly endive grown in two different seasons. Our outcomes are consistent with those obtained by Adamczewska-Sowińska and Uklańska [25] who, by investigating the effects of different type and doses of nitrogen fertilization on endive, found that marketable yield increased with increasing doses of nitrogen as compared to unfertilized control. However, in our study the types of nitrogen source tested (CaCN_2 and standard fertigation) were equally effective in increasing head fresh weight and other parameters such as head height, stem diameter, visual quality, number of leaves and head dry matter when compared to unfertilized control. These results agree with those observed in lettuce, another member of the *Asteraceae* family, by Montemurro et al. [26] who, by comparing dicyandiamide, a nitrification inhibitor (NI), with urea did not find significant differences in terms of yield. Our results are also consistent with those of Di Gioia et al. [13] who found the same leaf area index, fresh yield and dry weight when comparing two lettuce types either fertilized with calcium cyanamide or with a standard split application of soluble nitrogen.

There are some reports demonstrating that applying CaCN_2 or other stabilized fertilizers that contain NIs is able to raise the yield in potato [27], soybean [28], maize [29], and wheat [30]. However, as hypothesized by Frye [31], Di Gioia et al. [13] and Sabatino et al. [11] the application of NIs may cause an increase in yield traits only at low N application rates or when N is lost by leaching and/or denitrification, causing a shortage in N level quite strong to reduce crop yield in absence of NI.

Our findings demonstrated that autumn season positively affected yield traits and plant visual quality. Currently, there is a mounting proof indicating the additive and synergetic effects of natural bioactive compounds on human health, especially from plant sources since they are able to diminish the risk of several dysfunctions linked to oxidative stress [32–34]. Our findings demonstrated that, compared with standard fertigation, a CaCN_2 pre-transplant treatment, in spring-summer season, can improve the ascorbic acid and polyphenol contents in curly endive. Whereas, the CaCN_2 pre-transplant treatment in fall season, positively effects ascorbic acid and TSS content. Our findings are coherent with those acquired by Mercelle [35], Stefanelli et al. [36] and Sabatino et al. [11], who indicated negative effects of high N administrations on nutritional and functional fruit traits. Consequently, we may hypothesize that curly endive plants in plots treated with the highest dose of CaCN_2 had higher TSS, ascorbic acid and total phenolic contents, caused by a slow nitrification process defined by CaCN_2 application and the remains of ammonium-N in the soil which can take part in reducing nutritional stresses procured by the risks of excesses or N deficiency in specific phases of the growing cycle. Moreover, our data revealed that, except for 100AN treatment, spring-summer season positively influenced total phenolic content in curly endive plants. Thus, since stress conditions prompt phenolics accumulation [37–41] and considering that the optimum growth temperature for curly endive is about 15–18 °C [42], we may hypothesize that the higher total phenolic content observed in plants from the spring-summer cycle could be due to the high temperatures (thermal stress), which are usual in the Mediterranean climate regions during spring-summer time.

Our findings showed that, in the fall season, N_{acc} decreased as the percentage of nitrogen supplied by CaCN_2 increased. This is in line with the results of Pleysier et al. [43] who, by inquiring the consequences of the application of CaCN_2 four weeks before maize and rice plantation, found that N_{acc} decreased with increasing doses of nitrogen supplied via CaCN_2 . This response was attributed by these authors to N loss by volatilization between application and planting time. Besides, our results demonstrated that autumn growing season positively affected N_{acc} compared to the spring-summer season. This could be related to the fact that autumn temperatures in the Mediterranean region, differently from the tropics, do not reach quite high levels to provoke considerable losses of ammonium-N by volatilization. Furthermore, our results are, also, in accord with those of Colonna et al. [44], who by investigating on the nutritional traits of ten leafy green vegetables harvested at two light intensities, found lower protein and $\text{NO}_3\text{-N}$ content when plants are grown at lower PAR level. Our results concerning NUE indices, either in terms of REC_N or in terms of NUE expressed as

PE_N, demonstrated that both growing seasons and nitrogen supplied via CaCN₂ play a major role. As regards the NUE expressed as PFP_N and AE_N, the application of CaCN₂ did not result in any improvement. However, PFP_N and AE_N indices were considerably higher in autumn season than in the spring-summer season. Our findings are in line with those of Chen et al. [12] and with those of Pleyrier et al. [43], who reported that CaCN₂ or other NI can potentially enhance NUE. Nevertheless, our results are partially coherent with those of Di Gioia et al. [13], who found that applying CaCN₂ improved NUE neither in terms of REC_N nor in terms of NUE expressed as PFP_N, AE_N, and PE_N. Since, the experimental fertilization schemes proposed in this study improved imperative features in curly endive, they seem to be feasible for curly endive commercial production in both spring-summer and fall seasons.

5. Conclusions

Our study suggests that a pre-transplant application of CaCN₂ can be an alternative and valuable source of N for curly endive production. Overall, CaCN₂ did not improve crop yield. Nevertheless, CaCN₂, especially when used at 100 kg N ha⁻¹ (100_{CC}), effectively increased ascorbic acid and total phenolic in spring-summer season and ascorbic acid and TSS in fall season. Simultaneously, CaCN₂ pre-transplant treatment reduced N content and N_{acc}. Furthermore, CaCN₂ improved NUE indices both in terms of REC_N and PE_N. Our results also showed that autumn season increased yield traits, plant visual quality, ascorbic acid content and NUE indices as compared to the spring-summer season.

Supplementary Materials: The following are available online at <http://www.mdpi.com/2073-4395/10/9/1357/s1>, Table S1: Significance of two-way ANOVA analysis (nitrogen source × year).

Author Contributions: Conceptualization, L.S., F.D. and G.I.; methodology, L.S., G.I. and F.D.; investigation, L.S., G.I., T.T., S.L.B. and B.B.C.; resources, F.D.; data curation, L.S., G.I., B.B.C., M.C. and Y.R.; writing—original draft preparation, L.S., G.I., M.C. and Y.R.; writing—review and editing, L.S., G.I. and Y.R.; supervision, F.D. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Kalloo, G.; Bergh, B.O. *Genetic Improvement of Vegetable Crops*; Pergamon Press: Oxford, UK, 1993; pp. 535–564.
2. Gajc-Wolska, J.; Kowalczyk, K.; Nowecka, M.; Mazur, K.; Metera, A. Effect of organic mineral fertilizers on the yield and quality of endive. *Acta Sci. Pol. Hortorum Cultus* **2012**, *11*, 189–200.
3. D’Antuono, L.F.; Ferioli, F.; Manco, M.A. The impact of sesquiterpene lactones and phenolics on sensory attributes: An investigation of a curly endive and escarole germplasm collection. *Food Chem.* **2016**, *199*, 238–245. [[CrossRef](#)]
4. Moncada, A.; Miceli, A.; Sabatino, L.; Iapichino, G.; D’Anna, F.; Vetrano, F. Effect of molybdenum rate on yield and quality of lettuce, escarole, and curly endive grown in a floating system. *Agronomy* **2018**, *8*, 171. [[CrossRef](#)]
5. Sabatino, L.; Ntatsi, G.; Iapichino, G.; D’Anna, F.; De Pasquale, C. Effect of selenium enrichment and type of application on yield, functional quality and mineral composition of curly endive grown in a hydroponic system. *Agronomy* **2019**, *9*, 207. [[CrossRef](#)]
6. Di Gioia, F.; Gonnella, M.; Santamaria, P. Contribution of leafy vegetables to dietary nitrate intake and regulations. In *Nitrate in Leafy Vegetables: Toxicity and Safety Measures*; Umar, S., Anjum, N.A., Khan, N.A., Eds.; I.K. International Publishing House Pvt. Ltd.: New Delhi/Bangalore, India, 2013; pp. 2–16.
7. Colla, G.; Kim, H.J.; Kyriacou, M.C.; Roupael, Y. Nitrates in fruits and vegetables. *Sci. Hortic.* **2018**, *237*, 221–238. [[CrossRef](#)]
8. Bletsos, F.A. Use of grafting and calcium cyanamide as alternatives to methyl bromide soil fumigation and their effects on growth, yield, quality and fusarium wilt control in melon. *J. Phytopathol.* **2005**, *153*, 155–161. [[CrossRef](#)]

9. Zhu, B.L.; Ma, J.W.; Ye, X.Z.; Xia, Z.M. Effects of lime-nitrogen on soil ameliorate and vegetables production. *J. Zhejiang Univ. Agric. Life Sci.* **2001**, *27*, 339–342.
10. Lijing, W.; Tongle, H.U.; Lijing, J.I.; Keqiang, C. Inhibitory efficacy of calcium cyanamide on the pathogens of replant diseases in strawberry. *Front. Agric. China* **2007**, *1*, 183–187.
11. Sabatino, L.; D’Anna, F.; Prinzivalli, C.; Iapichino, G. Soil solarization and calcium cyanamide affect plant vigor, yield, nutritional traits, and nutraceutical compounds of strawberry grown in a protected cultivation system. *Agronomy* **2019**, *9*, 513. [[CrossRef](#)]
12. Chen, D.; Suter, H.; Islam, A.; Edis, R.; Freney, J.R.; Walker, C.N. Prospects of improving efficiency of fertiliser nitrogen in Australian agriculture: A review of enhanced efficiency fertilisers. *Aust. J. Soil Res.* **2008**, *46*, 289–301. [[CrossRef](#)]
13. Di Gioia, F.; Gonnella, M.; Buono, V.; Ayala, O.; Cacchiarelli, J.; Santamaria, P. Calcium cyanamide effects on nitrogen use efficiency, yield, nitrates, and dry matter content of lettuce. *Agron. J.* **2017**, *109*, 354–362. [[CrossRef](#)]
14. Stone, D.A. The effect of starter fertilizers on the growth and nitrogen use efficiency of onion and lettuce. *Soil Use Manag.* **2000**, *16*, 42–48. [[CrossRef](#)]
15. Escobar-Gutierrez, A.J.; Burns, I.G.; Lee, A.; Edmondson, R.N. Screening lettuce cultivars for low nitrate content during summer and winter production. *J. Hortic. Sci. Biotechnol.* **2002**, *77*, 232–237. [[CrossRef](#)]
16. Chen, B.M.; Wang, Z.W.; Li, S.X.; Wang, G.X.; Song, H.X.; Wang, X.N. Effect of nitrate supply on plant growth, nitrate accumulation, metabolic nitrate concentration and nitrate reductase activity in three leafy vegetables. *Plant Sci.* **2004**, *167*, 635–643. [[CrossRef](#)]
17. Karam, F.; Mounzer, O.; Sarkis, F.; Lahoud, R. Yield and nitrogen recovery of lettuce under different irrigation regimes. *J. Appl. Hort.* **2002**, *4*, 70–76. [[CrossRef](#)]
18. Poh, B.L.; Gazula, A.; Simonne, E.H.; Di Gioia, F.; Hochmuth, R.R.; Alligood, M.R. Use of reduced irrigation operating pressure in irrigation scheduling: I. Effect of operating pressure, irrigation rate and nitrogen rate on drip-irrigated fresh market tomato nutritional status and yields: Implications on irrigation and fertilization management. *Horttechnology* **2011**, *21*, 14–21.
19. Burns, I.G.; Zhang, K.; Turner, M.K.; Lynn, J.; Mc Clement, S.; Hand, P.; Pink, D. Genotype and environment effects on nitrate accumulation in a diversity set of lettuce accessions at commercial maturity: The influence of nitrate uptake and assimilation, osmotic interactions and shoot weight and development. *J. Sci. Food Agric.* **2011**, *91*, 2217–2233. [[CrossRef](#)]
20. Kyriacou, M.C.; Soteriou, G.A.; Colla, G.; Roupael, Y. The occurrence of nitrate and nitrite in Mediterranean fresh salad vegetables and its modulation by preharvest practices and postharvest conditions. *Food Chem.* **2019**, *285*, 468–477. [[CrossRef](#)]
21. Han, C.; Zhao, Y.; Leonard, S.W.; Traber, M. Edible coatings to improve storability and enhance nutritional value of fresh and frozen strawberries (*Fragaria × ananassa*) and raspberries (*Rubus ideaus*). *Postharvest Biol. Technol.* **2004**, *33*, 67–78. [[CrossRef](#)]
22. Slinkard, K.; Singleton, V.L. Total phenol analysis: Automation and comparison with manual methods. *AJEV* **1997**, *28*, 49–55.
23. Rivero, R.M.; Ruiz, J.M.; Garcia, P.C.; López-Lefebvre, L.R.; Sánchez, E.; Romero, L. Resistance to cold and heat stress: Accumulation of phenolic compounds in tomato and watermelon plants. *Plant Sci.* **2001**, *160*, 315–321. [[CrossRef](#)]
24. Greenwood, D.J.; Kubo, K.; Burns, I.G.; Draycott, A. Apparent recovery of fertilizer N by vegetable crops. *Soil Sci. Plant Nutr.* **1989**, *35*, 367–381. [[CrossRef](#)]
25. Adamczewska-Sowińska, K.; Uklańska, C.M. Effect of nitrogen fertilization on yield and quality of endive. *Veg. Crop. Res Bull.* **2009**, *70*, 193–201. [[CrossRef](#)]
26. Montemurro, F.; Capotorti, G.; Lacertosa, G.; Palazzo, D. Effects of urease and nitrification inhibitors application on urea fate in soil and nitrate accumulation in lettuce. *J. Plant Nutr.* **1998**, *21*, 245–252. [[CrossRef](#)]
27. Kelling, K.A.; Wolkowski, R.P.; Ruark, M.D. Potato response to nitrogen form and nitrification inhibitors. *Am. J. Potato Res.* **2011**, *88*, 459–469. [[CrossRef](#)]
28. Kaushal, T.; Onda, M.; Ito, S.; Yamazaki, A.; Fujikake, H.; Ohtake, N.; Sueyoshi, K.; Takahashi, Y.; Ohyama, T. Effect of deep placement of slow-release fertilizer (lime nitrogen) applied at different rates on growth, N₂ fixation and yield of soya bean (*Glycine max* L. Merr.). *J. Agron. Crop Sci.* **2006**, *192*, 417–426. [[CrossRef](#)]

29. Ball-Coelho, B.R.; Roy, R.C. Enhanced ammonium sources to reduce nitrate leaching. *Nutr. Cycl. Agroecosyst.* **1999**, *54*, 73–80. [[CrossRef](#)]
30. Sharma, S.N.; Kumar, R. Effects of dicyandiamide (DCD) blended with urea on growth, yield and nutrient uptake of wheat. *J. Agric. Sci.* **1998**, *131*, 389–394. [[CrossRef](#)]
31. Frye, W. Nitrification inhibition for N efficiency and environment protection. In Proceedings of the IFA International Workshop on Enhanced-Efficiency Fertilizers, Frankfurt, Germany, 28–30 June 2005; IFA: Paris, France, 2005; pp. 1–8.
32. Roupael, Y.; Kyriacou, M.C.; Petropoulos, S.A.; De Pascale, S.; Colla, G. Improving vegetable quality in controlled environments. *Sci. Hortic.* **2018**, *234*, 275–289. [[CrossRef](#)]
33. Kyriacou, M.C.; Roupael, Y. Towards a new definition of quality for fresh fruits and vegetables. *Sci. Hortic.* **2018**, *234*, 463–469. [[CrossRef](#)]
34. Roupael, Y.; Kyriacou, M.C. Enhancing quality of fresh vegetables through salinity eustress and biofortification applications facilitated by soilless cultivation. *Front. Plant Sci.* **2018**, *9*, 1254. [[CrossRef](#)] [[PubMed](#)]
35. Mercelle, R.D. Mineral nutrition and fruit quality. *Acta Hortic.* **1995**, *383*, 219–226. [[CrossRef](#)]
36. Stefanelli, D.; Goodwin, I.; Jones, R. Minimal nitrogen and water use in horticulture: Effects on quality and content of selected nutrients. *Food Res. Int.* **2010**, *43*, 1833–1843. [[CrossRef](#)]
37. Dixon, R.A.; Paiva, N.L. Stress-induced phenylpropanoid metabolism. *Plant Cell* **1995**, *7*, 1085–1097. [[CrossRef](#)] [[PubMed](#)]
38. Moglia, A.; Lanteri, S.; Comino, C.; Acquadro, A.; de Vos, R.; Beekwilder, J. Stress induced biosynthesis of dicaffeoylquinic acids in globe artichoke. *J. Agric. Food Chem.* **2008**, *56*, 8641–8649. [[CrossRef](#)]
39. Gisbert, C.; Prohens, J.; Raigón, M.D.; Stommel, J.R.; Nuez, F. Eggplant relatives as sources of variation for developing new rootstocks: Effects of grafting on eggplant yield and fruit apparent quality and composition. *Sci. Hortic.* **2011**, *128*, 14–22. [[CrossRef](#)]
40. Maršič, N.K.; Mikulič-Petkovšek, M.; Štampar, F. Grafting influences phenolic profile and carpometric traits of fruits of greenhouse-grown eggplant (*Solanum melongena* L.). *J. Agric. Food Chem.* **2014**, *62*, 10504–10514. [[CrossRef](#)]
41. Sabatino, L.; Iapichino, G.; Rotino, G.L.; Palazzolo, E.; Mennella, G.; D’Anna, F. *Solanum aethiopicum* gr. gilo and its interspecific hybrid with *S. melongena* as alternative rootstocks for eggplant: Effects on vigor, yield, and fruit physicochemical properties of cultivar ‘Scarlati’. *Agronomy* **2019**, *9*, 223. [[CrossRef](#)]
42. Tesi, R. *Orticultura Mediterranea Sostenibile*; Pàtron Editore: Bologna, Italy, 2010.
43. Pleysier, J.L.; Arora, Y.; Juo, A.S.R. Nitrogen leaching and uptake from calcium cyanamide in comparison to urea and calcium ammonium nitrate in an Ultisol from the humid tropics. *Fertil. Res.* **1987**, *12*, 193–199. [[CrossRef](#)]
44. Colonna, E.; Roupael, Y.; Barbieri, G.; De Pascale, S. Nutritional quality of ten leafy vegetables harvested at two light intensities. *Food. Chem.* **2016**, *199*, 702–710. [[CrossRef](#)]

