

## Notulae to the Italian flora of algae, bryophytes, fungi and lichens: 9

Sonia Ravera<sup>1</sup>, Alfredo Vizzini<sup>2</sup>, Marta Puglisi<sup>3</sup>, Slavomir Adamčík<sup>4</sup>, Michele Aleffi<sup>5</sup>, Gaetano Aloise<sup>6</sup>, Fabrizio Boccardo<sup>7</sup>, Ilaria Bonini<sup>8</sup>, Miroslav Caboň<sup>4</sup>, Immacolata Catalano<sup>9</sup>, Antonio B. De Giuseppe<sup>6</sup>, Luca Di Nuzzo<sup>10</sup>, Francesco Dovana<sup>11</sup>, Zuzana Fačková<sup>4</sup>, Gabriele Gheza<sup>12</sup>, Stefano Gianfreda<sup>13</sup>, Carmine Guarino<sup>14</sup>, Anna Guttová<sup>4</sup>, Raffaello Jon<sup>15</sup>, Jří Malíček<sup>16</sup>, Mario Marziano<sup>14</sup>, Cosimo Matino<sup>13</sup>, Pier Luigi Nimis<sup>17</sup>, Giulio Pandeli<sup>18</sup>, Luca Paoli<sup>19</sup>, Nicodemo G. Passalacqua<sup>6</sup>, Elena Pittao<sup>17</sup>, Silvia Popponesi<sup>20</sup>, Domenico Puntillo<sup>6</sup>, Francesco Sguazzin<sup>21</sup>, Giovanni Sicoli<sup>22</sup>, Chiara Vallese<sup>12</sup>

**1** Dipartimento di Scienze e Tecnologie Biologiche Chimiche e Farmaceutiche (STEBICEF), Università di Palermo, Via Archirafi 38, 90123, Palermo, Italy **2** Institute for Sustainable Plant Protection (IPSP) – CNR, Viale P.A. Mattioli 25, 10125, Torino, Italy **3** Dipartimento di Scienze Biologiche, Geologiche e Ambientali, Sezione di Biologia vegetale, Università di Catania, Via A. Longo 19, 95125, Catania, Italy **4** Department of Cryptogams, Institute of Botany, Plant Science and Biodiversity Centre, Slovak Academy of Sciences, Dúbravská cesta 9, SK-845 23, Bratislava, Slovakia **5** Dipartimento di Bioscienze e Medicina veterinaria, Unità di Biodiversità Vegetale e Gestione degli Ecosistemi, Laboratorio ed Erbario di Briologia, Università di Camerino, Via Pontoni 5, 62032, Camerino (Macerata), Italy **6** Museo di Storia Naturale della Calabria ed Orto Botanico, Università della Calabria, 87036, Arcavacata di Rende (Cosenza), Italy **7** Via Filippo Bettini 14/11, 16162, Genova, Italy **8** Dipartimento di Scienze della Vita, Università di Siena, Via P. A. Mattioli, 4, 53100, Siena **9** Dipartimento di Scienze e Tecnologie, Università degli Studi di Napoli Parthenope. Centro Direzionale Isola C4, 80143, Napoli, Italy **10** Dipartimento di Biologia, Università di Firenze, Via La Pira 4, 50121, Firenze, Italy **11** Dipartimento di Scienze della Vita e Biologia dei Sistemi, Università di Torino, Viale P.A. Mattioli 25, 10125, Torino, Italy **12** Dipartimento di Scienze Biologiche, Geologiche e Ambientali (BiGeA), Università di Bologna, Via Irnerio 42, 40126, Bologna, Italy **13** Istituto d'Istruzione Secondaria Superiore “Del Prete-Falcone”, Via Mazzini s.n., Sava (Taranto), Italy **14** Dipartimento di Scienze e Tecnologie (DST), Università del Sannio, Via de Sanctis, 82100 Benevento, Italy **15** Via Manin 94A, 20861 Brugherio (Monza Brianza), Italy **16** Institute of Botany, The Czech Academy of Sciences, Zámek 1, CZ-252 43 Průhonice, Czech Republic **17** Dipartimento di Scienze della Vita, Università degli Studi di Trieste, Via L. Giorgieri 10, 34127 Trieste, Italy **18** Via Botticini 15, 50143 Firenze, Italy **19** Dipartimento di Biologia, Università di Pisa, Via Luca Ghini 13, 56126 Pisa, Italy **20** Dipartimento di Chimica, Biologia e Biotecnologia, Università di Perugia, Via del Giochetto 6, 06126 Perugia, Italy **21** Via Selvotta 61, 33055, Muzzana del Turgnano (Udine), Italy **22** Dipartimento di Biologia, Ecologia e Scienze della Terra, Università della Calabria, 87036, Arcavacata di Rende (Cosenza), Italy

Corresponding author: Sonia Ravera (sonia.ravera@unipa.it)

---

Academic editor: L. Peruzzi | Received 31 March 2020 | Accepted 24 April 2020 | Published 13 May 2020

---

**Citation:** Ravera S, Vizzini A, Puglisi M, Adamčík S, Aleffi M, Aloise G, Boccardo F, Bonini I, Caboň M, Catalano I, De Giuseppe AB, Di Nuzzo L, Dovana F, Fačková Z, Gheza G, Gianfreda S, Guarino C, Guttová A, Jon R, Malíček J, Marziano M, Martino C, Nimis PL, Pandeli G, Paoli L, Passalacqua NG, Pittao E, Poponessi S, Puntillo D, Sguazzin F, Sicoli G, Vallese C (2020) Notulae to the Italian flora of algae, bryophytes, fungi and lichens: 9. Italian Botanist 9: 35–46. <https://doi.org/10.3897/italianbotanist.9.52762>

---

## Abstract

In this contribution, new data concerning bryophytes, fungi, and lichens of the Italian flora are presented. It includes new records and confirmations for the bryophyte genera *Encalypta*, *Grimmia*, and *Riccia*, for the fungal genera *Hericium*, *Inocybe*, *Inocutis*, *Pluteus*, and *Russula*, and for the lichen genera *Bryoria*, *Farnoldia*, *Hypocenomyce*, *Lecania*, *Paracollema*, *Peltigera*, *Sarcogyne*, and *Teloschistes*.

## Keywords

Ascomycota, Basidiomycota, Bryidae, Jungermanniidae, Marchantiidae

## How to contribute

The text of the records should be submitted electronically to: Cecilia Totti (c.totti@univpm.it) for algae, Marta Puglisi (mpuglisi@unict.it) for bryophytes, Alfredo Vizzini (alfredo.vizzini@unito.it) for fungi, Sonia Ravera (sonia.ravera@unipa.it) for lichens.

## Floristic records

### Bryophytes

#### *Encalypta ciliata* Hedw. (Encalyptaceae)

+ **VEN:** Il Castello, Livinallongo Val di Lana (Belluno) on rock in wet slit (UTM WGS84 32T 724444.5153886), 1720 m, 21 August 2016, *F. Sguazzin* (Bryophytorum Herbarium F. Sguazzin). – Species confirmed for the flora of Veneto.

*Encalypta ciliata* is a Circumpolar Boreal-montane floristic element (Dierßen 2001), recorded for several countries of the Mediterranean basin (Ros et al. 2013), but its range is subcosmopolitan (Smith 2004). According to Aleffi et al. (2008), this species is known in Italy for many Administrative Regions, including Veneto, where the records have not been confirmed over the last 50 years. According to Hodgetts (2015), *E. ciliata* is considered Endangered (EN) in Estonia; Vulnerable (VU) in Czech Republic, Hungary, Montenegro and Serbia; Critically Endangered (CR) in Ireland and Luxembourg; Data Deficient (DD) in Sardinia; Near Threatened (NT) in Sicily and Not Evaluated (NE) in Italy.

S. Poponessi, F. Sguazzin, M. Aleffi

***Grimmia elongata* Kaulf. (Grimmiaceae)**

+ **EMR:** Monte Prado, Tuscan-Emilian Apennine National Park, Villa Minozzo (Reggio Emilia), on rocks (UTM WGS84: 32T 612458.4900639), 2095 m, 23 July 2018, S. Poponessi (PERU). – Species new for the flora of Emilia-Romagna.

*Grimmia elongata* is a Circumpolar Arctic-montane species with main distribution in the mountain areas of Europe, while it is very rare in southern Europe and in the Mediterranean countries (Ros et al. 2013; Hodgetts 2015). It is also known for North America, North Africa, and Asia (Muñoz and Pando 2000; Greven 2003; Ignatova and Muñoz 2004; Manju and Rajesh 2011). *Grimmia elongata* is a cryophytic, acidophytic species preferably growing on acidic sandstones, rocks and rocky crevices, north-facing outcrops and ledges. It is closely related to *Grimmia donniana* Sm., from which it is distinguished for the reddish-brown colour of the cushions, mucicous or short-awned leaves with only one margin recurved and dioecious sexuality. According to Aleffi et al. (2008), the species is known in Italy for Val d'Aosta, Piemonte, Lombardia, Trentino-Alto Adige, Campania, and Sicilia.

S. Poponessi, M. Puglisi

***Riccia beyrichiana* Hampe ex Lehm. (Ricciaceae)**

+ **TOS:** Poggio alle Monache, Chiesanuova (Firenze), on wet paths in Mediterranean maquis (UTM 32T 676341.4842470), 224 m, 8 June 2019, *G. Pandeli* (SI); Collegramole Case Nuove Impruneta (Firenze), on soil in maquis with *Erica arborea* L. and *Arbutus unedo* L., (UTM WGS84 32T 676963.4842320), 198 m, 10 April 2017, *G. Pandeli* (SIENA); Poggio Valicaia Casignano Scandicci (Firenze), on wet paths in Mediterranean maquis (UTM WGS84 32T 637816.4983201), 360 m, 15 September 2019, *G. Pandeli* (SIENA). – Species new for the flora of Toscana.

*Riccia beyrichiana* grows on acidic soils and rocks in Mediterranean forests, on cliff tops, in wetlands and grasslands in exposed sites sometimes subjected to seasonal flooding. It is distinguished from the other species of *Riccia* for its large, channelled, long-persistent, shiny thallus with swollen margins. In Italy, it is reported for Piemonte, Lombardia, Trentino-Alto Adige, Umbria, Puglia, and Sardegna (Aleffi et al. 2008) and recently for Campania (Puglisi et al. 2015a, 2015b). Its global range includes Europe, SE Asia, N Africa, and N America (Ozenoglu Kiremit et al. 2016).

G. Pandeli, I. Bonini, M. Aleffi

**Fungi*****Hericium erinaceus* (Bull.) Pers. (Hericiaceae)**

+ **CAM:** Real Bosco di Capodimonte (Napoli), on living stem of *Quercus ilex* L. (UTM WGS 84: 33T 437027.4524498) 147 m, 28 November 2019, *M. Marziano* (NAP). – Species new for the flora of Campania.

*Hericium erinaceus* is considered as rare in Italy and is reported in the Red List as Endangered (Rossi et al. 2013), while it is widespread in America and Asia (Ginns 1985). Two specimens were found on the wounds of different stems of a living tree of *Quercus ilex* L. in a holm oaks forest in a historic royal park. This saprotrophic fungus produces white rot on living trees (Boddy et al. 2011). In Italy, it is known for Calabria (Siniscalco et al. 2018), Toscana (Corana et al. 2019), Sardegna, Emilia-Romagna, and Piemonte (Bernicchia and Padovan 1997). Other congener species found in Italy are *H. coralloides* (Scop.) Pers., *H. alpestre* Pers., and *H. cirrhatum* (Pers.) Nikol. (Saitta et al. 2011).

M. Marziano, C. Guarino

### ***Inocybe pelargonium* Kühner (Inocybaceae)**

+ **LIG**: near Via S. Lorenzo di Casanova, Geminiano (Genova), on calcareous soil with *Quercus ilex* L., (UTM WGS 84: 32T 494157.4922117), 375 m, 19 October 2015, F. Boccardo (Herb. GDOR 3715). – Species new for the flora of Liguria.

*Inocybe pelargonium* is associated mainly with *Picea abies* (L.) H.Karst, *Fagus sylvatica* L. or *Quercus* L. (Bandini et al. 2019). *Inocybe pelargonium* is recognized by having strikingly distinctive characters: coppery-brownish, coppery-ochraceous to orange colours on pileus, typical smell like *Pelargonium* leaves or sweet fruits, when cut sub-spermatoc, small spores (5.8–10.0 × 3.9–5.7 µm), and relatively short cystidia (28–65 µm) variable in shape, from sub-utriform, sub-fusiform to sub-clavate, generally crystalliferous at apex (Bandini et al. 2019).

F. Dovana, F. Boccardo

### ***Inocutis levis* (P.Karst.) Y.C.Dai (Hymenochaetaceae)**

+ **ITALY (CAL)**: Orto Botanico Università della Calabria, Rende (Cosenza), on the stem of 30-year-old declining poplar trees (*Populus nigra* and *Populus nigra* subsp. *italica*) (UTM WGS84: 33S 605990.4357046), 220 m, 11 October 2018, G. Sicoli, G. Aloise, N.G. Passalacqua (CLU No. F304). – Species new for the flora of Italy (Calabria).

*Inocutis levis* is a lignicolous fungus, showing sessile and pileate basidiomata, and a distinct granular core at the point of attachment to the plant tissues (Dai 2010). The maximum size we detected were 60 cm in width, 40 cm in projection, 4 and 10 cm in context and hymenial thickness, respectively. Spores are ellipsoid, yellowish and thick-walled, very similar to those produced by *Inocutis tamaricis* (Pat.) Fiasson & Niemelä, but differing from these because they remain cyanophilous even at maturity (Dai 2010; Sicoli and Mannarino 2017). The molecular comparison (nITS rDNA sequence) of our specimen with those available in GenBank (Boudagga et al. 2017; Hashemi et al. 2017) confirmed the identification. As far as we know, and despite a number of records for Asia and Africa (Karsten 1887; Salem and Michail 1980; Dai 2010; Tura et al. 2010; Boudagga et al. 2017; Hashemi et al. 2017), no indication of this species for Europe has been reported, so far.

G. Sicoli, G. Aloise, N.G. Passalacqua

***Pluteus semibulbosus* (Lasch) Quél. (Pluteaceae)**

+ **CAL**: Botanical Garden, University of Calabria, Rende (Cosenza), on a dead branch laying on the ground, reasonably belonging to *Quercus pubescens* Willd. (UTM WGS84: 33S 605956.4357271), 220 m, 14 October 2019, G. Sicoli, A.B. De Giuseppe, N.G. Passalacqua (CLU No. F309). – Species new for the flora of Calabria.

*Pluteus semibulbosus* is an agaricoid, lignicolous, saprotrophic fungus, showing pileate and stipitate basidiomata. The pileus, less than 3.0 cm in diameter, shows a whitish-cream to pale-skin translucent cuticle, darker in the centre, striate towards the margin. The stipe is central, white, cylindrical and more or less bulbous at the base, thus differing from the close *Pluteus plautus* (Weinm.) Gillet, which lacks the bulb. Caulocystidia are characteristically cylindrical to broadly clavate to broadly fusiform, whereas they are broadly clavate to pyriform or even ventricose in *P. plautus*. Moreover, pleurocystidia are broadly lageniform in contrast with the conical-fusiform, narrowly utriform to cylindrical pleurocystidia occurring in another close species, *Pluteus inquilinus* Romagn. (Kaiguzuz et al. 2019).

G. Sicoli, A.B. De Giuseppe, N.G. Passalacqua

***Russula nympharum* F.Hampe & Marxm. (Russulaceae)**

+ **ITALY (LIG)**: Portofino Vetta (Genova), in *Quercus ilex* L. forest (UTM WGS84: 32T 513144.4908678), 30 October 2009, R. Jon (R Jon 0482); San Martino di Noce-to (Genova), (UTM WGS84: 32T 513756.4911561), *Quercus* sp., 30 October 2009, R. Jon (R Jon 0483). – Species new for the flora of Italy (Liguria).

+ **ITALY (MAR)**: Serrina S of Carpegna (Urbino), in *Quercus cerris* L. forest, (UTM WGS84: 33T 285363.4849414), 16 October 2004, leg. A. Storgaard, rev. M. Vaneková (C-F-91172). – Species new for the flora of Italy (Marche).

+ **ITALY (PIE)**: Masserano (Biella), (UTM WGS84: 32T 439663.5049156), *Quercus* sp., 15 November 2009, R. Jon (R Jon 0501). – Species new for the flora of Italy (Piemonte)

+ **ITALY (TOS)**: Marsiliana, Riserva Statale (Grosseto), forest with *Quercus suber* L. and *Q. ilex*, (UTM WGS84: 32T 646434.4763924), 8 November 2016, L. Michelin (SAV F-4887); 2 km SE of the village, Frassine, Monterotondo Marittimo (Grosseto), forest with *Q. ilex* and *Q. cerris*, (UTM WGS84: 32T 646194.4775121), 10 November 2016, M. Caboň (SAV F-4992). – Species new for the flora of Italy (Toscana).

The original description of *Russula nympharum* (Adamčík et al. 2016) was based on four collections originating from France, Spain, and Belgium. This species is distinguished from its look-alike *Russula maculata* Quél. by lower spore ornamentation not exceeding 0.6 µm, more abundant pleurocystidia, mostly cylindrical terminal cells in pileipellis and broader, clavate pileocystidia near the pileus margin, on average wider than 7 µm. Our collections represent the first reports of *R. nympharum* from Italy. According to morphological and molecular (nITS rDNA) analysis of recently collected herbarium specimens, we identified five collections of *R. nympharum* originating from

four Italian Regions. The studied collections are deposited in the herbarium of the Slovak Academy of Sciences (SAV) and in the Copenhagen Herbarium (C).

M. Caboň, S. Adamčík, R. Jon

## Lichens

### *Bryoria furcellata* (Fr.) Brodo & D.Hawksw. (Parmeliaceae)

+ **TAA**: Valle Aurina, tourist path 700 m N of Casere fraz. di Predoi (Bolzano), on bark of *Larix decidua* Mill. (UTM WGS84: 33T 281783.5215413), 1900 m, 19 July 2009, J. Malíček (Herb. Malíček no. 2323). – New for the flora of Trentino-Alto Adige.

*Bryoria furcellata* is a hair lichen usually 3–5(–12) cm long, showing regularly isotomic dichotomous branching, axils usually broad towards base and acute towards tips. It seems to be a well-delimited species characterized by lateral spinules, abundant, fissural soralia which develop tufts of spinules and by the presence of fumarprotocetraric acid (Gilbert and Hawksworth 2009). It has been reported in Italy so far only from Veneto (Nimis et al. 1991) and Friuli Venezia Giulia (Nimis 2016). It is a mainly boreal-montane, circumpolar lichen of the Northern Hemisphere, extending south to Mexico and Central America, apparently rare in the Alps (Nimis et al. 2018).

J. Malíček, S. Ravera

### *Farnoldia micropsis* (A.Massal.) Hertel

+ **VEN**: Arabba (Belluno), near top of mountain 1 km S of Passo Pordoi, on dolomite rock (UTM WGS84: 32T 715995.5151106), 2400 m, 20 July 2009, leg. J. Malíček, det. J. Hafellner (Herb. Malíček no. 2326). – Species confirmed for the flora of Veneto.

*Farnoldia micropsis* is a crustose lichen with a white areolate thallus and black apothecia (up to 2 mm wide) adnate or between the areolae, characterized by I+ blue medulla. It is a circumpolar arctic-alpine species, widespread and common throughout the Alps (Nimis et al. 2018) on calcium-bearing rocks. Nevertheless, the only record from Veneto of *F. micropsis* dates back to the second half of the 19<sup>th</sup> century (Arnold 1876, Hertel 1967).

J. Malíček, S. Ravera

### *Hypocenomyce stoehadiana* Abbassi Maaf & Cl.Roux (Ophioparmaceae)

+ **CAM**: Cratere degli Astroni (Napoli), on bark of *Quercus robur* L. (UTM WGS 84: 33T 428026.4521603) 19 m, 1 November 2016, D. Puntillo (CLU No. 17157, 17454, 17466, 17479, 17817, 17818, 17877, 17911). – Species new for the flora of Campania.

*Hypocenomyce stoehadiana* is a squamulose species with a Mediterranean-Macaronesic distribution, found on ancient specimens of *Olea* L., *Quercus ilex* L., and *Quercus virgiliana* (Ten.) Ten. in warm-humid areas. It has often sterile while at the collection site a large number of thalli is provided with fruiting bodies. According to Nimis and



Martellos (2017), this species shows an exclusively Tyrrhenian distribution in Italy, although in Calabria three populations have been found on the Ionian coast (CLU No. 5672, 5724, 17557). This lichen is classified in the Italian Red List of epiphytic lichens as “Vulnerable” (Nascimbene et al. 2013).

D. Puntillo, I. Catalano, S. Ravera

***Lecania atrynooides* M.Knowles (Ramalinaceae)**

+ **TOS:** Cala San Quirico, Populonia (Livorno), on siliceous sandstone along the coast (UTM WGS84: 32T 621304.4759203), 3 m, 8 June 2018, *L. Paoli, Z. Fačkovcová* (SAV). – Species confirmed for the flora of Toscana.

*Lecania atrynooides* is a crustose Mediterranean-Atlantic species of siliceous rocks, generally growing in areas with a humid-warm climate, such as Tyrrhenian Italy. In Toscana it was reported only from the island of Capraia (Nimis et al. 1990). The identification of the new specimen from the area of Populonia has also been confirmed by sequencing of internal transcribed spacer (ITS) regions of ribosomal DNA (ITS1-5.8S-ITS2).

L. Paoli, Z. Fačkovcová, A. Guttová

***Paracollema italicum* (B. de Lesd.) Otálora, P.M.Jørg. & Wedin (Collemaataceae)**

+ **CAM:** Santuario S.S. Annunziata, Licusati frazione di Camerota (Salerno), on *Olea europaea* L. (UTM WGS84: 33T 530453.4434790), 410 m, 4 April 2010, *S. Ravera* (Herb. Ravera); Pisciotta (Salerno), on *Olea europaea* L. (UTM WGS84: 33T 519127.4440793), 230 m, 22 February 2011, leg. *S. Ravera, G. Brunialti*, det. *S. Ravera* (Herb. Ravera); Marina di Pisciotta (Salerno), on *Olea europaea* L. (UTM WGS84: 33T 519798.4439244), 40 m, 22 February 2011, leg. *S. Ravera, G. Brunialti*, det. *S. Ravera* (Herb. Ravera). – Species new for the flora of Campania.

It is an epiphytic cyanolichen, which prefers trunks of *Olea*, *Ulmus*, *Quercus ilex* L., and *Quercus pubescens* Willd. in mesophytic woodland and exposed situations, not subjected to direct sunlight. It usually constitutes communities with *Normandina pulchella* (Borrer) Nyl. and several cyanolichens within it dominate (Degelius 1954). In Italy, it is rarely found in the Mediterranean belt. This species is presently known only for a few localities in Lazio (Ravera 2001) and in Puglia (Von Brackel 2011), while it has not been reconfirmed in its *locus classicus* in Genova (Ravera and Giordani 2008). *Paracollema italicum* is included in the Italian Red List of epiphytic lichens as “Endangered” (Nascimbene et al. 2013).

S. Ravera

***Peltigera extenuata* (Vain.) Lojka (Peltigeraceae)**

+ **LOM:** Surroundings of Caruga, Valchiavenna (Sondrio), on a drystone wall covered by mosses (UTM WGS84: 32T 525689.5132912), 1240 m, September 2017, leg. *C. Vallese*, det. *R. Benesperi, C. Vallese* (Herb. Benesperi); Nasoncio, Gerola Alta (Sondrio), on a concrete wall covered with mosses at the edge of a secondary road (UTM WGS84: 32T

542796.5102441), 1087 m, 27 August 2019, leg. G. Gheza, det. R. Benesperi, C. Vallese (Herb. Nascimbene No. JN6789, Herb. Gheza). – Species new for the flora of Lombardia.

*Peltigera extenuata* is a foliose species characterized by a thallus with (mostly) rounded lobes and flocculent and pale rhizines becoming darker in the central part. The peculiar characteristic of this species is the presence of strictly laminal soredia (Goward et al. 1995). It is a terricolous species, ecologically and morphologically similar to *Peltigera didactyla* (With.) J.R.Laundon, but it differs in having a KC+ red reaction of the medulla and soralia (Nimis 2016). In the past, it was often referred to *Peltigera didactyla* var. *extenuata* (Nyl. ex Vain.) Goffinet & Hastings, but according to Goffinet et al. (2003) this taxon is currently recognised as a species, which has been recently reported for Italy (Matteucci and Vanacore Falco 2015, Ravera et al. 2019).

C. Vallese, G. Gheza, L. Di Nuzzo

### ***Sarcogyne praetermissa* K.Knudsen & Kocourk. (Acarosporaceae)**

+ **ITALY (ABR)**: Gran Sasso Massif, pass between Santo Stefano di Sessanio and Campo Imperatore (L'Aquila), limestone, dolomite, Alpine grassland (UTM WGS84: 33T 389500.4693000), c. 1680 m, 6 April 1997, leg. P.L. Nimis, M. Tretiach (sub *Sarcogyne privigna*), rev. P.L. Nimis, 27 January 2020 (TSB No. 27188); Rocca Caramanico (Pescaia), on limestone (UTM WGS84: 33T 418500.4661500), c. 1000 m, 12 August 1996, leg. P.L. Nimis, M. Tretiach (sub *Sarcogyne privigna* v. *callicola*), rev. P.L. Nimis, 27 January 2020 (TSB No. 24943). – Species new for the flora of Italy (Abruzzo).

+ **ITALY (MAR)**: Sibillini Mnts., Monte Vettore, near Valle Orsara (Ascoli Piceno), on calcareous boulders (UTM WGS84: 33T 362600.4743400), c. 1100 m, 7 August 1996, leg. P.L. Nimis, M. Tretiach (sub *Sarcogyne privigna*), rev. P.L. Nimis, 27 January 2020 (TSB No. 24064). – Species new for the flora of Italy (Marche).

+ **ITALY (MOL)**: Below Lupara along river Biferno (Campobasso), dry calcareous rocks, cultivations (UTM WGS84: 33T 478566.4621312), c. 280 m, 4 April 1997, leg. P.L. Nimis, M. Tretiach (sub *Sarcogyne privigna*), rev. P.L. Nimis, 27 January 2020 (TSB No. 27080). – Species new for the flora of Italy (Molise).

*Sarcogyne praetermissa* is a recently-described species which was previously treated as a calcicolous morph of *Sarcogyne hypophaea* (Nyl.) Arnold (syn.: *Sarcogyne privigna* auct., see Nimis 2016), from which it differs in the unsegmented apothecial margins, the stouter paraphyses, and the growth on calcareous substrata (Knudsen and Kocourková 2018, see also Roux et al. 2019). This species, hitherto reported from central and northern Europe, and from Montenegro, seems to be particularly frequent on limestone outcrops along the Adriatic side of the Italian Peninsula, from the submediterranean to the upper montane belt, but is likely to be more widespread, and should be looked for elsewhere.

P.L. Nimis, E. Pittao

### ***Teloschistes chrysophthalmus* (L.) Th.Fr. (Teloschistaceae)**

+ **PUG**: Bosco Cuturi, Manduria (Taranto) on *Phillyrea latifolia* L. and *Quercus ilex* L. (UTM WGS84: 33T 725580.4469003) and on *Pyrus amygdaliformis* Vill. (UTM



WGS84: 33T 725621.4468806), 93 m, 19 January 2020, *S. Gianfreda* (HERB GM); Bosco Rosamarina, Manduria (Taranto), on *Phillyrea angustifolia* L (UTM WGS84: 33T 726899.4468101), 86 m, 27 January 2020, *G. Matino* (HERB GM); near Sava (Taranto) on *Prunus dulcis* (Miller) D. A. Webb and *Pistacia lentiscus* L. (UTM WGS84: 33T 713504.4472856), 110 m, 28 January 2020, *S. Gianfreda* (HERB GM). – Species confirmed for the flora of Puglia.

*Teloschistes chrysophthalmus* is a fruticose lichen; it is quite showy due to lobes being orange-yellow to grey, mostly covered with numerous marginal fibrils which are also present around the pedicellate apothecia. This lichen is included in the Italian Red List of epiphytic lichens as “Near-threatened” (Nascimbene et al. 2013). It was much more common in the past and presently it is extinct in many Regions, especially in northern Italy (Nimis 2016). The last record in Puglia dates back to the late 19<sup>th</sup> century (Jatta 1889). We found some specimens in three localities on different substrates. These lichens grow on twigs of shrubs, on the terminal part, well exposed to the sun, and never found in undergrowth, in association with *Xanthoria parietina* (L.) Th.Fr., *Physcia adscendens* H.Olivier and *Parmotrema hypoleucinum* (J.Steiner) Hale. The density of *T. chrysophthalmus* seems to be extremely limited and its distribution is irregular, probably due to human interference (tree pruning, fires, use of herbicides and fertilizers).

S. Gianfreda, C. Matino

## Acknowledgements

The studies of M. Caboň and S. Adamčík were supported by the Slovak national project APVV 15-0210.

## References

- Adamčík S, Caboň M, Eberhardt U, Saba M, Hampe F, Slovák M, Kleine J, Marxmuller H, Jančovičová S, Pfister DH, Khalid AN, Kolařík M, Marhold K, Verbeken A (2016) A molecular analysis reveals hidden species diversity within the current concept of *Russula maculata* (Russulaceae, Basidiomycota). *Phytotaxa* 270: 71–88. <https://doi.org/10.11646/phytotaxa.270.2.1>
- Aleffi M, Tacchi R, Cortini Pedrotti C (2008) Check-list of the Hornworts, Liverworts and Mosses of Italy. *Bocconea* 22: 1–255.
- Arnold F (1876) Lichenologische Ausflüge in Tirol: XVI. Ampezzo. *Verhandlungen der Zoologisch-Botanischen Gesellschaft zu Wien* 26: 389–414.
- Bandini D, Oertel B, Ploch S, Ali T, Vauras J, Schneider A, Scholler M, Eberhardt U, Thines M (2019) Revision of some central European species of *Inocybe* (Fr.) Fr. subgenus *Inocybe*, with the description of five new species. *Mycological Progress* 18: 247–294. <https://doi.org/10.1007/s11557-018-1439-9>
- Berlese AN (1896) Sulla struttura e sviluppo della *Pileolaria terebinthi*. *Rivista di Patologia Vegetale*, Firenze: 287–294.

- Bernicchia A, Padovan F (1997) Preliminary distributional data on Hericiaceae (Basidiomycetes) in Italy. *Bocconea* 5: 853–859.
- Boddy L, Crockatt ME, Ainsworth AM (2011) Ecology of *Hericium cirrhatum*, *H. coralloides* and *H. erinaceus* in the UK. *Fungal Ecology* 4: 163–173. <https://doi.org/10.1016/j.funeco.2010.10.001>
- Boudagga S, Ouali Z, Mersni M, Bouslama L, Gargano M, Venturella G, Jaouani (2017) Cultivation of some mushrooms species originating from Tunisia and exploration of their valuable metabolites. 9<sup>th</sup> International Medicinal Mushrooms Conference, Advances in Medicinal Mushroom Science: Building Bridges between Western and Eastern Medicine, September 24–28, 2017, Palermo, Italy, 131–132.
- Corana F, Cesaroni V, Mannucci B, Baiguera RM, Picco AM, Savino E, Ratto D, Perini C, Kawagishi H, Girometta CE, Rossi P (2019) Array of Metabolites in Italian *Hericium erinaceus* Mycelium, Primordium, and Sporophore. *Molecules* 24: 1–3511. <https://doi.org/10.3390/molecules24193511>
- Dai Y-C (2010) Hymenochaetaceae (Basidiomycota) in China. *Fungal Diversity* 45: 131–343. <https://doi.org/10.1007/s13225-010-0066-9>
- Degelius G (1954) The Lichen Genus *Collema* in Europe. *Morphology, Taxonomy, Ecology. Symbolae Botanicae Upsalenses* 13: 277–280.
- Dierßen K (2001) Distribution, ecological amplitude and phytosociological characterization of European bryophytes. *Bryophytorum Bibliotheca* 56: 1–289.
- Gilbert OL, Hawksworth D (2009) *Bryoria* Brodo & D. Hawksw. (1977). In: Smith CW, Aptroot A, Coppins BJ, Fletcher A, Gilbert OL, James PW, Wolseley PA (Eds) *The Lichens of Great Britain and Ireland*. The British Lichen Society, Department of Botany, The Natural History Museum, Cromwell Road, London, 1046 pp.
- GINNS J (1985) *Hericium* in North America: cultural characteristics and mating behaviour. *Canadian Journal of Botany* 63: 1551–1563. <https://doi.org/10.1139/b85-215>
- Goffinet B, Miadlikowska J, Goward T (2003) Phylogenetic Inferences Based on nrDNA Sequences Support Five Morphospecies Within the *Peltigera didactyla* Complex (Lichenized Ascomycota). *The Bryologist* 106: 349–364. <https://doi.org/10.1639/01>
- Goward T, Goffinet B, Vitikainen O (1995) Synopsis of the genus *Peltigera* (lichenized Ascomycetes) in British Columbia, with a key to the North American species. *Canadian Journal of Botany* 73: 91–111. <https://doi.org/10.1139/b95-012>
- Greven HC (2003) *Grimmias of the World*. Backhuys Publishers, Leiden, 247 pp.
- Hashemi H, Mohammadi H, Abdollahzadeh J (2017) Symptoms and fungi associated with elm trees decline in Iran. *European Journal of Forest Research* 136: 857–879. <https://doi.org/10.1007/s10342-017-1075-y>
- Hertel H (1967) Revision einiger calciphiler Formenkreise der Flechtengattung *Lecidea*. *Beihefte zur Nova Hedwigia* 24: 92–96.
- Hodgetts NG (2015) Checklist and country status of European bryophytes – towards a new Red List for Europe. *Irish Wildlife Manuals*, No. 84. National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht. Ireland, 130 pp.
- Ignatova E, Muñoz J (2004) The genus *Grimmia* Hedw. (Grimmiaceae, Musci) in Russia. *Arctoa* 13: 101–182. <https://doi.org/10.15298/arctoa.13.13>
- Jatta A (1889) *Monographia Lichenum Italiae Meridionalis*. Tip. Vecchi, Trani, 261 pp.

- Karsten PA (1887) Fungi aliquot novi in Turkestanian A Dre. Walther lecti. Hedwigia 26: 1–112.
- Kaygusuz O, Türkekul I, Knudsen H, Çolak ÖF (2019) New records of *Pluteus* section *Hispidoderma* in Turkey Based on Morphological Characteristics and Molecular Data. Phytotaxa 413: 175–206. <https://doi.org/10.11646/phytotaxa.413.3.1>
- Knudsen K, Kocourková J (2018) *Sarcogyne praetermissa* (Acarosporaceae), a new calcicolous lichen species from Europe, with a key to the European Sarcogyne species. Herzogia 31: 133–139. <https://doi.org/10.13158/099.031.0109>
- Matteucci E, Vanacore Falco I (2015) The “Serenio Ameglio” Lichen collection: a virtual exploration of exsiccata conserved in the Aosta Valley Regional Natural Science Museum. Museologia scientifica 9: 57–61.
- Manju CN, Rajesh KP (2011) *Grimmia funalis* (Schwägr.) Bruch & Schimp. (Grimmiaceae; Bryophyta) from India. Taiwania 56: 81–83.
- Muñoz J, Pando F (2000) A world synopsis of the genus *Grimmia*. Monographs in Systematic Botany from the Missouri Botanical Garden 83: 1–133.
- Nascimbene J, Nimis PL, Ravera S (2013) Evaluating the conservation status of epiphytic lichens of Italy: a red list. Plant Biosystems 147: 898–904. <https://doi.org/10.1080/11263504.2012.748101>
- Nimis PL (2016) The Lichens of Italy – A second annotated catalogue. EUT Edizioni Università di Trieste, Trieste, 740 pp.
- Nimis PL, Martellos S (2017) ITALIC – The Information System on Italian Lichens. Version 5.0. University of Trieste, Dept. of Biology, Trieste. <http://dryades.units.it/italic>
- Nimis PL, Lazzarin G, Gasparo D (1991) Lichens as bioindicators of air pollution by SO<sub>2</sub> in the Veneto Region.
- Nimis PL, Tretiach M, De Marchi A (1990) Contributions to lichen floristics in Italy. V. The lichens of the islands of Capraia (Tuscan Archipelago). Cryptogamie, Bryologie, Lichénologie 11: 1–30.
- Nimis PL, Hafellner J, Roux C, Clerc P, Mayrhofer H, Martellos S, Bilovitz PO (2018) The lichens of the Alps – an annotated checklist. MycoKeys 31: 1–634. Studia Geobotanica. An international journal 11: 3–76. <https://doi.org/10.3897/mycokeys.31.23568>
- Ozenoglu Kiremit H, Kirmaci M, Kiremit F (2016) New findings of *Riccia* species (Marchantiophyta), in Turkey and Southwest Asia. Cryptogamie, Bryologie 37: 1–8. <https://doi.org/10.7872/cryb/v37.iss1.2016.19>
- Puglisi M, Campisi P, Dia MG, Privitera M (2015a) New or interesting regional bryophyte records for Italian bryoflora. Flora Mediterranea 25 (Special Issue): 193–198. <https://doi.org/10.7320/FIMedit25SI.193>
- Puglisi M, Minissale P, Sciandrello S, Privitera M (2015b) The bryophyte vegetation of the Mediterranean temporary ponds in Italy. Plant Sociology 52: 69–78.
- Ravera S (2001) Studia lichenologica in Central Italy. I. Interesting epiphytic species from Lazio (Latium Region). Allionia 38: 187–193.
- Ravera S, Giordani P (2008) *Collema italicum* B. de Lesd. Informatore Botanico Italiano 40 suppl. 1: 143–145.
- Ravera S, Puglisi M, Vizzini A, Totti C, Aleffi M, Barberis G, Benesperi R, Brackel W, Dagnino D, De Giuseppe AB, Fačková Z, Gheza G, Giordani P, Guttová A, Mair P, Mayrhofer H, Nascimbene J, Nimis PL, Paoli L, Passalacqua NG, Pittao E, Popponesi S, Prosser F,

- Ottonello M, Puntillo D, Puntillo M, Sicoli G, Sguazzin F, Spitale D, Tratter W, Turcato C, Vallese C (2019) Notulae to the Italian flora of algae, bryophytes, fungi and lichens: 7. Italian Botanist 7: 69–91. <https://doi.org/10.3897/italianbotanist.7.34285>
- Ros RM, Mazimpaka V, Abou-Salama U, Aleffi M, Blockeel TL, Brugués M, Cros RM, Dia MG, Dirkse GM, Draper I, El-Saadawi W, Erdag A, Ganeva A, Gabriel R, Gonzales-Mancebo JM, Granger C, Herrnstadt A, Hugonnot V, Khalil K, Kürschner H, Losada-Lima A, Luís L, Mifsus S, Privitera M, Puglisi M, Sabovlijević, Sèrgio C, Shabbara HM, Sim-Sim M, Sotiaux A, Tacchi R, Vanderpoorten A, Wernner O (2013) Mosses of the Mediterranean, an annotated checklist. Cryptogamie, Bryologie 34: 99–283. <https://doi.org/10.7872/cryb.v34.iss2.2013.99>
- Rossi G, Montagnani C, Gargano D, Peruzzi L, Abeli T, Ravera S, Cogoni A, Fenu G, Magrini S, Gennai M, Foggi B, Wagensommer RP, Venturella G, Blasi C, Raimondo FM, Orsenigo S [Eds] (2013) Lista Rossa della Flora Italiana. 1. Policy Species e altre specie minacciate. Comitato Italiano IUCN e Ministero dell’Ambiente e della Tutela del Territorio e del Mare, 54 pp.
- Roux C, Poumarat S, Gueidan C, Navarro-Rosinés P, Monnat J-Y, Houmeau J-M (2019) La Acarosporaceae de Okcidenta Eŭropo. Bulletin de la Société linnéenne de Provence 70: 107–167.
- Saccardo PA (1872) Florula spontanea Horti Botanici Patavini. Nuovo Giornale botanico italiano 4: 212–220.
- Saitta A, Bernicchia A, Gorjón SP, Altobelli E, Granito VM, Losi C, Lunghini D, Maggi O, Medardi G, Padovan F, Pecoraro L, Vizzini A, Persiani AM (2011) Biodiversity of wood-decay fungi in Italy. Plant Biosystems 145(4): 958–968. <https://doi.org/10.1080/11263504.2011.633114>
- Salem MA, Michail SH (1980) *Inonotus pseudohispidus* on *Populus nigra* in Egypt. Transactions of the British Mycological Society 74: 107–110. [https://doi.org/10.1016/S0007-1536\(80\)80015-5](https://doi.org/10.1016/S0007-1536(80)80015-5)
- Scalia G (1915) Rassegna dei casi fitopatologici, studiati nel Laboratorio di Patologia vegetale della Reale Scuola Enologica di Catania nel triennio 1910–1913: relazione del direttore prof. dott. Giuseppe Scalia. Monaco & Mollica, Catania, 40 pp.
- Sicoli G, Mannarino D (2017) *Inocutis tamaricis*, ospite “balneare” nel comune di Amantea (CS). Rivista di Micologia 60: 71–78.
- Siniscalco C, Bianco PM, Lavorato C, Rotella M, Parrettini GL, Marra E, Floccia F, Campana L (Eds.) (2018) Abbinamento delle componenti micologiche della Calabria ai sistemi di classificazione degli Habitat EUNIS e Natura 2000. Prima correlazione delle specie fungine delle foreste naturali. ISPRA, Manuali e linee guida n. 180/18, 206 pp.
- Smith AJE (2004) The moss flora of Britain and Ireland. 2<sup>nd</sup> edn. Cambridge University Press, Cambridge, 1012 pp. <https://doi.org/10.1017/CBO9780511541858>
- Trotter A (1908) Flora Italica Cryptogama. Pars I: Fungi. Uredinales: 40–41.
- Ṭura D, Zmitrovich IV, Wasser SP, Nevo E (2010) Checklist of Hymenomycetes (Aphyllophorales s.l.) and Heterobasidiomycetes in Israel. Mycobiology 38: 1–256. <https://doi.org/10.4489/MYCO.2010.38.4.256>
- Von Brackel W (2011) Lichenicolous fungi and lichens from Puglia and Basilicata (southern Italy). Herzogia 24: 65–101. <https://doi.org/10.13158/hea.24.1.2011.65>