



A short proof of a theorem of Juhász

Santi Spadaro¹

Department of Mathematics, Ben Gurion University of the Negev, Be'er Sheva 84105, Israel

ARTICLE INFO

Article history:

Received 24 February 2011

Received in revised form 29 May 2011

Accepted 7 June 2011

MSC:

54A25

Keywords:

Arhangel'skii Theorem

Increasing union

Free sequence

Elementary submodel

ABSTRACT

We give a simple proof of the increasing strengthening of Arhangel'skii Theorem. Our proof naturally leads to a refinement of this result of Juhász.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

The pair (X, τ) denotes a Hausdorff topological space. In Chapter 6 of his book [6], István Juhász proves the following theorem.

Theorem 1.1. ([6, 6.11]) *Suppose (X, τ) is such that $X = \bigcup_{\alpha < \lambda} X_\alpha$, where $X_\alpha \subset X_\beta$ whenever $\alpha < \beta$ and $t(X_\alpha) \cdot \psi(X_\alpha) \cdot L(X_\alpha) \leq \kappa$ for every $\alpha < \lambda$. Then $|X| \leq 2^\kappa$.*

Where $\psi(X)$, $t(X)$ and $L(X)$ are respectively the *pseudocharacter*, the *tightness* and the *Lindelöf number* of X (see [3] or [6]). This may be considered an *increasing strengthening* (in the sense of Juhász and Szentmiklóssy [8]) of the following theorem of Arhangel'skii and Shapirovskii.

Theorem 1.2 (Arhangel'skii and Shapirovskii). ([9]) $|X| \leq 2^{\psi(X) \cdot t(X) \cdot L(X)}$.

Juhász's proof of Theorem 1.1 is five pages long. We offer a simpler and shorter proof with some help from the technique of elementary submodels. The background on this technique needed to read this paper is quite basic and can be found in the first few sections of [2].

Recall the definition of a free sequence, which already had a crucial role in the original proof [1] of Arhangel'skii's famous theorem saying that the cardinality of a first-countable Lindelöf space never exceeds the continuum (a special case of Theorem 1.2).

Definition 1.3. A set $\{x_\alpha : \alpha < \kappa\}$ is called a *free sequence of length κ* if for every $\beta < \kappa$ we have $\overline{\{x_\alpha : \alpha < \beta\}} \cap \overline{\{x_\alpha : \alpha \geq \beta\}} = \emptyset$.

E-mail addresses: santi@cs.bgu.ac.il, santispadaro@yahoo.com.

¹ Supported by the Center for Advanced Studies in Mathematics at Ben Gurion University.

As usual, we let $F(X) = \sup\{|F| : F \subset X \text{ is a free sequence}\}$. It is known that $F(X) \leq L(X) \cdot t(X)$. Indeed, assume that $L(X) \cdot t(X) \leq \kappa$. If X contained a free sequence F of size κ^+ then by $L(X) \leq \kappa$, F would have a complete accumulation point. But no complete accumulation point of a free sequence can lie in the closure of an initial segment of it. And this contradicts $t(X) \leq \kappa$.

2. The main proof

Before proving Theorem 1.1 we need a very simple lemma. Define $\Phi(X) = \sup\{L(X \setminus \{x\}) : x \in X\}$.

Lemma 2.1. $\Phi(X) = L(X) \cdot \psi(X)$.

Proof. Since any open cover of $X \setminus \{x\}$ can be extended to a cover of X by the addition of a single open set, we have $L(X) \leq \Phi(X)$. If for some $x \in X$ we have $L(X \setminus \{x\}) \leq \kappa$ then, for every $y \neq x$, select an open set U_y containing y such that $x \notin \overline{U_y}$. Then $\mathcal{U} = \{U_y : y \neq x\}$ covers $X \setminus \{x\}$ and hence we can find a subcover \mathcal{V} having cardinality $\leq \kappa$. Then $\bigcap\{X \setminus \overline{U} : U \in \mathcal{U}\} = \{x\}$, which proves that $\psi(x, X) \leq \kappa$. So, taking suprema we have that $\psi(X) \leq \Phi(X)$, and hence $\psi(X) \cdot L(X) \leq \Phi(X)$.

To prove the other direction, fix $x \in X$ suppose that $L(X) \cdot \psi(X) = \kappa$. Let \mathcal{U} be an open collection such that $|\mathcal{U}| \leq \kappa$ and $\bigcap \mathcal{U} = \{x\}$. Then $X \setminus \{x\} = \bigcup\{X \setminus U : U \in \mathcal{U}\}$ and $L(X \setminus U) \leq \kappa$ for every $U \in \mathcal{U}$. Thus $L(X \setminus \{x\}) \leq \kappa$. \square

Proof of Theorem 1.1. If $\lambda \leq 2^\kappa$ then we are done by Theorem 1.2, so we can assume that $\lambda = (2^\kappa)^+$. Let M be an elementary submodel of $H(\mu)$, where μ is a large enough regular cardinal, such that $[M]^\kappa \subset M$, $|M| = 2^\kappa$, $\{\kappa, \tau, \lambda\} \subset M$ and $\{X_\alpha : \alpha < \lambda\} \in M$. Call a set $C \subset X$ bounded if $|C| \leq 2^\kappa$.

Claim 1. If $C \in [X \cap M]^{\leq \kappa}$ then \overline{C} is bounded.

Proof of Claim 1. The proof of this claim is an extension of the proof of Subclaim 2 of Example 2.2 in [2]. We will achieve Claim 1 if we can prove that $\overline{C} \subset X \cap M$. So, suppose that this is not true and choose $p \in \overline{C} \setminus M$. For every $\theta < \lambda$ such that $p \in X_\theta$ we can use $\psi(X_\theta) \leq \kappa$ to find open sets $\{U_\alpha^\theta : \alpha < \kappa\}$ such that $X_\theta \setminus \{p\} = \bigcup_{\alpha < \kappa} X_\theta \setminus U_\alpha^\theta$. By $L(X_\theta \setminus U_\alpha^\theta) \leq \kappa$ we can find relative open sets $\{V_{\alpha\beta}^\theta : \beta < \kappa\}$ in X_θ covering $X_\theta \setminus U_\alpha^\theta$ such that $p \notin \overline{V_{\alpha\beta}^\theta}$ for every $\alpha, \beta < \kappa$.

Then we have $\overline{C} \cap (X_\theta \setminus \{p\}) = \bigcup_{\alpha, \beta < \kappa} \overline{C_{\alpha\beta}^\theta} \cap X_\theta$ for every $\theta < \lambda$, where $C_{\alpha\beta}^\theta = V_{\alpha\beta}^\theta \cap C$.

Note now that by κ -closedness of M , $C_{\alpha\beta}^\theta \in M$ for every α, β and θ . Moreover, since $p \notin M$

$$(\forall \theta \in \lambda \cap M) \left(\overline{C} \cap X_\theta \cap M = \bigcup_{\alpha, \beta < \kappa} \overline{C_{\alpha\beta}^\theta} \cap X_\theta \cap M \right).$$

So

$$M \models (\forall \theta < \lambda) \left(\overline{C} \cap X_\theta = \bigcup_{\alpha, \beta < \kappa} \overline{C_{\alpha\beta}^\theta} \cap X_\theta \right)$$

which implies

$$H(\mu) \models (\forall \theta < \lambda) \left(\overline{C} \cap X_\theta = \bigcup_{\alpha, \beta < \kappa} \overline{C_{\alpha\beta}^\theta} \cap X_\theta \right)$$

which is a contradiction because $p \in X_\theta$ for some $\theta < \lambda$. Δ

Now we claim that $X \subset M$. Suppose not and choose $p \in X \setminus M$.

Claim 2. The collection $\mathcal{U} = \{U \in M \cap \tau : p \notin U\}$ is an open cover of $X \cap M$.

Proof of Claim 2. Fix $x \in X \cap M$ and let $\mathcal{V} = \{V \in \tau : x \notin \overline{V}\}$. Note that $\mathcal{V} \in M$ and \mathcal{V} covers $X \setminus \{x\}$. Suppose you have constructed subcollections $\{\mathcal{V}_\alpha : \alpha < \beta\}$ of \mathcal{V} such that $\mathcal{V}_\alpha \in M$, $|\mathcal{V}_\alpha| \leq \kappa$ for every $\alpha < \beta$ and a set $\{x_\alpha : \alpha < \beta\} \subset X \cap M$ such that $Cl_{X \setminus \{x\}}(\{x_\alpha : \alpha < \gamma\}) \subset \bigcup_{\alpha < \gamma} \mathcal{V}_\alpha$ for every $\gamma < \beta$. By Claim 1, $Cl_{X \setminus \{x\}}(\{x_\alpha : \alpha < \beta\})$ is bounded and hence there is $\lambda_\beta < \lambda$ such that $Cl_{X \setminus \{x\}}(\{x_\alpha : \alpha < \beta\}) \subset X_{\lambda_\beta}$. By Lemma 2.1, $L(Cl_{X \setminus \{x\}}(\{x_\alpha : \alpha < \beta\})) \leq \kappa$, so there is a subcollection \mathcal{V}_β of \mathcal{V} such that $|\mathcal{V}_\beta| \leq \kappa$ and $Cl_{X \setminus \{x\}}(\{x_\alpha : \alpha < \beta\}) \subset \bigcup \mathcal{V}_\beta$. If $\bigcup_{\alpha \leq \beta} \mathcal{U}_\alpha$ does not cover $X \setminus \{x\}$ then we can fix a point $x_\beta \in ((X \setminus \{x\}) \cap M) \setminus \bigcup_{\alpha \leq \beta} \mathcal{U}_\alpha$. If the induction doesn't stop at an ordinal below κ^+ then $F = \{x_\alpha : \alpha < \kappa^+\}$ is a free sequence of length κ^+ in $X \setminus \{x\}$. Since F is bounded we can choose $\theta < \lambda$ such that $F \subset X_\theta$. Again by Lemma 2.1 we have $L(X_\theta \setminus \{x\}) \leq \kappa$. But then F cannot converge to x , because any κ^+ sized subset of a space of Lindelöf number κ has

a complete accumulation point. Therefore, there is an open neighbourhood G of x which misses κ^+ many points of F and $F \setminus G$ is a free sequence in X of cardinality κ^+ . But that contradicts $F(X_\theta) \leq \kappa$.

So there is a subcollection $\mathcal{W} \subset \mathcal{V}$ such that $|\mathcal{W}| \leq \kappa$ and $X \setminus \{x\} \subset \bigcup \mathcal{W}$. Now, by elementarity, we can assume $\mathcal{W} \in M$, but then we also have $\mathcal{W} \subset M$. Let now $W \in \mathcal{W}$ such that $p \in W$. Then the set $U = X \setminus \overline{W} \in M$ is an open neighbourhood of x such that $p \notin U$. \triangle

Suppose that for some $\beta < \kappa^+$ we have constructed a set $\{x_\alpha : \alpha < \beta\} \subset X \cap M$ and subcollections $\{\mathcal{U}_\alpha : \alpha < \beta\}$ of \mathcal{U} such that $\mathcal{U}_\alpha \in M$, $|\mathcal{U}_\alpha| \leq \kappa$ and $\{x_\gamma : \gamma < \alpha\} \subset \bigcup_{\gamma < \alpha} \mathcal{U}_\gamma$, for every $\alpha < \beta$. Since $\{x_\alpha : \alpha < \beta\}$ is bounded we have $L(\overline{\{x_\alpha : \alpha < \beta\}}) \leq \kappa$ and hence we can find a subcollection \mathcal{U}_β of \mathcal{U} of cardinality not exceeding κ such that $\overline{\{x_\alpha : \alpha < \beta\}} \subset \bigcup \mathcal{U}_\beta$. If $\bigcup_{\alpha \leq \beta} \mathcal{U}_\alpha$ does not cover X then we can pick a point $x_\beta \in X \cap M \setminus \bigcup_{\alpha \leq \beta} \mathcal{U}_\alpha$. If we didn't stop then $\{x_\alpha : \alpha < \kappa^+\}$ would be a free sequence of size κ^+ in X . But that can't happen since $\{x_\alpha : \alpha < \kappa^+\}$ is bounded. So there is a subcollection $\mathcal{V} \subset \mathcal{U}$ such that $|\mathcal{V}| \leq \kappa$ and $X \cap M \subset \bigcup \mathcal{V}$. But since $\mathcal{V} \in M$ we have that $M \models X \subset \bigcup \mathcal{V}$ and hence $H(\mu)$ thinks that there is $V \in \mathcal{V}$ such that $p \in V$. But that's a contradiction. \square

3. Remarks and acknowledgements

Note that in the proof of Theorem 1.1 we never used $t(X) \leq \kappa$, but only $F(X) \leq \kappa$. Therefore, we actually proved the increasing strengthening of the following theorem.

Theorem 3.1. $|X| \leq 2^{\psi(X) \cdot F(X) \cdot L(X)}$.

Theorem 3.1 was independently proved by Juhász [7] and the author in his dissertation [10].

Besides in the entire Chapter 6 of Juhász's book and in Juhász and Szentmiklóssy's paper [8], cardinal functions on unions of chains were also extensively studied by Mikhail Tkachenko, both in a general topological context (see [11–13]) and with Torres Falcón in the presence of a topological group structure (see [14], and also Torres Falcón's paper [15]). The study of unions of chains is very useful in obtaining reflection theorems for cardinal functions, as proved by Hajnal and Juhász's paper [4] and Hodel and Vaughan's paper [5].

In Chapter 6 of his dissertation [10] the author claimed to have a simple proof of the increasing strengthening of Theorem 3.1. That proof however still relied on two lemmas from Juhász's book [6], while the present one is more self-contained and even shorter. The author would like to thank his PhD advisor, Gary Gruenhage, for valuable discussion, István Juhász for sending him the seminar slides where Theorem 3.1 was proved, Mikhail Tkachenko for bringing the reference [13] to his attention and the Center for Advanced Studies in Mathematics at Ben Gurion University for financial support.

References

- [1] A. Arhangel'skii, The power of bicomacta with first axiom of countability, *Soviet Math. Dokl.* 10 (1969) 951–955.
- [2] A. Dow, An introduction to applications of elementary submodels to topology, *Topology Proc.* 13 (1) (1988) 17–72.
- [3] R. Engelking, *General Topology*, second ed., Sigma Ser. Pure Math., vol. 6, Heldermann Verlag, Berlin, 1989.
- [4] A. Hajnal, I. Juhász, Having a small weight is determined by the small subspaces, *Proc. Amer. Math. Soc.* 79 (4) (1980) 657–658.
- [5] R. Hodel, J. Vaughan, Reflection theorems for cardinal functions, Special issue in honor of Howard H. Wicke, *Topology Appl.* 100 (1) (2000) 47–66.
- [6] I. Juhász, Cardinal Function in Topology – Ten Years Later, *Math. Centre Tracts*, vol. 123, Mathematisch Centrum, Amsterdam, 1980.
- [7] I. Juhász, Private communication.
- [8] I. Juhász, Z. Szentmiklóssy, Increasing strengthenings of cardinal function inequalities, *Fund. Math.* 126 (3) (1986) 209–216.
- [9] B. Shapírovskii, On discrete subspaces of topological spaces; weight, tightness and Suslin number, *Soviet Math. Dokl.* 13 (1972) 215–219.
- [10] S. Spadaro, Discrete sets, free sequences and cardinal properties of topological spaces, Doctoral dissertation, Auburn University, 2009.
- [11] M.G. Tkachenko, Chains and cardinals, *Soviet Math. Dokl.* 19 (2) (1978) 382–385.
- [12] M.G. Tkachenko, The behavior of cardinal invariants under the union operation for a chain of spaces, *Moscow Univ. Math. Bull.* 33 (4) (1978) 39–46.
- [13] M.G. Tkachenko, Sum theorems for the tightness and π -character in the class of compact spaces, *Comment. Math. Univ. Carolin.* 24 (1) (1983) 51–62.
- [14] M.G. Tkachenko, Y. Torres Falcón, Unions of chains in dyadic compact spaces and topological groups, in: *Proceedings of the First Joint Japan–Mexico Meeting in Topology*, Morelia, 1999, *Topology Appl.* 121 (1–2) (2002) 25–32.
- [15] Y. Torres Falcón, Unions of chains of subgroups of a topological group, *Appl. Gen. Topol.* 2 (2) (2001) 227–235.