

Pachydema lopadusanorum n. sp. (Coleoptera Melolonthidae) from Lampedusa Island (Sicily Channel, Central Mediterranean Sea, Italy)

Ignazio Sparacio¹, Tommaso La Mantia^{2*} & Michele Bellavista³

¹Via Principe di Paternò 3, 90144 Palermo, Italy; e-mail: edizionidanaus@gmail.com

²Università degli Studi di Palermo Dip. SAAF, Viale delle Scienze ed. 4, 90128 Palermo, Italy; email: tommaso.lamantia@unipa.it

³Via C. De Grossis 7, 90135 Palermo, Italy; e-mail: michele.bellavista@gmail.com

*Corresponding author

ABSTRACT

In this paper the populations of *Pachydema* Castelnau, 1832 (Coleoptera Melolonthidae) living in Lampedusa Island (Sicily Channel, Central Mediterranean Sea, Italy), so far attributed to *P. hirticollis* (Fabricius, 1787) of North Africa, are examined. The comparison of the main morphological characters between these two populations allowed to attribute those of Lampedusa to a new species that is described in the present work. Faunistic and biological observations on these species are provided.

KEY WORDS

Scarabaeoidea; *Pachydema*; taxonomic; new species; North Africa; C-Mediterranean.

Received 11.08.2018; accepted 24.09.2018; printed 30.09.2018; published online 05.10.2018

INTRODUCTION

The genus *Pachydema* Laporte de Castelnau, 1832 (Coleoptera Melolonthidae) includes about 100 species distributed throughout the Canary Islands, North Africa, Oman (1 species) and the Middle East (Barraud, 1985; Lacroix, 2005; Král & Smetana, 2006). Many species of this genus are endemic, uncommon, and with a marked sexual dimorphism.

Among them, *P. hirticollis* (Fabricius, 1787), reported from Algeria, Tunisia, and Libya, is also present in Italy in Lampedusa and Pantelleria islands, Sicily Channel (Ragusa, 1887, 1893, 1907; Barraud, 1977, 1985; Arnone et al., 1995; Carpaneto & Piattella, 1995; Sparacio, 1995; Král & Smetana, 2006; Lacroix, 2007; Ballerio et al., 2010).

The comparison between the Lampedusa and the North African populations of *P. hirticollis* made it possible to highlight some peculiar morphological differences that allowed to attribute those from Lampedusa to a distinct species herein described.

ACRONYMS AND ABBREVIATIONS. Marcello Arnone collection, Palermo, Italy (CMA); Michele Bellavista collection, Palermo, Italy (CMB); Ignazio Sparacio collection, Palermo, Italy (CIS); Collection of Museum für Naturkunde der Humboldt Universität, Berlin, Germany (ZMHB); Collection of Museo Civico di Storia Naturale “Giacomo Doria”, Genova, Italy (MCSNG); Collection of Museum National d’Histoire Naturelle, Paris, France (MNHN); ex/x: specimen/s. Unless otherwise stated, the collector of the beetles in the field is the owner of the collection.

RESULTS

Systematics

Ordo COLEOPTERA Linnaeus, 1758
 Familia MELOLONTHIDAE Samouelle, 1819
 Subfamilia MELOLONTINAE Samouelle, 1819
 Tribus TANYPROCTINI Erichson, 1847
 Genus *Pachydema* Laporte de Castelnau, 1832

Pachydema lopadusanorum n. sp. (Fig. 1)

TYPE MATERIAL. Holotypus male: Lampedusa Island (Sicily, Italy), 13.V.1983 (CIS); paratypes, idem, 16 males and 1 female (CIS); idem, legit I. Sparacio, 3 males and 1 female (CMB); Lampedusa Island, Vallone Imbriacola, 19.IV.1987, leg. M. Arnone, 10 males (CIS); idem, 19.IV.1987, leg. M. Arnone, 42 males and 8 females (CA); idem, 18.II.2017, legit T. La Mantia, 2 females (CIS); Lampedusa 1902 - Horn; Lampedusa [V.1886 leg. L. Failla], 4 males and 1 female, collection E. Ragusa (CMC).

OTHER EXAMINED MATERIAL. *Pachydema hirticollis* (Fig. 2). Tunisia. Tunis - quedenfeldti, Monastir, Brsk. - Coll. Brenske, 1 ex (ZMHB, syntype Collection Brenske of *P. quedenfeldti* Brenske, 1889). Tunisia, Kamait, 23.IV.1882, G. e L. Doria, Museo Civico di Genova, 7 males and 3 females (MCSNG). Tunisia, 4.VI.1972, G. Fiori, Museo Genova ex coll. G. Fiori (1987), 2 females (MCSNG). Biserta, 23.IV.1994, leg. F. Tagliaferri, 8 males and 1 females (MCSNG). Capo Bon, Korbous, 15.V.1987, 3 males and 1 female (CIS); idem, 26 e 29.IV.1998, 108 males and 18 females (CIS); Sousse, 5.V.1992, 2 females (CIS).

Pachydema demoflysi Normand, 1938.
 Tunisia, Sbeitla, 16.V.1996, H. Pierotti, 3 males (MCSNG).

DESCRIPTION OF HOLOTYPE. Male. Length 12 mm. Head, antennae, palpi, legs, pronotum, and ventral surface black in colour; elytrae red. Pubescence long and yellowish, very dense on the ventral surface.

Head with deep and dense punctures and long and yellowish pubescence. Clypeus wider than long (width clypeus/length clypeus = 1.9), fore margin high, slightly emarginated in the middle, with large, very dense and irregular punctures. An-

tennae 10-segmented (Fig. 3), club 5-segmented, shorter of the funicle; scape elongated and dilated distally, 2° segment short and roundish, shorter than the 3°; 3°–4° elongated; 5° short and transverse. Maxillary palpi with last article fusiform with pointed apex.

Pronotum convex, transverse (width pronotum/length pronotum = 2.1), sub-trapezoidal, maximum width at its posterior third, covered by long and yellowish pubescence, that of the fore margin is directed mainly backwards; pronotal lateral margins subrectilinear in the posterior half after narrow and straight forward; the fore margin is slightly protruding at the middle with fore angles normally shaped; the pronotal bases is widely protruding in the backwards, posterior angles little obtuse; punctures large, deep, and dense.

Scutellum triangular with slightly arched sides, and with a line of large and confluent punctures on the sides. Elytrae a little elongated, slightly rounded on the sides with maximum width at posterior third and yellowish setae on the lateral margins; punctures large and dense, confluent on disc; interstriae obsolete, the 1° is flat, hardly visible. Humeral callus visible.

Anterior tibiae tridentate, with a little basal tooth, 2° and 3° tarsomeres flat and rounded (Fig. 5). Posterior tibiae enlarged at the middle of the inferior margin and, after, strongly dilated towards the apex; posterior tarsi longer than tibiae (length tarsi/length tibiae = 1.5). First posterior tarsomere shorter than the 3° tarsomere. Claws bifid.

Pygidium pointed-rounded at apex, finely micro-reticulated with small punctures. Metathoracic wings fully developed.

Aedeagus (Fig. 9) rather long and robust, the parameres apices (in lateral view) are a little elongated and a little pointed.

VARIABILITY. The paratypes males have no substantial morphological differences compared to the holotype. Length 11–13.2 mm. The 1° elytral interstria is always hardly visible, often absent distally. The paratype females are bigger (length 14–16.5), with a more convex dorsum, wider distally, reddish-brown in colour, sometimes very dark; antennae and tarsi shorter; punctures of the pronotum large, deep and dense; elytrae smooth with punctures very small and spaced. Propygidium with small punctures on microreticulated surface; pygidium with punctures medium sized and dense.




Figure 1. *Pachydemalopadusanorum* n. sp. from Lampedusa island (length 11.8 mm). Figure 2. *Pachydemahirticollis* from Tunisia, Capo Bon, Korbous (length 12.4 mm). Figure 3. Antennae of *P. lopadusanorum* n. sp. Figure 4. Antennae of *P. hirticollis*. Figure 5. Anterior tarsomeres of *P. lopadusanorum* n. sp. Figure 6. Anterior tarsomeres of *P. hirticollis*. Figure 7. Syotype of *P. quedenfeldti* (from Tunisia, Monastir).

ETYMOLOGY. Latin name of type locality: “*lopadusanorum*”, that live in Lampedusa. The species is named after the people from Lampedusa island (Sicily Channel, Italy), and in particular to Vincenzo Billeci, Giuseppe Maraventano, Elena Prazzi, Francesco Sanguedolce, and Gerry Sorrentino, who fight for the conservation of nature and human values on that small patch of land surrounded by the sea.

DISTRIBUTION AND BIOLOGY. *Pachydema lopadusanorum* n. sp. is endemic to Lampedusa Island (Sicily Channel, Central Mediterranean Sea, Italy). Adults are active in spring and summer and can be found in flight or on flowers (Sparacio, 1995; Arnone et al., 1995; Ballerio et al., 2010; all sub *P. hirticollis*).

STATUS AND CONSERVATION. The few populations and restricted distribution makes *P. lopadusanorum* n. sp. as “Vulnerable”, according to the Categories and Criteria of the IUCN Red List of Threatened Species (IUCN, 2017).

COMPARATIVE NOTES. *Pachydema lopadusanorum* n. sp. appears distinct morphologically from the North African populations of *P. hirticollis* as follows:

1. Body more elongated, maximum width of the elytrae towards the middle. Second segment of the antennal funicle elongated and roundish, almost as long as the 3°. Pronotum less transverse, more rounded at the sides with medium-size and spaced punctures. First elitral interstria wide, convex, clearly visible throughout the length of the elytrae. Second and third anterior tarsomeres roundish and elongated. Propygidium and pygidium of the females with bigger punctures. Aedeagus (Fig. 8) slender, the parameres apices (in lateral view) are long and pointed.....*Pachydema hirticollis*

2. Body shorter and wider, maximum width of the elytrae at the apical third. 2° segment of the antennal funicle less elongated and more roundish, shorter than the 3°. Pronotum larger and straighter on the sides, with punctures bigger and denser. 1° elitral interstria is flat, not very visible, often absent distally. Second and third anterior tarsomeres roundish. Propygidium and pygidium of the females with smaller punctures. Aedeagus (Fig. 9) shorter, the parameres apices (in lateral view) a little

elongated and a little pointed.....
.....*Pachydema lopadusanorum* n. sp.

Pachydema hirticollis is a moderately variable species morphologically. Some different taxa are considered as synonymous of *P. hirticollis* (locus typicus: Africa Dom Vahl) and they are summarized by Baraud (1985), Král & Smetana (2006), and Lacroix (2007). They, all of North African origin, are: *P. barbara* Rambur, 1843 (locus typicus: “Côte de Barbarie”), *P. nigricans* Laporte, 1832 (locus typicus: “femelle, Barbarie, Tunis”), *P. quedenfeldti* Brenske, 1889 (locus typicus: Monastir), *P. rufipennis* Burmeister, 1855 (locus typicus: “In Algerien”), *P. spreta* Fairmaire, 1860 (locus typicus: Tunis).

One syntype of *P. quedenfeldti* (from Tunisia, Monastir; type material is made of four specimens - ZMHB) shows the pronotum with more straight sides, but all the other characters correspond to




Figure 8. Aedeagus, lateral and dorsal view, of *Pachydema hirticollis*. Figure 9. Aedeagus, lateral and dorsal view, of *Pachydema lopadusanorum* n. sp.

those of other North African populations of *P. hirticollis* (Fig. 7).

Pachydemus demoflysi from Tunisia is distinct from these two species due to the smaller size, the narrower clypeus, smaller punctures of pronotum, and the shape of the aedeagus (see also Barraud, 1985 and Lacroix, 2007).

Pachydemus decorosa Normand, 1936 (locus typicus: Le Kef) is similar to *P. hirticollis* and is little known. It is cited by Král & Smetana (2006) and Lacroix (2007); for morphological differences see Barraud (1985).

REMARKS. *Pachydemus hirticollis* occurs in Algeria, Tunisia, and Libya. For Italy, Ragusa (1887, 1893, 1907) reported the capture of a pair of *P. hirticollis* in Lampedusa by Luigi Failla Tedaldi in May 1886. The occurrence of this taxon in Lampedusa was confirmed in the following bibliography (Bertolini, 1899; Luigioni, 1929; Porta, 1932; Barraud, 1977, 1986; Du Chatenet, 1986; Carpaneto & Piattella, 1995; Sparacio, 1995; Král & Smetana, 2006; Lacroix, 2007). In particular, Gridelli (1960: Cala Madonna and Monte Parrino, IV-V.1954-1956) and Arnone et al. (1995: Cala Galera, Vallone Imbriacola, Spiaggia dei Conigli) provided several localities within Lampedusa. Arnone et al. (1995), moreover, indicated *P. hirticollis* also for Pantelleria, but this population has not been studied.

All these records from *P. hirticollis* for Lampedusa island must now refer to *P. lopadusanorum* n. sp.

For these species in Tunisia and Lampedusa, see also: <http://www.entomologiitaliani.net/public/forum/phpBB3/viewtopic.php?t=29334> and <http://www.entomologiitaliani.net/public/forum/phpBB3/viewtopic.php?t=43785>.

ACKNOWLEDGEMENTS

We are grateful to Marcello Arnone (Palermo, Italy), Alberto Ballerio (Brescia, Italy), Marcello Romano (Capaci, Italy), Roberto Poggi (Museo Civico di Storia Naturale, Genoa, Italy), Giorgio Sabella and Fabio Viglianisi (Dipartimento di Biologia Animale University of Catania, Italy), Joachim Willers (Museum für Naturkunde der Humboldt Universität, Berlin, Germany), and to the staff of Legambiente (Lampedusa, Italy) for

their collaboration in the various phases of this study.

REFERENCES

- Arnone M., 2010. Quinto contributo alla revisione della collezione coleotterologica di Enrico Ragusa: Scarabaeoidea. Il Naturalista siciliano, 34: 61–172.
- Arnone M., Carpaneto G.M. & Piattella E., 1995. Arthropoda di Lampedusa, Linosa e Pantelleria. Coleoptera Scarabaeoidea. Il Naturalista siciliano, 19 (Suppl.): 447–468.
- Ballerio A., Rey A., Uliana M., Rastelli M., Rastelli S., Romano M. & Colacurcio L., 2010. Piccole Faune. Coleotteri Scarabaeoidei d'Italia. DVD. M. Serra Tarantola ed., Brescia, 1-13 + dvd pp.
- Barraud J., 1977. Faune de l'Europe occidentale: Belgique, France, Grand-Bretagne, Italie, Peninsule ibérique. IV. Coléoptères Scarabaeoidea. Nouvelle Revue d'Entomologie, Toulouse, 7 (suppl.): 1–352.
- Barraud J., 1985. Encyclopédie Entomologique XLVI. Coléoptères Scarabaeoidea. Faune du Nord de l'Afrique du Maroc au Sinaï. Édition Lechevalier, Paris, 1–651.
- Bertolini S., 1899. Catalogo dei Coleotteri d'Italia. Rivista Italiana di Scienze Naturali, Siena, 144 pp.
- Carpaneto G.M. & Piattella E., 1995. Coleoptera Polyphaga V (Lucanoidea, Scarabaeoidea). In: Minelli A., Ruffo S. & La Posta S. (Eds.), Checklist delle specie della fauna italiana. Calderini, Bologna, 50: 1–18.
- Du Chatenet G., 1986. Guide des Coléoptères d'Europe. Delachaux & Niestlé, Paris, 480 pp.
- Gridelli e., 1960. Coleoptera, 369–407 pp. In: E. Zavattari e coll., Biogeografia delle Isole Pelagie. Rendiconti Accademia Nazionale dei Lincei, 11: 1–471.
- IUCN, 2017. The IUCN Red List of Threatened Species. Version 2017–3. <http://www.iucnredlist.org> (Last access: 05.VIII.2018).
- Lacroix M., 2007. Pachydeminae du Monde (Coleoptera, Melolonthidae). Genera et Catalogue commenté. Collections Hennetons, Lacroix, Paris, 450 pp.
- Luigioni P., 1929. I Coleotteri d'Italia. Catalogo sinonimico-topografico-bibliografico. Memorie della pontificia Accademia delle Scienze Nuovi Lincei, Roma 13: 1–1160.
- Král D. & Smetana A., 2006. Pachydemini. Pp. 199–207. In: Löbl I. & Smetana A. (Eds.), Catalogue of Palaearctic Coleoptera. Vol. 3. Scarabaeoidea, Scirtidae, Dascilloidea, Buprestoidea, Byrrhoidea. Apollo Books, Stenstrup, 690 pp.
- Porta A., 1932. Fauna Coleopterorum Italica. 5. Rhynchophora-Lamellicornia. Stabilimento Tipografico Piacentino, Piacenza, 476 pp.

- Ragusa E., 1887. Coleotteri nuovi o poco conosciuti della Sicilia. *Il Naturalista siciliano*, 7: 25–28.
- Ragusa E., 1893. Catalogo ragionato dei coleotteri di Sicilia. Scarabaeidae. *Il Naturalista siciliano*, 12: 233–239.
- Ragusa E., 1907. Coleotteri nuovi o poco conosciuti della Sicilia. *Il Naturalista siciliano*, 19: 242–252.
- Sparacio I., 1995. Coleotteri di Sicilia. I. L'Epos Società Editrice, Palermo, 238 pp.