TOWER SETS AND OTHER CONFIGURATIONS WITH THE COHEN-MACAULAY PROPERTY

GIUSEPPE FAVACCHIO, ALFIO RAGUSA AND GIUSEPPE ZAPPALÀ

ABSTRACT. Some well-known arithmetically Cohen-Macaulay configurations of linear varieties in \mathbb{P}^r as k-configurations, partial intersections and star configurations are generalized by introducing tower schemes. Tower schemes are reduced schemes that are finite union of linear varieties whose support set is a suitable finite subset of \mathbb{Z}_+^c called tower set. We prove that the tower schemes are arithmetically Cohen-Macaulay and we compute their Hilbert function in terms of their support. Afterwards, since even in codimension 2 not every arithmetically Cohen-Macaulay squarefree monomial ideal is the ideal of a tower scheme, we slightly extend this notion by defining generalized tower schemes (in codimension 2) and we show that the support of these configurations (the generalized tower set) gives a combinatorial characterization of the primary decomposition of the arithmetically Cohen-Macaulay squarefree monomial ideals.

INTRODUCTION

In the last few years a large number of researchers in algebraic geometry in order to produce projective schemes with suitable Hilbert functions and graded Betti numbers constructed special configurations of linear varieties related to some subsets of \mathbb{Z}^{c}_{\perp} . Among these should be cited the partial intersection schemes introduced first in [MR] and generalized in any codimension in [RZ] and the k-configurations defined in [GS] and [GHS] to obtain maximal graded Betti numbers with respect to a fixed Hilbert function. On the other hand, to study the extremal Hilbert functions for fat point schemes in the plane, secant varieties of some classical algebraic varieties and some properties of the symbolic powers of ideals, the star configurations were defined and deeply investigated (see for instance [AS], [GHM]). All these configurations lead to aCM ideals, mostly monomial and squarefree. Looking at what all these configurations have in common, in this paper we define the tower sets (Definition 2.1), suitable finite subsets of \mathbb{Z}_{+}^{c} , on which are supported the tower schemes (Definition 2.3), which generalize all the previous mentioned configurations. These tower sets enclose the combinatorial aspects of such configurations.

Also for these schemes we are able to prove that they have the aCM property (Theorem 2.6). Moreover, we compute the Hilbert function of the tower schemes in terms of its tower set support. At this point one can believe that, at least for monomial squarefree ideals, all aCM ideals can be constructed in this way. Unfortunately, already in codimension 2,

²⁰¹⁰ Mathematics Subject Classification. 13 H 10, 14 N 20, 13 D 40.

Key words and phrases. Cohen-Macaulay, Monomial ideals, Configurations.

this is false as we show in Example 3.4. So the question which arises is to find the right configuration which could characterize all the aCM monomial squarefree ideals in a polynomial ring. Here we give a complete answer in codimension 2 (Theorems 3.19, 3.32 and 3.35) defining a slight modification of the tower schemes (generalized tower sets and schemes, see Definitions 3.12 and 3.13). The codimension bigger than 2 case remains open.

After preliminaries and basic facts, in section 2 we introduce tower sets and tower schemes and we prove that all these schemes are aCM (Theorem 2.6). Then we show that every tower scheme has the same Hilbert function as a corresponding tower scheme supported on a left segment whose Hilbert function was computed in [RZ] (see Proposition 2.11 and Corollary 2.12). Section 3 is devoted to give a combinatorial characterization for aCM squarefree monomial ideals of codimension 2. To do that we give a slight generalization of tower sets and tower schemes (Definitions 3.12 and 3.13). Then we prove numerous preparatory results about these sets and schemes and finally in Theorems 3.19 and 3.35 we prove the stated characterization.

1. NOTATION AND PRELIMINARIES

Throughout the paper k will be a field and $R := k[x_1, \ldots, x_n] = \bigoplus_d R_d$ will be the standard graded polynomial k-algebra.

We will denote by $\mathbb{Z}_+ := \{r \in \mathbb{Z} \mid r > 0\}$. If $r \in \mathbb{Z}_+$ we will set $[r] := \{1, \ldots, r\}$. If $c, r \in \mathbb{Z}_+$ we will denote by $C_{c,r}$ the set of the subsets of [r] of cardinality c.

Moreover, we will set $\pi_i : \mathbb{Z}_+^c \to \mathbb{Z}_+$ the projection on the *i*-th component. On the set \mathbb{Z}_+^c we will use the following standard partial order. If $\alpha, \beta \in \mathbb{Z}_+^c$, $\alpha \leq \beta$ iff $\pi_i(\alpha) \leq \pi_i(\beta)$ for every $i \in [c]$.

We will denote by $(\mathbb{Z}_{+}^{c})^{*} := \{(a_{1}, \ldots, a_{c}) \in \mathbb{Z}_{+}^{c} \mid a_{i} \neq a_{j} \text{ for every } i \neq j\}.$ Let $T \subset \mathbb{Z}_{+}^{c}$ be a finite set. Let $1 \leq t \leq c-1$ be an integer and let $\alpha \in \mathbb{Z}_{+}^{t}$. We set

$$T_{\alpha} := \{ \gamma \in \mathbb{Z}_{+}^{c-t} \mid (\gamma, \alpha) \in T \}$$

and

$$T^{\alpha} := \{ \gamma \in \mathbb{Z}_{+}^{c-t} \mid (\alpha, \gamma) \in T \}$$

Definition 1.1. The function $\varphi : (\mathbb{Z}_{+}^{c})^{*} \to C_{c,n}$ such that $\varphi(a_{1}, \ldots, a_{c}) = \{a_{1}, \ldots, a_{c}\}$, will be called *forgetful* function. A function $\omega : C_{c,n} \to (\mathbb{Z}_{+}^{c})^{*}$ will be called *ordinante* iff $\varphi \circ \omega = \operatorname{id}_{C_{c,n}}$.

Let $L \subset \mathbb{Z}_+^c$ be a finite set. L is said *left segment* if for every $\alpha \in L$ and $\beta \in \mathbb{Z}_+^c$ with $\beta \leq \alpha$ it follows that $\beta \in L$.

Let $L \subset \mathbb{Z}_{+}^{c}$ be a left segment. The set $\{\alpha_{1}, \ldots, \alpha_{r}\} \subseteq L$ is called *set* of generators for L if for every $\alpha \in L$, $\alpha \leq \alpha_{i}$ for some i. The element $(\max \pi_{1}(L), \ldots, \max \pi_{c}(L)) \in \mathbb{Z}_{+}^{c}$ is said the *size* of L.

Let $L \subset \mathbb{Z}_{+}^{c}$ be a left segment of size (m_1, \ldots, m_c) , with c < n. For $1 \leq i \leq c$, let $\mathcal{F}_i = \{f_{i1}, \ldots, f_{im_i}\}$ be c families of generic linear forms belonging to R. For every $\alpha = (a_1, \ldots, a_c) \in L$ we set $I_{\alpha} := (f_{1a_1}, \ldots, f_{ca_c})$. We recall that the scheme defined by the ideal $I_L(\mathcal{F}_1, \ldots, \mathcal{F}_c) := \bigcap_{\alpha \in L} I_{\alpha}$ is called *partial intersection*, with support on L and with respect to the families $\mathcal{F}_1, \ldots, \mathcal{F}_c$.

Towers sets

If $\alpha \in \mathbb{Z}_+^c$ we set $v(\alpha) := \sum_{i=1}^c \pi_i(\alpha)$. If $L \subset \mathbb{Z}_+^c$ is a left segment, the Hilbert function of L is

$$H_L(i) := |\{ \alpha \in L \mid v(\alpha) = i + c \}|.$$

We remind that H_L coincides with the Hilbert function of a partial intersection supported on L (for details about left segments and partial intersections see [RZ]).

In the sequel, if M is a matrix of rank r, with entries in R, we will denote by I(M) the ideal generated by the minors of size r in M.

2. Towers sets

Many recent papers dealt with special configurations of linear subvarieties of projective spaces which raised up to Cohen-Macaulay varieties, for instance partial intersections studied in [RZ], k-configurations studied in [GHS], star configurations studied in [GHM]. In this section we would like to generalize all these configurations in such a way to preserve the Cohen-Macaulayness.

Definition 2.1. Let $T \subset \mathbb{Z}^c_+$ be a finite set. We say that T is a *tower set* if for every $t \in [c-1]$ and for every $\alpha, \beta \in \mathbb{Z}^t_+$, with $\alpha < \beta, T_\alpha \neq \emptyset$, we have $T_\alpha \supseteq T_\beta$.

Note that when c = 1 every finite subset of \mathbb{Z}_+ is a tower set.

Remark 2.2. If $T \subset \mathbb{Z}_+^c$ is a tower set and $\alpha \in \mathbb{Z}_+^t$ then $T_\alpha \subset \mathbb{Z}_+^{c-t}$ is also a tower set.

Definition 2.3. Let $T \subset \mathbb{Z}_{+}^{c}$ be a tower set. Let $R := k[x_{1}, \ldots, x_{n}]$, with $2 \leq c \leq n-1$. Let $\mathcal{F}_{i} = \{f_{ij} \mid j \in \pi_{i}(T)\}, 1 \leq i \leq c$, where $f_{ij} \in R_{d_{ij}}$, such that f_{ij} and f_{ih} are coprime when $j \neq h$ and for every $\alpha = (a_{1}, \ldots, a_{c}) \in T$ the sequence $(f_{1a_{1}}, \ldots, f_{ca_{c}})$ is regular. We will denote by I_{α} the complete intersection ideal generated by $f_{1a_{1}}, \ldots, f_{ca_{c}}$. We set

$$I_T(\mathcal{F}_1,\ldots,\mathcal{F}_c):=\bigcap_{lpha\in T}I_lpha.$$

It defines a *c*-codimensional subscheme of \mathbb{P}^n called *tower scheme*, with support on *T*, with respect to the families $\mathcal{F}_1, \ldots, \mathcal{F}_c$.

Note that if T is a c-left segment then T is a tower set, so every partial intersection is a tower scheme.

Recently many people investigated special subschemes called star configurations. We recall that a star configuration is defined as follows. Let $R := k[x_1, \ldots, x_n], s, c \in \mathbb{Z}_+$ such that $c \leq \min\{s, n-1\}$. Take a set \mathcal{F} consisting of s forms $f_1, \ldots, f_s \in R$ such that any c of them are a regular sequence. If $s \geq a_1 > \ldots > a_c \geq 1$ are integers and $\alpha = \{a_1, \ldots, a_c\}$ we set $I_{\alpha} := (f_{a_1}, \ldots, f_{a_c})$. A star configuration is the subscheme $V_c(\mathcal{F}, \mathbb{P}^n) \subset \mathbb{P}^n$ defined by the ideal $\bigcap I_{\alpha}$ where α runs over all the subsets of [s] of cardinality c. For more details on star configurations see, for instance, [GHM].

Remark 2.4. A star configuration is a particular tower scheme. Namely, let

$$T = \{ (a_1, \dots, a_c) \in \mathbb{Z}_+^c \mid s \ge a_1 > \dots > a_c \ge 1 \},\$$

Towers sets

and let us consider the families of forms

$$\mathcal{F}_i = (f_{s-i+1}, \dots, f_{c-i+1}), \quad \text{for } 1 \le i \le c.$$

T is trivially a tower set and $V(I_T(\mathcal{F}_1,\ldots,\mathcal{F}_c)) = V_c(\mathcal{F},\mathbb{P}^n).$

In the quoted papers it was shown that partial intersections, k-configurations and star configurations are all aCM schemes. Now we prove that every tower scheme is an aCM scheme and this generalizes those results. We need the following lemma, which is a slight generalization of Lemma 1.6 in [RZ].

Lemma 2.5. Let $c, r \geq 2$ be integers. Let $V_1 \supseteq \ldots \supseteq V_r$ be (c-1)codimensional aCM subschemes of \mathbb{P}^n and $A_j = V(f_j)$ hypersurfaces, $1 \leq j \leq r$, with deg $f_j = d_j$. We set $Y_i := V_i \cap A_i$ and let us suppose that Y_i is
c-codimensional for each i and that Y_i and Y_j have no common components
for $i \neq j$. We set $d := \sum_{i=1}^r d_i$, $Y := Y_1 \cup \ldots \cup Y_{r-1}$ and $X := Y \cup Y_r$. Then
the following sequence of graded R-modules

$$0 \to I_{Y_r}(-(d-d_r)) \xrightarrow{f} I_X \xrightarrow{\varphi} I_Y/(f) \to 0$$

is exact, where $f = \prod_{i=1}^{r-1} f_i$ and φ is the natural map. Moreover

$$I_X = I_{V_1} + f_1 I_{V_2} + f_1 f_2 I_{V_3} + \dots + f_1 \dots f_{r-1} I_{V_r} + (f_1 \dots f_r).$$

Proof. The proof is analogous to that of Lemma 1.6 in [RZ]. We report it for convenience of the reader.

Observe that the exactness of the above sequence in the middle depends on the fact that f is regular modulo I_{Y_r} , since Y_i and Y_j have no common components for $i \neq j$. So, the only not trivial fact to prove is that the map φ is surjective. For this we use induction on r. For r = 2, since V_1 is aCM, $I_Y = I_{V_1} + (f_1)$, therefore every element in $I_Y/(f_1)$ looks like $z + (f_1)$ with $z \in I_{V_1} \subseteq I_{V_2}$. Hence, $z \in I_{Y_1} \cap I_{Y_2} = I_X$. So, φ is surjective and the sequence is exact. Now, from the exactness of this sequence it follows that I_X is generated by I_{V_1} and $f_1I_{Y_2}$, i.e. $I_X = I_{V_1} + f_1I_{V_2} + (f_1f_2)$.

Let us suppose the lemma true for r-1. This means, in particular, that $I_Y = I_{V_1} + f_1 I_{V_2} + \dots + f_1 \dots f_{r-2} I_{V_{r-1}} + (f_1 \dots f_{r-1})$. Therefore, every element $z \in I_Y/(f_1 \dots f_{r-1})$ has the form $x + (f_1 \dots f_{r-1})$ with $x \in I_{V_1} + f_1 I_{V_2} + \dots + f_1 \dots f_{r-2} I_{V_{r-1}}$. Hence, $x \in I_{V_r}$ which implies $x \in I_Y \cap I_{Y_r} = I_X$. Again, by the exactness of our sequence we get that I_X is generated by $f_1 \dots f_{r-1} I_{Y_r}$ and by $I_{V_1} + f_1 I_{V_2} + \dots + f_1 \dots f_{r-2} I_{V_{r-1}}$, i.e. $I_X = I_{V_1} + f_1 I_{V_2} + \dots + f_1 \dots f_{r-1} I_{V_r} + (f_1 \dots f_r)$.

We are ready to prove our result.

Theorem 2.6. Every tower scheme is aCM.

Proof. Let X be a tower scheme of codimension c. To show that X is aCM we use induction on c. The property is trivially true for c = 1, so we can assume that every tower scheme of codimension c-1 is aCM. Let T be the support of X and let $\mathcal{F}_i = \{f_{ij} \mid j \in \pi_i(T)\}$, for $1 \leq i \leq c$, be the families defining X. Let $\pi_c(T) = \{a_1, \ldots, a_s\}$, with $a_1 < \ldots < a_s$. For every $i \in [s]$ we denote by V_i the tower scheme of codimension c-1 supported on T_{a_i} . If i < j then $V_i \supseteq V_j$. By inductive hypotheses each V_i is aCM. Moreover we

Towers sets

denote by A_i the hypersurface defined by f_{ca_i} , with $i \in [s]$ and $Y_i = V_i \cap A_i$. Note that by the hypotheses on \mathcal{F}_h 's Y_i is aCM of codimension c.

Therefore we have $X = \bigcup_{1 \le i \le s} Y_i$. Now we use induction on s. For s = 1 $X = Y_1$ is aCM. Suppose that $Y = \bigcup_{1 \le i \le s-1} Y_i$ is aCM and show that $X = Y \cup Y_s$ is aCM. Applying the previous lemma we get the exact sequence

 $0 \to I_{Y_s}(-\deg f) \to I_X \to I_Y/(f) \to 0$

where $f = \prod_{i=1}^{s-1} f_{ca_i}$ from which we see that a resolution of I_X can be obtained as direct sum of the resolutions of $I_{Y_s}(-\deg f)$ and $I_Y/(f)$; since both have resolutions of length c the same is true for I_X and we are done.

Our next aim is to compute the Hilbert function of a tower scheme in the case when the defining families consist of linear forms. To do this, if T is a tower set, we define a map $\sigma : T \to \mathbb{Z}_+^c$ as follows. Let $\alpha = (a_1, \ldots, a_c) \in T$, we set

$$h_1(\alpha) := |\{i \mid i \le a_1, (i, a_2, \dots, a_c) \in T\}|,$$

$$h_j(\alpha) := |\{i \mid i \le a_j, T_{(i, a_{j+1}, \dots, a_c)} \neq \emptyset\}|, \text{ for } 2 \le j \le c$$

and finally

$$\sigma(a_1,\ldots,a_c):=(h_1(\alpha),\ldots,h_c(\alpha)).$$

The map σ is trivially injective. We set $T^{\#} := \sigma(T)$.

Proposition 2.7. For every tower set $T, T^{\#}$ is a left segment.

Proof. Let $\alpha' = (a'_1, \ldots, a'_c) \in T^{\#}$ and $\beta' = (b'_1, \ldots, b'_c) \in \mathbb{Z}^c_+$, such that $\beta' \leq \alpha'$. We have to prove that $\beta' \in T^{\#}$, i.e. we have to find $\beta \in T$ such that $\sigma(\beta) = \beta'$. Let $\alpha = (a_1, \ldots, a_c) \in T$ be such that $\sigma(\alpha) = \alpha'$, hence $a'_i = h_i(\alpha)$. Since $b'_c \leq h_c(\alpha)$, there is a unique element b_c such that $T_{b_c} \neq \emptyset$ and $|\{i \mid i \leq b_c, T_i \neq \emptyset\}| = b'_c$. Now, since $b'_{c-1} \leq h_{c-1}(\alpha)$, we have that $T_{(b'_{c-1}, a_c)} \neq \emptyset$ and, since T is a tower set, $T_{(b'_{c-1}, b_c)} \neq \emptyset$, therefore there is a unique element b_{c-1} such that $T_{(b_{c-1}, b_c)} \neq \emptyset$ and $|\{i \mid i \leq b_{c-1}, T_{(i, b_c)} \neq \emptyset\}| = b'_{c-1}$. By iterating the same argument we will set b_j the unique element such that $T_{(b_j, b_{j+1}, \ldots, b_c)} \neq \emptyset$ and $|\{i \mid i \leq b_j, T_{(i, b_{j+1}, \ldots, b_c)} \neq \emptyset\}| = b'_j$, for $1 \leq j \leq c$. Now we set $\beta = (b_1, \ldots, b_c)$. By definition $\beta \in T$ and $\sigma(\beta) = \beta'$.

Remark 2.8. Note that if $T, U \subset \mathbb{Z}^c_+$ are tower sets such that $T \subseteq U$ then $T^{\#} \subseteq U^{\#}$.

Proposition 2.9. Let $T \subset \mathbb{Z}_+^c$ be a tower set. Let $X = V(I_T(\mathcal{F}_1, \ldots, \mathcal{F}_c))$. Let Y be a tower scheme supported on $T^{\#}$, with respect to the same families $\mathcal{F}_1, \ldots, \mathcal{F}_c$. Then $H_X = H_Y$.

Proof. If c = 1 then T is a finite subset of \mathbb{Z} , say r = |T|, so $T^{\#} = [r]$. Therefore I_X and I_Y are principal ideals generated by a form of same degree, hence $H_X = H_Y$.

So we may assume that $c \geq 2$ and we proceed by induction on c. We consider the set $\pi_c(T) = \{m_1, \ldots, m_s\}, m_1 < \ldots < m_s$. Since T is a tower set, $T_{m_1} \supseteq \ldots \supseteq T_{m_s}$ are (c-1)-tower sets. Let X_i be the scheme defined by

 $I_{T_{m_i}}(\mathcal{F}_1,\ldots,\mathcal{F}_{c-1})$. Then each X_i is an aCM scheme of codimension c-1 by Theorem 2.6 and $X_1 \supseteq \ldots \supseteq X_s$. Moreover, by Remark 2.8, $(T_{m_1})^{\#} \supseteq \ldots \supseteq (T_{m_s})^{\#}$. Let Y_i be the scheme defined by $I_{(T_{m_i})^{\#}}(\mathcal{F}_1,\ldots,\mathcal{F}_{c-1})$. By the inductive hypothesis $H_{X_i} = H_{Y_i}$. Now, let $\mathcal{F}_c = \{f_1,\ldots,f_s\}$, we set $\overline{X_i} := X_i \cap V(f_i)$ and $\overline{Y_i} = Y_i \cap V(f_i)$. Since X_i and Y_i are aCM then $H_{\overline{X_i}} = H_{\overline{Y_i}}$. Finally, using induction on s and the exact sequences (see Lemma 2.5)

$$0 \to I_{\overline{X}_s}(-\delta) \to I_X \to I_{\overline{X}_i \cup \dots \cup \overline{X}_{s-1}}/(f_1 \dots f_{s-1}) \to 0$$
$$0 \to I_{\overline{Y}_s}(-\delta) \to I_Y \to I_{\overline{Y}_i \cup \dots \cup \overline{Y}_{s-1}}/(f_1 \dots f_{s-1}) \to 0$$
$$\log(f_{x_s}(-\delta)) \quad \text{we get the conclusion}$$

where $\delta = \deg(f_1 \dots f_{s-1})$, we get the conclusion.

Proposition 2.9 allows us to find a formula for the Hilbert function of a tower scheme.

Remark 2.10. Note that, according to the exact sequence of Lemma 2.5, the Hilbert function of a tower scheme depends on the tower set and on the degrees of the forms in the families.

Now we associate to a tower scheme X, supported on a left segment L, a partial intersection Y with support on a suitable left segment L_D such that $H_X = H_Y$.

If L is a left segment of size (a_1, \ldots, a_c) and $D = \{d_{ij}\}, 1 \leq i \leq c$ and $1 \leq j \leq a_i$ are positive integers, we define a new left segment, which we will be denoted by L_D , in the following way. If L is (minimally) generated by K_1, \ldots, K_r , then L_D is the left segment generated by K'_1, \ldots, K'_r where, if $K_i = (k_1, \ldots, k_c)$ then $K'_i = (\sum_{j=1}^{k_1} d_{1j}, \ldots, \sum_{j=1}^{k_c} d_{cj})$. Thus, let X be a tower scheme supported on the left segment L and let

Thus, let X be a tower scheme supported on the left segment L and let $\mathcal{F}_i = \{f_{ij} \mid j \in \pi_i(L)\}$, for $1 \leq i \leq c$, be the families defining X. Set now $\pi_c(L) = \{a_1, \ldots, a_c\}$, with $a_1 < \ldots < a_c$ and $D = \{d_{ij}\}$ where $d_{ij} = \deg f_{ij}$, $1 \leq i \leq c$ and $1 \leq j \leq a_i$. Since, by Remark 2.10, H_X depends just on D we may assume that $f_{ij} = \prod_{h=1}^{d_{ij}} l_{ij}^h$, where each l_{ij}^h is a linear form. Now we denote by Y the partial intersection supported on L_D with respect the c ordered families of linear forms

$$\mathcal{L}_{i} = (l_{i11}, \dots, l_{i1d_{i1}}, l_{i21}, \dots, l_{i2d_{i2}}, \dots, l_{ia_{i1}}, \dots, l_{ia_{id_{ia_i}}}).$$

Proposition 2.11. Given a tower scheme X supported on the left segment L with respect to the families of forms $\mathcal{F}_i = \{f_{ij} \mid j \in \pi_i(L)\}$, for $1 \leq i \leq c$ and $D = \{d_{ij}\}$ where $d_{ij} = \deg f_{ij}$. Then $H_X = H_{L_D}$.

Proof. By definition $I_X = \bigcap_{(j_1,\ldots,j_c)\in L}(f_{1j_1},\ldots,f_{cj_c})$. Now we denote by Y the partial intersection supported on L_D with respect the c ordered families of linear forms $\mathcal{L}_i = (l_{i11},\ldots,l_{i1d_{i1}},l_{i21},\ldots,l_{i2d_{i2}},\ldots,l_{ia_i1},\ldots,l_{ia_id_{ia_i}}), 1 \leq i \leq c$. Now if α is an integer such that $1 \leq \alpha \leq \sum_{s=1}^{a_i} d_{is}$ we set

$$t_{\alpha} := \max\{j \mid d_{i1} + \ldots + d_{ij} < \alpha\} + 1$$

and

$$h_{\alpha} := \alpha - \sum_{s=1}^{t_{\alpha}-1} d_{is}$$

and $p_{i\alpha} := l_{it_{\alpha}h_{\alpha}}$. Then, with this notation

$$I_Y = \bigcap_{(\alpha_1,\dots,\alpha_c)\in L_D} (p_{1\alpha_1},\dots,p_{c\alpha_c}).$$

It is a matter of computation to show that $I_X = I_Y$ and this completes the proof.

In the next corollary we lead back the computation of the Hilbert function of a tower scheme to that of a partial intersection. The Hilbert function of a partial intersection was explicitly computed in [RZ].

Corollary 2.12. If X is a tower scheme supported on a tower set T with respect to families of forms of degrees D, then $H_X = H_{(T^{\#})_D}$.

Proof. It follows just using Propositions 2.9 and 2.11.

3. Generalized tower sets: a characterization of ACM property

In this section we will generalize tower sets in such a way to characterize aCM squarefree monomial ideal of codimension 2.

Let $I \subset k[x_1, \ldots, x_n]$ be an equidimensional squarefree monomial ideal of codimension c and let $I = \mathfrak{p}_1 \cap \ldots \cap \mathfrak{p}_t$ be its primary decomposition. Each \mathfrak{p}_i is a prime ideal of the type $(x_{a_{i1}}, \ldots, x_{a_{ic}})$. So we can consider the subset $S(I) := \{\{a_{i1}, \ldots, a_{ic}\} \mid 1 \leq i \leq t\}$ of $C_{c,n}$.

Vice versa to $\mathcal{S} \subseteq C_{c,n}$ we can associate an equidimensional squarefree monomial ideal

$$I_{\mathcal{S}} := \bigcap_{\{a_1,\dots,a_c\}\in\mathcal{S}} (x_{a_1},\dots,x_{a_c}).$$

Definition 3.1. Let $S \subseteq C_{c,n}$. We will say S aCM if I_S is an aCM ideal.

Definition 3.2. Let $S \subseteq C_{c,n}$. We will say that S is towerizable if there exists a permutation τ on [n] and an ordinante function $\omega : C_{c,n} \to (\mathbb{Z}_+^c)^*$ such that $\tau(\omega(S))$ is a tower set.

Remark 3.3. Let $S \subseteq C_{c,n}$. Note that S is towerizable if there exists a tower set T and families $\mathcal{F}_i \subseteq \{x_1, \ldots, x_n\}$, such that $\mathcal{S}(I_T(\mathcal{F}_1, \ldots, \mathcal{F}_c)) = S$.

By Theorem 2.6 if S is towerizable then S is aCM, however there are aCM equidimensional squarefree monomial ideals I such that S(I) is not towerizable. Here it is an example in codimension 2.

Example 3.4. Let $S = \{\{1,2\},\{3,4\},\{5,6\},\{4,6\},\{1,4\},\{1,6\}\}$. Then it is easy to check that I_S is the determinantal ideal generated by the order 3 minors of the matrix

$$\begin{pmatrix} x_1 & 0 & 0 \\ x_2 & x_3 & x_5 \\ 0 & x_4 & 0 \\ 0 & 0 & x_6 \end{pmatrix}$$

so S is aCM. Let us suppose that S is towerizable. Then there exists a tower scheme X with support on a tower set T such that $S(I_X) = S$. Of course |T| = 6 and there is not a variable x_k such that the ideal (x_k) contains 4 of the 6 minimal primes of I_S . Consequently, $|\pi_2(T)| \leq 3$ and for every $a \in \pi_2(T) |T_a| \leq 3$ so we have only three possibilities

- 1) $\pi_2(T) = \{a, b\}$ with $|T_a| = 3$ and $|T_b| = 3$;
- 2) $\pi_2(T) = \{a, b, c\}$ with $|T_a| = 2$, $|T_b| = 2$ and $|T_c| = 2$;
- 3) $\pi_2(T) = \{a, b, c\}$ with $|T_a| = 3$, $|T_b| = 2$ and $|T_c| = 1$.

The first two cases cannot occur since $I_X = I_S$ does not contain monomials of degree two.

Therefore $T_a \supset T_b \supset T_c$ and $T_a = \{h_1, h_2, h_3\}, T_b = \{h_1, h_2\}, T_c = \{h_1\}$ for some h_i 's and thus $T = \{(h_1, a), (h_2, a), (h_3, a), (h_1, b), (h_2, b), (h_1, c)\}$. But the numbers 2, 3 and 5 belong each to one only element of S whereas in T there are only two such numbers, precisely h_3 and c.

Because of the previous example it is natural to ask which sets $S \subseteq C_{c,n}$ are aCM. We will give a characterization in codimension 2 (see Theorems 3.19 and 3.35).

Definition 3.5. Let $h \in \mathbb{Z}_+$. Let $S \subseteq C_{2,n}$. We set

$$\mathcal{S}: h := \{ A \in \mathcal{S} \mid h \notin A \}.$$

If $S \subseteq \mathbb{Z}^2_+$ we set

$$S: h := \{ \alpha \in S \mid \pi_1(\alpha) \neq h \text{ and } \pi_2(\alpha) \neq h \}.$$

Remark 3.6. Note that if S is aCM then S : h is aCM. Indeed, if M is an Hilbert-Burch matrix for I_S then $I_{S:h}$ is generated by the maximal minors of the matrix obtained from M by replacing x_h with 1.

In the sequel we will use the following result which shows that if $S \subseteq C_{2,n}$ is aCM then also the scheme obtained by replacing $(x_i, x_j) \supseteq I_S$ with (h_i, h_j) , generic complete intersections, is aCM.

Proposition 3.7. Let $S \subseteq C_{2,n}$ be an aCM set, with $I_S \subset k[x_1, \ldots, x_n]$. Let $h_1, \ldots, h_n \in k[y_1, \ldots, y_m]$ be forms, such that depth $(h_i, h_j) = 2$ for every $\{i, j\} \in S$ and

$$depth(h_i, h_j, h_u, h_v) \ge 3$$

for every $\{i, j\}, \{u, v\} \in S, \{i, j\} \neq \{u, v\}$. Then the ideal $J = \bigcap_{\{i, j\} \in S} (h_i, h_j)$ is aCM.

Proof. We consider the following vectors

$$\underline{x} = (x_1, ..., x_n) \text{ and } \underline{h} = (h_1, ..., h_n).$$

Since $I_{\mathcal{S}}$ is aCM we can consider $M = M(\underline{x})$, an Hilbert-Burch matrix for $I_{\mathcal{S}}$. We claim that $N = M(\underline{h})$ is an Hilbert-Burch matrix for J. We have to prove that J = I(N). Let $g \in I(N)$ be a maximal minor of N. Then $g = f(\underline{h})$, with $f(\underline{x}) \in I_{\mathcal{S}}$. Therefore $f(\underline{x}) = \lambda_i(\underline{x})x_i + \mu_j(\underline{x})x_j$ for every $\{i, j\} \in S$, consequently $f(\underline{h}) \in (h_i, h_j)$ for every $\{i, j\} \in S$. So $I(N) \subseteq J$.

To conclude the proof it is enough to show that $\deg I(N) = \deg J$. By the generality of the forms h_1, \ldots, h_n , we have that

$$\deg J = \sum_{\{i,j\}\in\mathcal{S}} (\deg h_i) (\deg h_j).$$

Now we proceed by induction on n. If n = 2 then $S = \{1, 2\}$ and $I(N) = (h_1, h_2)$. So we can suppose that deg $I(N) = \sum_{\{i,j\}\in \mathcal{S}} (\deg h_i)(\deg h_j)$, when $S \subseteq C_{2,n-1}$. We can write $S = (S : n) \cup S_{(n)}$ where $S : n = \{\alpha \in S \mid n \notin \alpha\}$

and $\mathcal{S}_{(n)} = \mathcal{S} \setminus (\mathcal{S} : n)$. By Remark 3.6, $\mathcal{S} : n$ is aCM. Let $\widehat{M}(\underline{x})$ be an Hilbert-Burch matrix for $I_{\mathcal{S};n}$. We set $\widehat{N} = \widehat{M}(\underline{h})$. Therefore

$$I_{\mathcal{S}} = I(M) = (I(M) : x_n) \cap (x_n, \prod_{u \in \mathcal{S}_{(n)}} x_u) = I(\widehat{M}) \cap (x_n, \prod_{u \in \mathcal{S}_{(n)}} x_u).$$

Hence, using the inductive hypothesis, we get

$$\deg I(N) = \deg(I(\widehat{N})) + \deg(h_n, \prod_{u \in \mathcal{S}_{(n)}} h_u)) =$$

$$\sum_{\{i,j\}\in(\mathcal{S}:n)} (\deg h_i)(\deg h_j) + (\deg h_n) \sum_{\{u,n\}\in\mathcal{S}_{(n)}} \deg h_u = \sum_{\{i,j\}\in\mathcal{S}} (\deg h_i)(\deg h_j).$$

We recall that if $T \subseteq \mathbb{Z}^2_+$ and $i \in \mathbb{Z}_+$ then

$$T^{i} = \{ j \in \pi_{2}(T) \mid (i, j) \in T \}$$

and

$$T_i = \{ j \in \pi_1(T) \mid (j, i) \in T \}.$$

Remark 3.8. Let $T \subset (\mathbb{Z}^2_+)^*$ be a tower set. Then

- 1) a < b and $T_a \neq \emptyset \Rightarrow (a, b) \notin T$. Indeed, the assumption implies $T_a \supseteq T_b$. Since $(a, a) \notin T$, we have $a \notin T_a$, therefore $a \notin T_b$ i.e. $(a, b) \notin T$.
- 2) a < b and $(b, a) \in T \Rightarrow (a, b) \notin T$. Indeed, the assumption implies $T_a \neq \emptyset$ so, by the previous item, $(a, b) \notin T$.
- 3) $\{(a,b),(b,a)\} \not\subseteq T$ for every a and b. It follows by item 2.

Note that by item 3, $|T| = |\varphi(T)|$ where φ is the forgetful function.

Proposition 3.9. Let $T \subset \mathbb{Z}^2_+$ be a tower set. Then T^i and T^h are comparable under inclusion for every i and h.

Proof. Let $j \in T^i$ be such that $j \notin T^h$, we have to show that $T^h \subset T^i$. Let $k \in T^h$, i.e. $h \in T_k$; but $h \notin T_j$ therefore since T is a tower set we have that $T_j \subset T_k$, so $i \in T_k$ i.e. $(i,k) \in T$ that implies that $k \in T^i$.

Proposition 3.10. Let $T \subset (\mathbb{Z}^2_+)^*$ be a tower set.

- 1) Let $h \in \pi_2(T)$ be such that $T_h \supseteq T_j$ for every $j \in \pi_2(T)$. Then $h \notin \pi_1(T)$.
- 2) Let $h \in \pi_1(T)$ be such that $T^h \supseteq T^i$ for every $i \in \pi_1(T)$. Then $h \notin \pi_2(T)$.

Proof. 1) If $(h, j) \in T$ then $h \in T_j \subseteq T_h$, i.e. $(h, h) \in T$.

2) Using Proposition 3.9 the proof is analogous to item 1.

Let $T \subset (\mathbb{Z}^2_+)^*$ be a tower set. Let $h \in \pi_1(T) \cap \pi_2(T)$. We set

$$F_T(h) := \{ j \in \pi_2(T) \mid T_h \subset T_j \text{ and } (h, j) \notin T \}.$$

Note that if $j \in F_T(h)$ then j < h.

Definition 3.11. Let $U \subseteq C_{2,n}$. We say that U is *connected* if for every $A, B \in U$ there is $C \in U$ such that $A \cap C \neq \emptyset$ and $B \cap C \neq \emptyset$. Let $S \subset (\mathbb{Z}^2_+)^*$. We say that S is *connected* if $\varphi(S)$ is connected.

Definition 3.12. Let $S \subset (\mathbb{Z}^2_+)^*$ be a finite set. We say that S is a generalized tower set if

- 1) S is connected;
- 2) $S = T \cup S_0$ where T is a tower set

and S_0 has the following further properties

- 3) for every $(i, j) \in S_0, i \notin \pi_1(T) \cup \pi_2(T)$ and $j \in \pi_1(T) \cap \pi_2(T)$;
- 4) for every $(i, j) \in S_0$ and $h \in F_T(j)$, $(i, h) \in S_0$.

Definition 3.13. Let $S \subset (\mathbb{Z}_{+}^{2})^{*}$ be a generalized tower set. Let $R = k[x_{1}, \ldots, x_{n}], n \geq 3$. Let $\mathcal{F}_{i} = \{f_{ij} \mid j \in \pi_{i}(S)\}, 1 \leq i \leq 2$, where each f_{ij} is a form satisfying such conditions of genericity: for every $(a_{1}, a_{2}) \in S, f_{1a_{1}}, f_{2a_{2}}$ are coprime and for every $(a_{1}, a_{2}), (b_{1}, b_{2}) \in S$, with $\{a_{1}, a_{2}\} \neq \{b_{1}, b_{2}\}, depth(f_{1a_{1}}, f_{2a_{2}}, f_{1b_{1}}, f_{2b_{2}}) \geq 3$. If $\alpha = (a_{1}, a_{2}) \in S$, we will denote by I_{α} the complete intersection ideal generated by $f_{1a_{1}}, f_{2a_{2}}$. We set

$$I_S(\mathcal{F}_1, \mathcal{F}_2) := \bigcap_{\alpha \in S} I_{\alpha}.$$

It defines a 2-codimensional subscheme of \mathbb{P}^n called *generalized tower scheme*, with support on S, with respect to the families $\mathcal{F}_1, \mathcal{F}_2$.

In the sequel if $S \subset (\mathbb{Z}^2_+)^*$ we will set for short $I_S := I_{\varphi(S)}$, consequently S will be said aCM if I_S is aCM.

In order to prove our results on the Cohen-Macaulayness of such schemes we need several lemmas.

Lemma 3.14. Let $S = T \cup S_0$ be a generalized tower set. Let $i \in \pi_1(S_0)$ and let $m = \min\{j \mid (i, j) \in S_0\}$. Then $F_T(m) = \emptyset$.

Proof. Let $s \in F_T(m)$; then s < m and by Definition 3.12, item 4, $(i, s) \in S_0$, which is a contradiction.

Lemma 3.15. With the above notation, if $h \in \pi_1(T) \cap \pi_2(T)$ then for every $j \in \pi_1(T) \cap \pi_2(T) \setminus \{h\}$ we have $F_{T:h}(j) \subseteq F_T(j)$.

Proof. If $b \in F_{T:h}(j)$ then $(T:h)_j \subset (T:h)_b$ and $(j,b) \notin T:h$, with $j \neq h$ and $b \neq h$, so $(j,b) \notin T$. Moreover there is a such that $(a,b) \in T:h$ and $(a,j) \notin T:h$. Since $a \neq h$ this implies that $T_b \not\subseteq T_j$. Since T is a tower set we get that $T_j \subset T_b$.

Lemma 3.16. Let $S = T \cup S_0$ be a generalized tower set. Let $h \in \pi_1(T) \cap \pi_2(T)$. Then S : h is a generalized tower set with respect to the decomposition $S : h = (T : h) \cup (S_0 : h)$.

Proof. Of course $S : h = (T : h) \cup (S_0 : h)$.

- 1) Since S is connected then S:h is connected too.
- 2) Let $a, b \in \pi_2(T:h)$, a < b. Let $i \in (T:h)_b$; then $(i,b) \in T:h$ i.e. $i \in T_b \subseteq T_a$; since $i \neq h$ and $b \neq h$ then $i \in (T:h)_a$.

- 3) Let $(i, j) \in S_0$: h. Of course $i \notin \pi_1(T:h) \cup \pi_2(T:h)$. Moreover $(\pi_1(T) \cap \pi_2(T)) \setminus \{h\} = \pi_1(T:h) \cap \pi_2(T:h)$. Indeed, $\pi_1(T:h) \cap \pi_2(T:h) \subseteq (\pi_1(T) \cap \pi_2(T)) \setminus \{h\}$ trivially; if $k \in (\pi_1(T) \cap \pi_2(T)) \setminus \{h\}$ then $k \neq h, k \in T_b$ where $T_b \supseteq T_u$ for every u and $k \in T^a$ where $T^a \supseteq T^v$ for every v, (see Proposition 3.9), so by Proposition 3.10, $a \neq h$ and $b \neq h$ i.e. $(a, k), (k, b) \in T: h$, hence item 3 is clear.
- 4) Let $(i, j) \in S_0$: h and $u \in F_{T:h}(j)$. By Lemma 3.15, $u \in F_T(j)$, so $(i, u) \in S_0$. Since $i \neq h$ and $u \neq h$ we get $(i, u) \in S_0$: h.

Lemma 3.17. Let $U \subseteq C_{2,n}$ and $a \in [n]$. Then $I_{U:a} = I_U : (x_a)$.

Proof. $I_U = \bigcap_{\{i,j\} \in U} (x_i, x_j)$. Then

$$I_U: (x_a) = \bigcap_{\{i,j\} \in U} \left((x_i, x_j) : (x_a) \right) = \bigcap_{\{i,j\} \in U:a} (x_i, x_j) = I_{U:a}.$$

Lemma 3.18. Let $S = T \cup S_0$ be a generalized tower set. For every $a \in \pi_1(S_0)$, there is $(a, h) \in S_0$ such that $I_{S:a} + (x_h)$ is a complete intersection ideal of height 2.

Proof. We proceed step by step.

1) In this first step we show that the assertion is equivalent to prove that for all $\{i, j\} \in \varphi(S : a)$ we have either $\{i, h\} \in \varphi(S : a)$ or $\{j, h\} \in \varphi(S : a)$.

At the beginning we observe that if $I_{S:a} + (x_h)$ is equidimensional of height 2 and \mathfrak{p} is a minimal prime in its primary decomposition then $x_h \in \mathfrak{p}$, so $I_{S:a} + (x_h) = \bigcap_i (x_h, x_i) = (x_h, \prod_i x_i)$ that is a complete intersection.

On the other hand to show that $I_{S:a} + (x_h)$ is equidimensional of height 2 it is enough to prove that for all $\{i, j\} \in \varphi(S : a)$ we have either $\{i, h\} \in \varphi(S : a)$ or $\{j, h\} \in \varphi(S : a)$. In fact let $\mathfrak{p} =$ (x_i, x_j, x_h) be a prime ideal containing $I_{S:a} + (x_h)$, so $\{i, j\} \in \varphi(S :$ a), consequently $\{i, h\} \in \varphi(S : a)$ or $\{j, h\} \in \varphi(S : a)$ i.e. \mathfrak{p} contains a prime ideal of height 2 containing $I_{S:a} + (x_h)$.

Now let $a \in \pi_1(S_0)$, and let $m_1 = \min S^a$; we set

$$U(S^a) := \{ m \in S^a \mid T_m = T_{m_1} \text{ and } F_T(m) = \emptyset \}.$$

Note that by Lemma 3.14, $m_1 \in U(S^a)$.

We claim that the integer h which we are looking for can be found in $U(S^a)$. In the next two steps we prove properties of $U(S^a)$ for our claim.

- 2) Let $m, n \in U(S^a)$ then $T^m = T^n$. If $\alpha \in T^m \setminus T^n$, i.e. $(m, \alpha) \in T$ and $(n, \alpha) \notin T$ we have $T_\alpha \supset T_m = T_n$. Now since $\alpha \notin F_T(n) = \emptyset$ and $(n, \alpha) \notin T$ we should have $T_\alpha \subseteq T_n$, a contradiction.
- 3) If $F_T(m) = \emptyset$ and $(\alpha, \beta) \in T$ then either $(\alpha, m) \in T$ or $(m, \beta) \in T$. Since $\beta \notin F_T(m) = \emptyset$ we get either $T_\beta \subseteq T_m$, hence $(\alpha, m) \in T$, or $(m, \beta) \in T$.

In the remaining steps we will find the integer h working by induction on $|U(S^a)|$.

- 4) If $|U(S^a)| = 1$ then $U(S^a) = \{m_1\}$, and we would like to show that for all $\{i, j\} \in \varphi(S : a)$ we have either $\{i, m_1\} \in \varphi(S : a)$ or $\{j, m_1\} \in \varphi(S : a)$. If $\{i, j\} \in \varphi(T)$ then by item 3 we are done. So we can assume that $\{i, j\} \in \varphi(S_0)$, with $(i, j) \in S_0$. Of course we can suppose that $j \neq m_1$. Since $j \notin F_T(m_1)$ then either $(m_1, j) \in T$ or $T_j \subseteq T_{m_1}$. If $(m_1, j) \in T$ we are done; otherwise we can assume that $T_j \subseteq T_{m_1}$.
- if $T_j \subset T_{m_1}$ then either $(j, m_1) \in T$ and we are done, or $(j, m_1) \notin T$ then $m_1 \in F_T(j)$, so by item 4 in Definition 3.12 we get $(i, m_1) \in S_0$ and we are done again.
- if $T_{m_1} = T_j$, let us suppose that

$$\{i, m_1\}, \{m_1, j\} \notin \varphi(S).$$

Since S is connected then, taking (i, j) and (a, m_1) , we get $(a, j) \in S_0$. Since $U(S^a) = \{m_1\}$ and $j \neq m_1$ we have $F_T(j) \neq \emptyset$. Now take $k \in F_T(j)$; then $T_{m_1} = T_j \subset T_k$, so $k < m_1$. Since $(a, j) \in S_0$ and $k \in F_T(j)$ by item 4 in Definition 3.12, we have $(a, k) \in S_0$, and this contradicts the minimality of m_1 .

Let now $U(S^a) = \{m_1, \dots, m_p\}, p > 1.$

- 5) At first we prove that $U(S^a) \setminus \{m_p\} = U((S:m_p)^a)$. The inclusion $U(S^a) \setminus \{m_p\} \subseteq U((S:m_p)^a)$ follows directly by definition of $U(S^a)$ and by Lemma 3.15. Let $m \in U((S:m_p)^a)$ then $m \neq m_p$ and $(T:m_p)_m = (T:m_p)_{m_1}$; since $T_{m_1} = T_{m_p}$ we get $m_p \notin T_{m_1}$ therefore $T_m \supseteq T_{m_1}$. By the minimality of m_1 we have $T_m = T_{m_1}$. Now if $b \in F_T(m)$ then $T_{m_1} = T_m \subset T_b$, so $b < m_1$. Since S is a generalized tower set we get $(a,b) \in S_0$ and this contradicts the minimality of m_1 .
- 6) By the inductive hypothesis there exists $m \in U(S^a) \setminus \{m_p\}$ such that for any $(i, j) \in (S : a) : m_p$ we have either

$$\{m, i\}$$
 or $\{m, j\} \in \varphi((S : a) : m_p)$.

We will prove that either m_p or m is the wanted element. Let us suppose that there exist $(\alpha, \beta), (u, v) \in S$ such that

$$\{m, \alpha\}, \{m, \beta\} \notin \varphi((S:a)) \text{ and } \{m_p, u\}, \{m_p, v\} \notin \varphi((S:a)).$$

Note that $\beta = m_p$, since otherwise $(\alpha, \beta) \in S : m_p$. Now since $v \neq m_p$ $(u, v) \in S_0 : m_p$, hence by hypothesis on m it should be either $(u, m) \in S_0$ or $\{v, m\} \in \varphi(T)$; but the last assertion is false since, by item 2, $T_m = T_{m_p}$ and $T^m = T^{m_p}$. This implies that (α, m_p) and $(u, m) \in S$ and this contradicts the connection of S.

Finally we are ready to prove the announced result.

Theorem 3.19. If S is a generalized tower set then S is aCM.

Proof. $S = T \cup S_0$, where T is a tower set and for S_0 the properties of Definition 3.12 hold. We proceed by induction on $r = |\pi_1(S_0)|$. If $S_0 = \emptyset$ then S = T which is aCM by Theorem 2.6. Now we can suppose that the assertion is true up to r - 1. Take $a \in \pi_1(S_0)$. Note that $S : a = T \cup (S_0 : a)$

and S: a is a generalized tower set with respect to this decomposition. Then by inductive hypothesis S: a is aCM. We can write $I_S = I_{S:a} \cap (x_a, f)$ where $f = \prod_{i \in S^a} x_i$. Using the exact sequence

$$0 \to I_S \to I_{S:a} \oplus (x_a, f) \to I_{S:a} + (x_a, f) \to 0$$

it is enough to show that proj-dim $(I_{S:a} + (x_a, f)) \leq 3$, i.e. proj-dim $(I_{S:a} + (f)) \leq 2$. To do that we use induction on deg $f = |S^a|$. If $f = x_h$ then, by Lemma 3.18, $I_{S:a} + (x_h)$ is a complete intersection ideal of height 2 and we are done.

If deg $f = |S^a| > 1$, by Lemma 3.18, there is $h \in S^a$ such that $I_{S:a} + (x_h)$ is a complete intersection ideal of height 2. We can write

$$I_{S:a} + (f) = ((I_{S:a} : (x_h)) + (f_h)) \cap (I_{S:a} + (x_h)),$$

where $f_h = f/x_h$. In fact $I_{S:a} + (f) \subseteq ((I_{S:a} : (x_h)) + (f_h)) \cap (I_{S:a} + (x_h))$ trivially. Let $g \in ((I_{S:a} : (x_h)) + (f_h)) \cap (I_{S:a} + (x_h))$ be a monomial. If $g \in I_{S:a}$ we are done, otherwise $g \in (x_h)$. If $g \in (f_h)$ then $g \in (f)$ and we are done again. Otherwise $x_h g \in I_{S:a}$; since $I_{S:a}$ is a monomial squarefree ideal, we get $g \in I_{S:a}$.

By Lemma 3.17 we have that $I_{S:a}: (x_h) = I_{(S:h):a}$. Since by Lemma 3.16 S:h is a generalized tower set, observing that $f_h = \prod_{j \in (S:h)^a} x_j$, we can apply the inductive hypothesis to assert that

proj-dim
$$((I_{S:a}: (x_h)) + (f_h)) \le 2.$$

We set $J := (I_{S:a} : (x_h)) + (f_h)$. Now let us consider the exact sequence

 $0 \to I_{S:a} + (f) \to J \oplus (I_{S:a} + (x_h)) \to J + (x_h) \to 0.$

Since proj-dim $(J + (x_h)) \leq 3$ and $I_{S:a} + (x_h)$ is a complete intersection ideal of height 2, we can conclude that proj-dim $(I_{S:a} + (f)) \leq 2$.

Now we want to give a converse of the previous theorem. More precisely we want to prove that every monomial squarefree aCM ideal of height two is supported on a suitable generalized tower set. To do this we introduce some preparatory material.

Definition 3.20. Let $U \subseteq C_{2,n}$. We will say that U is generalized towerizable set if there exists an ordinante function $\omega : C_{2,n} \to (\mathbb{Z}^2_+)^*$ and a permutation τ on $\pi_2(\omega(U))$ such that $\tau\omega(U)$ is a generalized tower set.

Let $U \subseteq C_{2,n}$ be an aCM set; then, by the Hilbert-Burch theorem, I_U is a determinantal ideal generated by the maximal minors of a matrix of size $(r+1) \times r$.

Lemma 3.21. If $I \subset R$ is an aCM monomial ideal of height 2 then it admits a Hilbert-Burch matrix of the form

where D_i and M_{ij} are monomials, $D_i \neq 0$ and $M_{ij} \neq 0$ and D_i is in the position (i,i) (we enumerate the rows from 0 to r and the columns from 1 to r).

Proof. We take the minimal monomial set G of generators for I, then the first syzygy module is minimally generated by a set Φ of r elements acting each only on two of such generators. Moreover there are at least two generators in G on which only one syzygy acts. Let f_0 be one of these generators and let ϕ_1 be the syzygy acting on f_0 and let f_1 be the other generator on which acts ϕ_1 . Now we call $\phi_2, \ldots, \phi_{\alpha_1}$ all the other syzygies in Φ acting respectively on f_1 and $f_2, \ldots, f_{\alpha_1} \in G$. By iterating this procedure we get our matrix. \Box

Definition 3.22. An Hilbert-Burch matrix of the type as in Lemma 3.21 will be called a matrix of *standard form*.

Definition 3.23. Let $\mathcal{M} = (m_{ij})$ be an Hilbert-Burch matrix of *standard* form of size $(r+1) \times r$. Let $\sigma : [r] \to \{0, \ldots, r-1\}$ be the application such that $\sigma(j)$ is the only integer less than j such that $m_{\sigma(j)j} \neq 0$.

From now on we set M_i for $M_{\sigma(i),i}$.

Remark 3.24. Note that $\sigma(1) = 0$, $\sigma(2) = 1$ and, for j > 2, $\sigma(j) \ge \sigma(j-1) > 0$.

Given $j \in \{1, \ldots, r\}$ we denote with m(j) the set

$$m(j) := \{j, \sigma(j), \sigma^2(j), \dots, \sigma^h(j)\},\$$

where h is the only integer such that $\sigma^h(j) = 1$. We write $u \notin m(j)$ to mean $u \in [n] \setminus m(j)$.

Remark 3.25. Note that if $i \in m(j)$ then $m(i) \subseteq m(j)$.

We denote by f_i the determinant of the matrix obtained by removing the row *i* for $0 \le i \le r$. By the Hilbert-Burch theorem we have that

$$\{f_0,\ldots,f_r\}$$

is a minimal set of generators for I. Note that $f_0 = D_0 \cdots D_r$. In the following proposition will compute all the other generators.

Proposition 3.26. For any $i \in \{1, ..., r\}$, with the above notation, we have

$$f_i = \prod_{j \in m(i)} M_j \cdot \prod_{j \notin m(i)} D_j.$$

Proof. Let $i \in \{1, \ldots, r\}$, and let H be the square matrix given by \mathcal{M} without the row containing D_i . Since M_i is the only entry in the *i*-th column of H, we compute the determinant f_i by using the Laplace expansion along its *i*-th column. Thus $f_i = M_i G_1$, where G_1 is the determinant of the matrix H_1 obtained from H by deleting the row $\sigma(i)$ and the *i*-th column. Note that $M_{\sigma(i)}$ is the only entry in the $\sigma(i)$ -th column of H_1 , hence $f_i = M_i M_{\sigma(i)} G_2$, where G_2 is the determinant of the matrix H_2 obtained from H_1 by deleting the row $\sigma^2(i)$ and the $\sigma(i)$ -th column. So, by iterating this computation, we get $f_i = \prod_{j \in m(i)} M_j \cdot G'$, where G' is the determinant of the matrix H' obtained from H by deleting the rows $\sigma(j)$ and the columns j, for all $j \in m(i)$. Finally, we observe that H' is an upper triangular matrix, therefore $G' = \prod_{j \notin m(i)} D_j$.

Now we want to construct a generalized tower set starting from a n Hilbert-Burch matrix of standard form \mathcal{M} .

Definition 3.27. We define

$$U'_{\mathcal{M}} := \{\{i, j\} \in C_{2, 2r} \mid i < j \le r \text{ and } i \notin m(j)\}$$

and

$$U''_{\mathcal{M}} := \{\{i, j\} \in C_{2,2r} \mid i \le r < j \text{ and } j - r \in m(i)\}$$

Finally we set

$$U_{\mathcal{M}} := U'_{\mathcal{M}} \cup U''_{\mathcal{M}}.$$

- **Proposition 3.28.** 1) If $\{u, v\} \in U'_{\mathcal{M}}$ then either $u \notin m(i)$ or $v \notin m(i)$ for every $1 \le i \le r$.
 - 2) If $\{u, v\} \in U''_{\mathcal{M}}$, with u < v, then either $u \notin m(i)$ or $v r \in m(i)$ for every $1 \le i \le r$.
- *Proof.* 1) Let $u, v \in m(i)$ with u < v. Then $v = \sigma^h(i)$ and $u = \sigma^k(i)$ with h < k. Then $u = \sigma^{k-h}\sigma^h(i) = \sigma^{k-h}(v)$, i.e. $u \in m(v)$, hence $\{u, v\} \notin U'_{\mathcal{M}}$.
 - 2) Let $\{u, v\} \in U''_{\mathcal{M}}$, with u < v, such that $u \in m(i)$. Since $v r \in m(u)$ we get $v r \in m(i)$.

Proposition 3.29. $U_{\mathcal{M}}$ is connected.

Proof. Let $\{i, j\}, \{u, v\} \in U_{\mathcal{M}}$, with i < j, u < v and u < i. If $u \notin m(i)$ then $u < i \leq r$, so $\{u, i\} \in U'_{\mathcal{M}}$. If $u \in m(i)$, we have to consider two cases.

If $v \leq r$ then $\{u, v\} \in U'_{\mathcal{M}}$, hence $u \notin m(v)$. Since $u \in m(i)$ we get $m(i) \not\subseteq m(v)$ i.e. $i \notin m(v)$. Moreover, by Proposition 3.28 we have $v \notin m(i)$ and so $\{i, v\} \in U'_{\mathcal{M}}$.

If v > r then, by Proposition 3.28, we get $v - r \in m(i)$, so $\{i, v\} \in U''_{\mathcal{M}}$. \Box

Now we set $\mathcal{T} := U'_{\mathcal{M}} \cup \{\{i, j\} \in U''_{\mathcal{M}} \mid j - r \in m(r)\}$ and $\mathcal{S}_0 := U_{\mathcal{M}} \setminus \mathcal{T}$, so $\mathcal{S}_0 = \{\{i, j\} \in C_{2,2r} \mid i \leq r < j \text{ and } j - r \in m(i) \setminus m(r)\}.$

For every $i \in [r]$, we set $\mu_i := \max(m(i) \cap m(r))$. We set

$$\mathcal{V}_i^{(1)} := \{\{u, v\} \in U'_{\mathcal{M}} \mid \mu_u = \mu_v = \mu_i\}$$

and

$$r_i^{(1)} := \max \bigcup_{\alpha \in \mathcal{V}_i^{(1)}} \alpha \cup \{0\}.$$

Note that $r_i^{(1)} < r$. Moreover, we set

$$\mu_i^{(1)} := \begin{cases} \max(m(i) \cap m(r_i^{(1)})) & \text{if } r_i^{(1)} > 0\\ 0 & \text{otherwise} \end{cases}.$$

Now by induction let us suppose that we defined $\mu_i^{(k)}$ for $1 \le k < h$. Then we set

$$\mathcal{V}_i^{(h)} := \{\{u, v\} \in \mathcal{V}_i^{(h-1)} \mid \mu_u^{(h-1)} = \mu_v^{(h-1)} = \mu_i^{(h-1)}\}$$

and

$$r_i^{(h)} := \max \bigcup_{\alpha \in \mathcal{V}_i^{(h)}} \alpha \cup \{0\}.$$

Note that if $r_i^{(h-1)} > 0$ then $r_i^{(h)} < r_i^{(h-1)}$. If $r_i^{(h-1)} = 0$ then $r_i^{(h)} = 0$. Moreover we set

$$\mu_i^{(h)} := \begin{cases} \max(m(i) \cap m(r_i^{(h)})) & \text{if } r_i^{(h)} > 0\\ 0 & \text{otherwise} \end{cases}$$

In the sequel we will set $\mu_i^{(0)} := \mu_i$. Note that by definition the sequence $(\mu_i^{(h)})_{h>0}$ vanishes definitively.

Lemma 3.30. If $\{i, j\} \in U'_{\mathcal{M}}$ then there exists h such that $\mu_i^{(h)} \neq \mu_j^{(h)}$. Proof. If $\mu_i^{(h)} = \mu_j^{(h)}$ for every h then $\mathcal{V}_i^{(h)} = \mathcal{V}_j^{(h)} \neq \emptyset$ for every h. Conse-

Proof. If $\mu_i^{(n)} = \mu_j^{(n)}$ for every h then $\mathcal{V}_i^{(n)} = \mathcal{V}_j^{(n)} \neq \emptyset$ for every h. Consequently $(\mu_i^{(h)})_{h\geq 0}$ should not be definitively null, a contradiction.

For every $\{i, j\} \in U''_{\mathcal{M}}$, with i > j, we set $\omega(\{i, j\}) := (i, j)$.

For every $\{i, j\} \in U'_{\mathcal{M}}$, let t be the smallest integer such that $\mu_i^{(t)} \neq \mu_j^{(t)}$ (see Lemma 3.30). Then we set

$$\omega(\{i,j\}) := \begin{cases} (i,j) \text{ if } \mu_i^{(t)} < \mu_j^{(t)} \\ (j,i) \text{ if } \mu_i^{(t)} > \mu_j^{(t)} \end{cases}$$

Now we set $\overline{U} := \omega(U_{\mathcal{M}}) \subset (\mathbb{Z}^2_+)^*$.

Observe that for $1 \leq i \leq r$, $(r+1,i) \in \overline{U}$, hence $\pi_2(\overline{U}) = [r]$.

Lemma 3.31. Let $i, j \in [r]$, with $\mu_i \leq \mu_j$. Let h > r be such that $(h, i) \in \omega(\mathcal{T})$. Then $(h, j) \in \omega(\mathcal{T})$.

Proof. Since $\{h, i\} \in \mathcal{T}$ and h > r we get $h - r \in m(i) \cap m(r)$. On the other hand, $\mu_i \leq \mu_j$ implies that $m(i) \cap m(r) \subseteq m(j) \cap m(r)$. Therefore $h - r \in m(j) \cap m(r)$, so $(h, j) \in \omega(\mathcal{T})$.

Theorem 3.32. $U_{\mathcal{M}}$ is a generalized towerizable set.

Proof. Let us consider \overline{U} . It is enough to find a permutation τ on [2r] such that $\tau(\overline{U})$ is a generalized tower set.

We set $\overline{T} := \omega(\mathcal{T})$. We want to show that the set $\{\overline{T}_i\}_{1 \leq i \leq r}$ is totally ordered by inclusion. Let $i, j \in [r], i \neq j$.

Case $1 : \mu_i^{(k)} = \mu_j^{(k)}$ for every $k \ge 0$. In this case we will show that $\overline{T}_i = \overline{T}_j$. Of course it is enough to show that $\overline{T}_i \subseteq \overline{T}_j$. Take $h \in \overline{T}_i$ i.e. $(h,i) \in \overline{T}$. If h > r by Lemma 3.31 $(h,j) \in \overline{T}$. If $h \le r$ let t be the smallest integer such that $\mu_h^{(t)} < \mu_i^{(t)} = \mu_j^{(t)}$. By the minimality of $t, r_i^{(t)} = r_h^{(t)}$, hence from $m(h) \cap m(r_h^{(t)}) \subset m(i) \cap m(r_i^{(t)}) = m(j) \cap m(r_j^{(t)})$ it follows that $j \notin m(h)$. On the other hand if $h \in m(j)$, since $\mu_i^{(k)} = \mu_j^{(k)}$ for every $k \ge 0$, by Lemma 3.30, $\{i, j\} \notin U'_{\mathcal{M}}$ i.e. either $i \in m(j)$ or $j \in m(i)$. If $j \in m(i)$ then $h \in m(i)$ and this contradicts that $\{h, i\} \in \mathcal{T}$. If $i \in m(j)$, since both h and i belong to m(j) then either $h \in m(i)$ or $i \in m(h)$, again a contradiction with $\{h, i\} \in \mathcal{T}$. Therefore $h \notin m(j)$. This together with $j \notin m(h)$, as we saw, implies that $\{h, j\} \in \mathcal{T}$. By the inequality $\mu_h^{(t)} < \mu_j^{(t)}$, we get $(h, j) \in \overline{T}$.

Case 2 : let t be the smallest integer such that $\mu_i^{(t)} \leq \mu_j^{(t)}$. We claim that $\overline{T}_i \subseteq \overline{T}_j$. Take $h \in \overline{T}_i$ i.e. $(h, i) \in \overline{T}$. If h > r, using again Lemma 3.31, we

are done. So we can assume $h \leq r$. Let *s* be the smallest integer such that $\mu_h^{(s)} < \mu_i^{(s)}$. Assume $s \leq t$ (the same argument will work in the case s > t). We have to prove that $(h, j) \in \overline{T}$. From the inequalities $\mu_h^{(s)} < \mu_i^{(s)} \leq \mu_j^{(s)}$ it is enough to show that $\{h, j\} \in \mathcal{T}$ i.e. $h \notin m(j)$ and $j \notin m(h)$. Using the same inequalities we get

$$m(h) \cap m(r_h^{(s)}) \subset m(i) \cap m(r_i^{(s)}) \subseteq m(j) \cap m(r_j^{(s)}).$$

As above, by the minimality of s, we have $r_h^{(s)} = r_i^{(s)} = r_j^{(s)}$, so we can deduce that $j \notin m(h)$. On the other hand, if $h \in m(j)$, since there is $k \in m(j) \cap m(r_j^{(s)})$, with $k \notin m(h)$, we obtain that $h \in m(k)$ (note that we are using the fact that $h, k \in m(j)$ and $k \notin m(h)$). Since $m(k) \subseteq m(j) \cap m(r_j^{(s)})$, we get $h \in m(r_j^{(s)})$, so $h \in m(i)$, a contradiction, hence $h \notin m(j)$ and we are done.

Let τ be a permutation on [2r], such that $\tau(i) < \tau(j)$ whenever $1 \leq i, j \leq r$ and $\overline{T}_i \supseteq \overline{T}_j$. Moreover $\tau(i) = i$ for $i \geq r+1$. We denote by $S = \tau \omega(U_{\mathcal{M}})$. We want to show that S is a generalized tower set i.e. we need to prove the four properties stated in Definition 3.12.

1) Clearly S is connected by Lemma 3.29.

2) Now set $T := \tau(\overline{T})$ and $S_0 = S \setminus T$. By the properties of τ , T is a tower set.

3) Note that $\pi_2(S) = \pi_2(T) = [r]$ and $\pi_1(T) \subseteq [r] \cup \{r+1 \leq i \leq 2r \mid i-r \in m(r)\}$. Therefore if $(i,j) \in S_0$, $\{\tau^{-1}(i), \tau^{-1}(j)\} \in U''_{\mathcal{M}}$. This implies that i > r so $i \notin \pi_2(T)$. On the other hand since $\{\tau^{-1}(i), \tau^{-1}(j)\} \notin \mathcal{T}$, we have $i-r \notin m(r)$ i.e. $i \notin \pi_1(T)$. Let $u = \tau^{-1}(j)$. Then $(i,u) \notin \overline{T}$, so $\{i,u\} \in U''_{\mathcal{M}}$. We claim that $(u,r) \in \overline{T}$, which will imply that $j \in \pi_1(T)$. We need to show that $u \notin m(r)$. We know that $\{i,u\} \in U''_{\mathcal{M}}$, so $i-r \in m(u)$. If $u \in m(r)$ then $i-r \in m(r)$ therefore $(i,j) \in T$, a contradiction. Consequently $j \in \pi_1(T) \cap \pi_2(T)$.

4) Now we would like to prove that if $(i, \tau(j)) \in S_0$ and $\tau(u) \in F_T(\tau(j))$ then $(i, \tau(u)) \in S_0$, i.e. $\{i, u\} \in S_0$ hence we have to show that $i - r \in m(u)$ and $i - r \notin m(r)$. Since $(i, \tau(j)) \in S_0$ we have $i - r \in m(j) \setminus m(r)$. Since $\tau(u) \in F_T(\tau(j))$ we have $T_{\tau(j)} \subset T_{\tau(u)}$ and $(\tau(j), \tau(u)) \notin T$. From this we deduce that $\{u, j\} \notin U'_{\mathcal{M}}$, consequently either $j \in m(u)$ or $u \in m(j)$. If $j \in m(u)$ since $i - r \in m(j)$ we get $i - r \in m(u)$ and we are done. If $u \in m(j)$ since also $i - r \in m(j)$ we get either $u \in m(i - r)$ or $i - r \in m(u)$. If $u \in m(i - r)$ and $i - r \notin m(u)$, take $v \in \overline{T}_u \setminus \overline{T}_j$ and consider $\{u, v\}$, $\{i, j\} \in U_{\mathcal{M}}$. Since $U_{\mathcal{M}}$ is connected we have that one of the following sets must belong to $U_{\mathcal{M}}$:

$$\{u, i\}, \{u, j\}, \{v, i\}, \{v, j\}.$$

Note that $\{u, i\} \notin U_{\mathcal{M}}$ since $i - r \notin m(u)$. Moreover $\{u, j\} \notin U_{\mathcal{M}}$ (see above). If $\{v, i\} \in U_{\mathcal{M}}$, then $(i, v) \in \omega(\mathcal{S}_0)$, so $i - r \in m(v)$, consequently $u \in m(v)$, which contradicts that $\{u, v\} \in U_{\mathcal{M}}$.

If $\{v, j\} \in U_{\mathcal{M}}$, then $(j, v) \in \omega(S_0)$. Therefore $j \in \overline{T}_v \setminus \overline{T}_u$ and $v \in \overline{T}_u \setminus \overline{T}_v$, a contradiction since \overline{T}_u and \overline{T}_v are comparable by inclusion. Consequently $i - r \in m(u)$ and we are done. Remark 3.33. We note that if a Hilbert-Burch matrix of standard form $\mathcal{M} = (m_{i,j})$ is bidiagonal (i.e. the only non zero entries are $m_{i,i}$ and $m_{i,i+1}$) then $U_{\mathcal{M}}$ is a towerizable set. This follows by Theorem 3.32, since, using the same notation as above, $\mathcal{S}_0 = \emptyset$. Vice versa if $U_{\mathcal{M}}$ is a towerizable set then it is easy to build a bidiagonal Hilbert-Burch matrix of standard form \mathcal{M}' such that $I(\mathcal{M}) = I(\mathcal{M}')$. This generalizes a result of Ahn and Shin in [AS].

In order to get our main result we need the following Lemma.

Lemma 3.34. With the same terminology as above, let \mathcal{M} be a Hilbert-Burch matrix of standard form. Let $\{i, j\}, \{u, v\}$ be two different elements in $U_{\mathcal{M}}$. Then, if we set $H_i := D_i$ and $H_{r+i} := M_i$ for i = 1, ..., r,

$$depth(H_i, H_i, H_u, H_v) \ge 3.$$

Proof. We need to distinguish three possibilities. If $\{i, j\}, \{u, v\} \in U'_{\mathcal{M}}$ then $i, j, u, v \leq r$, since $\prod_{h=1..r} D_h$ is a minimal generator of $I(\mathcal{M})$ and $|\{i, j, u, v\}| \geq 3$ we are done. If $\{i, j\}, \{u, v\} \in U''_{\mathcal{M}}, \text{ say } i, u \geq r$, then if j = v we have $i - r \in m(j)$ and $u - r \in m(j)$. So, by Proposition 3.26, $H_i H_u$ is a factor of f_j hence H_i, H_u are coprime as f_j is squarefree, then depth $(H_i, H_j, H_u) = 3$. If j < v, then $i - r \in m(j)$ and $v \notin m(j)$. So, by Proposition 3.26, $H_i H_v$ is a factor of f_j hence H_i, H_v are coprime as f_j is squarefree, then depth $(H_i, H_j, H_v) = 3$. If $\{i, j\} \in U'_{\mathcal{M}}, \{u, v\} \in U''_{\mathcal{M}}$, with v < u, when $|\{i, j, v\}| = 3$ we are done, otherwise say v = j then $H_u H_i$ is a factor of f_j so depth $(H_i, H_j, H_u) = 3$.

Collecting all the results of this section we are ready to proof the main result.

Theorem 3.35. Let $I \subset k[x_1, \ldots, x_n]$ be a monomial squarefree of height 2. Then I is aCM iff it defines a generalized tower scheme.

Proof. Let us suppose that I is aCM. Then $I = I(\mathcal{M})$ for some \mathcal{M} of standard form of size $(r+1) \times r$ (see Lemma 3.21 and Definition 3.22). By Theorem 3.32 $U_{\mathcal{M}}$ is a generalized towerizable set. Let ω and τ be as in the proof of Theorem 3.32. Let $S := \tau(\omega(U_{\mathcal{M}}))$, which is a generalized tower set. For $j \in \pi_1(S)$, we set

$$f_{1j} := \begin{cases} D_{\tau^{-1}(j)} & \text{ for } j \le r\\ M_{j-r} & \text{ for } j > r \end{cases}.$$

For $j \in \pi_2(S) = [r]$, we set $f_{2j} := D_{\tau^{-1}(j)}$. Also we set

$$\mathcal{F}_i := \{ f_{ij} \mid j \in \pi_i(S) \}, \ i = 1, 2.$$

By Lemma 3.34, \mathcal{F}_1 and \mathcal{F}_2 satisfy the conditions of genericity required by Definition 3.13.

We claim that $I_S(\mathcal{F}_1, \mathcal{F}_2) = I$. At first we show that $I \subseteq I_S(\mathcal{F}_1, \mathcal{F}_2)$. Indeed, let g_k be the maximal minor obtained from \mathcal{M} by deleting the kth row and take any $(i, j) \in S$. We need to show that $g_k \in (f_{1i}, f_{2j})$ for every k. Since $g_0 = \prod_{1 \leq i \leq r} D_i$ and $f_{2j} = D_{\tau^{-1}(j)}, g_0 \in (f_{1i}, f_{2j})$. Assume $k \geq 1$. Since $(i, j) \in S$ we have $\{\tau^{-1}(i), \tau^{-1}(j)\} \in U_{\mathcal{M}} = U'_{\mathcal{M}} \cup U''_{\mathcal{M}}$. If

References

 $\{\tau^{-1}(i), \tau^{-1}(j)\} \in U'_{\mathcal{M}}$ then $(f_{1i}, f_{2j}) = (D_{\tau^{-1}(i)}, D_{\tau^{-1}(j)})$ and by Proposition 3.28 we get either $\tau^{-1}(i) \notin m(k)$ or $\tau^{-1}(j) \notin m(k)$ therefore by Proposition 3.26, $g_k \in (f_{1i}, f_{2j})$. We proceed analogously if $\{\tau^{-1}(i), \tau^{-1}(j)\} \in U''_{\mathcal{M}}$.

Now we show that $I_S(\mathcal{F}_1, \mathcal{F}_2) \subseteq I$. Let $g \in I_S(\mathcal{F}_1, \mathcal{F}_2)$ be a squarefree monomial. Let

$$E_g = \{0\} \cup \{1 \le h \le r \mid g \in (M_h) \text{ and } g \notin (D_h)\}.$$

We set $e := \max E_g$. We claim that $g \in (g_e)$.

Assume e = 0 and let $h \in [r]$. If $g \in (M_h)$ then $g \in (D_h)$ by the maximality of e. If $g \notin (M_h)$ then $g \in (D_h)$ since $\{r + h, h\} \in U''_{\mathcal{M}}$.

Assume e > 0. Remind that $g_e = \prod_{j \in m(e)} M_j \cdot \prod_{j \notin m(e)} D_j$. If $g \in (M_h)$ then $g \in (D_h)$ for h > e by the maximality of e. If $g \notin (M_h)$ then $g \in (D_h)$ since $\{r + h, h\} \in U'_{\mathcal{M}}$. Note that if e = 1 we are done. So we can assume e > 1. If $h \notin m(e)$ and h < e then $\{e, h\} \in U'_{\mathcal{M}}$, consequently $g \in (D_e, D_h)$, but $g \notin (D_e)$ so $g \in (D_h)$. Now let $h \in m(e)$, namely $\{r + h, e\} \in U'_{\mathcal{M}}$. Therefore $g \in (M_h, D_e)$, but $g \notin (D_e)$ so $g \in (M_h)$. Since g_e is a squarefree monomial, we showed that $g \in (g_e)$.

Vice versa let us suppose that I defines a generalized tower scheme. This means that $I = I_S(\mathcal{F}_1, \mathcal{F}_2)$ where S is a generalized tower set and $\mathcal{F}_1, \mathcal{F}_2$ are families of monomials, satisfying the conditions of Definition 3.13. By Theorem 3.19, S is aCM. So we get that I is aCM just applying Proposition 3.7.

References

- [AS] J. Ahn, and Y. S. Shin. The minimal free resolution of a star-configuration in \mathbb{P}^n and the Weak Lefschetz Property. J. Korean Math. Soc 49.2 (2012): 405-417.
- [GHM] A. V. Geramita, B. Harbourne, and J. Migliore. Star configurations in \mathbb{P}^n . J. of Algebra **376** (2013): 279-299.
- [GHS] A. V. Geramita, T.Y. Harima, S. Shin, *Extremal Point Sets and Gorenstein Ideals*. Adv. in Math. **152**(1) (2000), 78-119.
- [GS] A. V. Geramita, S. Shin, k-configurations in \mathbb{P}^3 all have extremal resolutions. J. of Algebra **213** (1999): 351–368.
- [MR] R. Maggioni, A. Ragusa, Connections between Hilbert function and geometric properties for a finite set of points in P², Le Matematiche, **39** (1984), no. 1-2, 153–170.
- [RZ] A. Ragusa, G. Zappalà, Partial intersection and graded Betti numbers, Beiträge zur Algebra und Geometrie, 44 (2003), no. 1, 285–302.

(G. Favacchio) Dip. di Matematica e Informatica, Università di Catania,

VIALE A. DORIA 6, 95125 CATANIA, ITALY E-mail address: favacchio@dmi.unict.it Fax number: +39095330094

(A. RAGUSA) DIP. DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI CATANIA,
 VIALE A. DORIA 6, 95125 CATANIA, ITALY
 E-mail address: ragusa@dmi.unict.it

Fax number: +39095330094

(G. ZAPPALÀ) DIP. DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI CATANIA, VIALE A. DORIA 6, 95125 CATANIA, ITALY

References

E-mail address: zappalag@dmi.unict.it *Fax number:* +39095330094