



An equivalent formulation of 0-closed sesquilinear forms

ROSARIO CORSO 

Abstract. In 1970, McIntosh introduced the so-called 0-closed sesquilinear forms and proved a corresponding representation theorem. In this paper, we give a simple equivalent formulation of 0-closed sesquilinear forms. The main underlying idea is to consider minimal pairs of non-negative dominating forms.

Mathematics Subject Classification. Primary 47A07; Secondary 47A10.

Keywords. Sesquilinear forms, 0-closed forms, Representation theorem, Minimal forms.

1. Introduction. Let \mathcal{H} be a Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$. Let $\mathcal{D}_1, \mathcal{D}_2$ be two subspaces of \mathcal{H} . A *sesquilinear form* (or, more simply, a *form*) \mathfrak{t} on $\mathcal{D}_1 \times \mathcal{D}_2$ is a map $\mathfrak{t} : \mathcal{D}_1 \times \mathcal{D}_2 \rightarrow \mathbb{C}$ which is linear in the first component and anti-linear in the second one. If $\mathcal{D}_1 = \mathcal{D}_2 = \mathcal{D}$, we write $\mathfrak{t}[f] := \mathfrak{t}(f, f)$ for $f \in \mathcal{D}$. A sesquilinear form \mathfrak{s} on $\mathcal{D} \times \mathcal{D}$ is said to be *non-negative* if $\mathfrak{s}[f] \geq 0$ for every $f \in \mathcal{D}$; *strictly-positive* if there exists $c > 0$ such that $\mathfrak{s}[f] \geq c\|f\|^2$ for every $f \in \mathcal{D}$. If \mathfrak{s} is non-negative, we denote by $\ker(\mathfrak{s})$ the subspace $\{f \in \mathcal{D} : \mathfrak{s}[f] = 0\}$.

Given a sesquilinear form \mathfrak{t} on $\mathcal{D}_1 \times \mathcal{D}_2$, with \mathcal{D}_2 dense in \mathcal{H} , it is possible to construct an operator T with domain

$$D(T) = \{f \in \mathcal{D}_1 : \exists h \in \mathcal{H}, \mathfrak{t}(f, g) = \langle h, g \rangle, \forall g \in \mathcal{D}_2\} \quad (1.1)$$

and defined as $Tf = h$, for all $f \in D(T)$, where h is the element in (1.1). The operator T is called *associated* to \mathfrak{t} and then the following representation holds

$$\mathfrak{t}(f, g) = \langle Tf, g \rangle, \quad \forall f \in D(T), g \in \mathcal{D}_2. \quad (1.2)$$

In the last decades, several theorems about the representation (1.2) have been given [1–5, 9, 11, 13–17]. The topic of the representation is connected to the Lebesgue decomposition (see [6–8, 12, 19]) as motivated in [8].

One of the classical representation theorems has been given for the so-called *closed* sesquilinear forms [14, Ch. VI]. We recall that a non-negative¹ sesquilinear form \mathfrak{s} on $\mathcal{D} \times \mathcal{D}$ is closed if, for any sequence of vectors $\{f_n\}$ of \mathcal{H} such that $f_n \rightarrow f$ and $\mathfrak{s}[f_n - f_m] \rightarrow 0$, one has $f \in \mathcal{D}$ and $\mathfrak{s}[f_n - f] \rightarrow 0$. The representation theorem for closed sesquilinear forms is useful, for instance, to define the Friedrichs extension of densely defined positive operators [14, Ch. VI] and a special sum of two operators [18].

In this paper, we specifically focus on 0-closed forms introduced and treated in [16, 17] by McIntosh in 1970. Hence, first of all, we recall the definition. Let \mathcal{D}_1 and \mathcal{D}_2 be dense subspaces of \mathcal{H} . A sesquilinear form \mathfrak{t} on $\mathcal{D}_1 \times \mathcal{D}_2$ is called *0-closed* [16, 17] if

- \mathcal{D}_1 and \mathcal{D}_2 can be made into Hilbert spaces \mathcal{H}_1 and \mathcal{H}_2 continuously embedded in \mathcal{H} with inner products $\langle \cdot, \cdot \rangle_1$ and $\langle \cdot, \cdot \rangle_2$ and norms $\| \cdot \|_1$ and $\| \cdot \|_2$, respectively;
- \mathfrak{t} is bounded with respect to $\| \cdot \|_1$ and $\| \cdot \|_2$, i.e., there exists $C > 0$ such that

$$|\mathfrak{t}(f, g)| \leq C \|f\|_1 \|g\|_2, \quad \forall f \in \mathcal{H}_1, g \in \mathcal{H}_2; \tag{1.3}$$

- the bounded operator $A : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ satisfying

$$\mathfrak{t}(f, g) = \langle Af, g \rangle_2, \quad \forall f \in \mathcal{H}_1, g \in \mathcal{H}_2, \tag{1.4}$$

is a bijection,²

For 0-closed forms, the following representation theorem holds.

Theorem 1.1 ([16, 17]). *Let \mathfrak{t} be a 0-closed form on $\mathcal{D}_1 \times \mathcal{D}_2$, where \mathcal{D}_1 and \mathcal{D}_2 are dense subspaces of \mathcal{H} . Then the operator T associated to \mathfrak{t} is densely defined, closed, and 0 belongs to the resolvent set of T . Moreover, also the sesquilinear form \mathfrak{t}^* given by*

$$\mathfrak{t}^*(f, g) = \overline{\mathfrak{t}(g, f)}, \quad \forall f \in \mathcal{D}_2, g \in \mathcal{D}_1,$$

is 0-closed and its associated operator is T^* .

The main scope of this paper is to prove an equivalent formulation of 0-closed forms (Theorem 2.4). In particular, we will employ the concept of minimal pairs of non-negative sesquilinear forms which dominate a given sesquilinear form (Definition 2.1). The auxiliary results Lemma 2.3 and Proposition 3.1 give some characterizations of minimal pairs assuming that the non-negative sesquilinear forms are closed.

2. The equivalent formulation. Throughout the paper, we denote by $\ker(S)$ and $R(S)$ the kernel and the range of an operator $S : D(S) \rightarrow \mathcal{H}_2$, respectively. We use the symbol $\mathcal{B}(\mathcal{H}_1, \mathcal{H}_2)$ for the set of bounded operators $S : \mathcal{H}_1 \rightarrow \mathcal{H}_2$. Firstly, we introduce the set of pairs of dominating forms. Let \mathcal{D}_1 and \mathcal{D}_2 be subspaces of \mathcal{H} and \mathfrak{t} a sesquilinear form on $\mathcal{D}_1 \times \mathcal{D}_2$. We denote by $\mathcal{M}(\mathfrak{t})$ the set of pairs $(\mathfrak{s}_1, \mathfrak{s}_2)$ of non-negative sesquilinear forms such that

¹Actually, the definition of closed forms can be given for a more general classes of forms, namely for semi-bounded or sectorial forms [14].

²The existence of the bounded operator $A : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ satisfying (1.4) is ensured by (1.3).

An equivalent formulation of 0-closed sesquilinear forms

- $\mathfrak{s}_1 : \mathcal{D}_1 \times \mathcal{D}_1 \rightarrow \mathbb{C}$ and $\mathfrak{s}_2 : \mathcal{D}_2 \times \mathcal{D}_2 \rightarrow \mathbb{C}$;
- for every $f \in \mathcal{D}_1$ and $g \in \mathcal{D}_2$, one has

$$|\mathfrak{t}(f, g)| \leq \mathfrak{s}_1[f]^{\frac{1}{2}} \mathfrak{s}_2[g]^{\frac{1}{2}}. \quad (2.1)$$

Definition 2.1. We say that a pair $(\mathfrak{s}_1, \mathfrak{s}_2) \in \mathcal{M}(\mathfrak{t})$ is *minimal* in $\mathcal{M}(\mathfrak{t})$ if, for every $(\mathfrak{p}, \mathfrak{q}) \in \mathcal{M}(\mathfrak{t})$ such that $\mathfrak{p} \leq \mathfrak{s}_1$ and $\mathfrak{q} \leq \mathfrak{s}_2$, we have $\mathfrak{p} = \mathfrak{s}_1$ and $\mathfrak{q} = \mathfrak{s}_2$.

Remarks 2.2. 1. If \mathfrak{s} is a non-negative sesquilinear form on $\mathcal{D} \times \mathcal{D}$, then trivially $(\mathfrak{s}, \mathfrak{s})$ belongs to $\mathcal{M}(\mathfrak{s})$ by the Cauchy-Schwarz inequality

$$|\mathfrak{s}(f, g)| \leq \mathfrak{s}[f]^{\frac{1}{2}} \mathfrak{s}[g]^{\frac{1}{2}},$$

and it is also minimal in $\mathcal{M}(\mathfrak{s})$.

2. If \mathfrak{t} is a sesquilinear form and $(\mathfrak{s}_1, \mathfrak{s}_2)$ is minimal in $\mathcal{M}(\mathfrak{t})$, then also $(\alpha \mathfrak{s}_1, \alpha^{-1} \mathfrak{s}_2)$ is minimal in $\mathcal{M}(\mathfrak{t})$ for any $\alpha > 0$. Anyway, even in the non-negative case, there might exist other less trivial minimal pairs. To make an example, let $\mathcal{H} = \mathbb{C}^2$ and \mathfrak{s} the non-negative form defined as follows

$$\mathfrak{s}(f, g) = 2f(1)\overline{g(1)} + 2f(2)\overline{g(2)}, \quad \forall f, g \in \mathbb{C}^2,$$

where $f = (f(1), f(2))$ and $g = (g(1), g(2))$. The pair $(\mathfrak{s}_1, \mathfrak{s}_2)$ made with

$$\mathfrak{s}_1(f, g) = 4f(1)\overline{g(1)} + f(2)\overline{g(2)}, \quad \forall f, g \in \mathbb{C}^2,$$

and

$$\mathfrak{s}_2(f, g) = f(1)\overline{g(1)} + 4f(2)\overline{g(2)}, \quad \forall f, g \in \mathbb{C}^2,$$

is in $\mathcal{M}(\mathfrak{s})$. Indeed, by the Cauchy-Schwarz inequality, for every $f, g \in \mathbb{C}^2$,

$$\begin{aligned} |s(f, g)| &= |2f(1) \cdot \overline{g(1)} + f(2) \cdot \overline{2g(2)}| \\ &\leq (4|f(1)|^2 + |f(2)|^2)^{\frac{1}{2}} (|g(1)|^2 + 4|g(2)|^2)^{\frac{1}{2}} \\ &= \mathfrak{s}_1[f]^{\frac{1}{2}} \mathfrak{s}_2[g]^{\frac{1}{2}}. \end{aligned}$$

Moreover, $(\mathfrak{s}_1, \mathfrak{s}_2)$ is minimal in $\mathcal{M}(\mathfrak{s})$. Indeed, let $(\mathfrak{p}, \mathfrak{q}) \in \mathcal{M}(\mathfrak{s})$ with $\mathfrak{p} \leq \mathfrak{s}_1$ and $\mathfrak{q} \leq \mathfrak{s}_2$. We have

$$2 = \mathfrak{s}((1, 0), (1, 0)) \leq \mathfrak{p}[(1, 0)]^{\frac{1}{2}} \mathfrak{q}[(1, 0)]^{\frac{1}{2}} \leq \mathfrak{s}_1[(1, 0)]^{\frac{1}{2}} \mathfrak{s}_2[(1, 0)]^{\frac{1}{2}} = 2,$$

so $\mathfrak{p}[(1, 0)]\mathfrak{q}[(1, 0)] = 4$. Since $\mathfrak{p} \leq \mathfrak{s}_1$ and $\mathfrak{q} \leq \mathfrak{s}_2$, we have $\mathfrak{p}[(1, 0)] = \mathfrak{s}_1[(1, 0)] = 2$ and $\mathfrak{q}[(1, 0)] = \mathfrak{s}_2[(1, 0)] = 1$. In the same way, we can prove that $\mathfrak{p}[(0, 1)] = \mathfrak{s}_1[(0, 1)] = 1$ and $\mathfrak{q}[(0, 1)] = \mathfrak{s}_2[(0, 1)] = 2$. Thus the non-negative forms \mathfrak{p} and \mathfrak{q} are completely determined knowing $\mathfrak{p}((1, 0), (0, 1))$ and $\mathfrak{q}((1, 0), (0, 1))$. In particular, denoting by $\mathfrak{s}_1 - \mathfrak{p}$ the sesquilinear form given by the difference between \mathfrak{s}_1 and \mathfrak{p} , we have

$$\begin{aligned} |\mathfrak{p}((1, 0), (0, 1))| &= |(\mathfrak{s}_1 - \mathfrak{p})((1, 0), (0, 1))| \\ &\leq (\mathfrak{s}_1 - \mathfrak{p})[(1, 0)]^{\frac{1}{2}} (\mathfrak{s}_1 - \mathfrak{p})[(0, 1)]^{\frac{1}{2}} = 0, \end{aligned}$$

where the first line holds because $\mathfrak{s}_1((1, 0), (0, 1)) = 0$ and the second line is valid by the Cauchy-Schwarz inequality since $\mathfrak{s}_1 - \mathfrak{p}$ is non-negative by the hypothesis on \mathfrak{p} . In conclusion, $\mathfrak{p} = \mathfrak{s}_1$. With similar arguments $\mathfrak{q} = \mathfrak{s}_2$, i.e., $(\mathfrak{s}_1, \mathfrak{s}_2)$ is minimal in $\mathcal{M}(\mathfrak{s})$.

In the end of Section 3, we will give another example, but in infinite dimension, showing several minimal pairs. Now we are going to prove a preliminary result (Lemma 2.3) which will be useful in the proof of our main claim, i.e., Theorem 2.4. We start with a particular representation of a sesquilinear form \mathfrak{t} on $\mathcal{D}_1 \times \mathcal{D}_2$ determined by a given pair of closed non-negative sesquilinear forms $(\mathfrak{s}_1, \mathfrak{s}_2)$ in $\mathcal{M}(\mathfrak{t})$. Let

$$\widetilde{\mathcal{D}}_1 = \ker(\mathfrak{s}_1)^\perp \cap \mathcal{D}_1 \quad \text{and} \quad \widetilde{\mathcal{D}}_2 = \ker(\mathfrak{s}_2)^\perp \cap \mathcal{D}_2, \tag{2.2}$$

where $\ker(\mathfrak{s}_1)^\perp$ and $\ker(\mathfrak{s}_2)^\perp$ are the orthogonal complements of $\ker(\mathfrak{s}_1)$ and $\ker(\mathfrak{s}_2)$ in \mathcal{H} , respectively. The restrictions $\widetilde{\mathfrak{s}}_1, \widetilde{\mathfrak{s}}_2$ of $\mathfrak{s}_1, \mathfrak{s}_2$ on $\widetilde{\mathcal{D}}_1, \widetilde{\mathcal{D}}_2$, respectively, are closed non-negative sesquilinear forms with $\ker(\widetilde{\mathfrak{s}}_1) = \{0\}$ and $\ker(\widetilde{\mathfrak{s}}_2) = \{0\}$. Now let

$$\langle f, f' \rangle_1 = \widetilde{\mathfrak{s}}_1(f, f') \quad \text{and} \quad \langle g, g' \rangle_2 = \widetilde{\mathfrak{s}}_2(g, g'), \quad f, f' \in \widetilde{\mathcal{D}}_1, g, g' \in \widetilde{\mathcal{D}}_2, \tag{2.3}$$

$$\|f\|_1 = \widetilde{\mathfrak{s}}_1[f]^\frac{1}{2} \quad \text{and} \quad \|g\|_2 = \widetilde{\mathfrak{s}}_2[g]^\frac{1}{2}, \quad f \in \widetilde{\mathcal{D}}_1, g \in \widetilde{\mathcal{D}}_2. \tag{2.4}$$

Since $\widetilde{\mathfrak{s}}_1$ and $\widetilde{\mathfrak{s}}_2$ are closed forms, $\widetilde{\mathcal{D}}_1$ and $\widetilde{\mathcal{D}}_2$ are complete with respect to the inner products $\langle \cdot, \cdot \rangle_1$ and $\langle \cdot, \cdot \rangle_2$ in (2.3), respectively. We denote by \mathcal{H}_1 and \mathcal{H}_2 the Hilbert spaces made in this way. By (2.1), the restriction of \mathfrak{t} on $\widetilde{\mathcal{D}}_1 \times \widetilde{\mathcal{D}}_2$ can be considered as a bounded sesquilinear form with respect to the norms $\|\cdot\|_1$ and $\|\cdot\|_2$ in (2.4) induced by $\widetilde{\mathfrak{s}}_1$ and $\widetilde{\mathfrak{s}}_2$, respectively. Then, by Riesz's theorem, there exists an operator $B \in \mathcal{B}(\mathcal{H}_1, \mathcal{H}_2)$ such that

$$\mathfrak{t}(f, g) = \langle Bf, g \rangle_2, \quad \forall f \in \widetilde{\mathcal{D}}_1, g \in \widetilde{\mathcal{D}}_2. \tag{2.5}$$

In particular, again by (2.1), B has norm³ $\|B\|$ less or equal 1. We have the following characterization.

Lemma 2.3. *Let \mathfrak{t} be a sesquilinear form on $\mathcal{D}_1 \times \mathcal{D}_2$, where \mathcal{D}_1 and \mathcal{D}_2 are subspaces of \mathcal{H} , and let $(\mathfrak{s}_1, \mathfrak{s}_2)$ be a pair of closed non-negative sesquilinear forms in $\mathcal{M}(\mathfrak{t})$. Then $(\mathfrak{s}_1, \mathfrak{s}_2)$ is minimal in $\mathcal{M}(\mathfrak{t})$ if and only if the operator B in (2.5) is a unitary operator.*

Proof. Let us assume that $(\mathfrak{s}_1, \mathfrak{s}_2)$ is minimal in $\mathcal{M}(\mathfrak{t})$. By (2.5),

$$|\mathfrak{t}(f, g)| \leq \|Bf\|_2 \|g\|_2 = \|Bf\|_2 \mathfrak{s}_2[g]^\frac{1}{2}, \quad \forall f \in \widetilde{\mathcal{D}}_1, g \in \widetilde{\mathcal{D}}_2. \tag{2.6}$$

Let $I : \mathcal{H} \rightarrow \mathcal{H}$ be the identity operator, $O_1 : \mathcal{H} \rightarrow \mathcal{H}$ and $O_2 : \mathcal{H} \rightarrow \mathcal{H}$ the orthogonal projections onto $\ker(\mathfrak{s}_1)^\perp$ and $\ker(\mathfrak{s}_2)^\perp$, respectively. We note that $\ker(\mathfrak{s}_1), \ker(\mathfrak{s}_2)$ are closed since $\mathfrak{s}_1, \mathfrak{s}_2$ are closed forms and that for $f \in \mathcal{D}_1, g \in \mathcal{D}_2$, we have $(I - O_1)f \in \ker(\mathfrak{s}_1) \in \mathcal{D}_1, (I - O_2)g \in \ker(\mathfrak{s}_2) \in \mathcal{D}_2$, so $O_1f \in \widetilde{\mathcal{D}}_1, O_2g \in \widetilde{\mathcal{D}}_2$. Let \mathfrak{p} be the sesquilinear form defined as follows

$$\mathfrak{p}(f, g) = \langle BO_1f, BO_1g \rangle_2, \quad \forall f, g \in \mathcal{D}_1.$$

Hence, the inequality (2.6) can be rewritten as

$$|\mathfrak{t}(f, g)| \leq \mathfrak{p}[f]^\frac{1}{2} \mathfrak{s}_2[g]^\frac{1}{2}, \quad \forall f \in \widetilde{\mathcal{D}}_1, g \in \widetilde{\mathcal{D}}_2. \tag{2.7}$$

³For simplifying the notation, we do not add a symbol to $\|B\|$ to specify the spaces \mathcal{H}_1 and \mathcal{H}_2 . We are going to do this also for other operators.

Considering that $\mathfrak{t}(f, g) = 0$ if $f \in \ker(\mathfrak{s}_1)$ or $g \in \ker(\mathfrak{s}_2)$ and that $\mathfrak{p}[O_1f] = \mathfrak{p}[f]$, $\mathfrak{s}_2[O_2g] = \mathfrak{s}_2[g]$ for every $f \in \mathcal{D}_1, g \in \mathcal{D}_2$, the inequality (2.7) extends by linearity for every $f \in \mathcal{D}_1$ and $g \in \mathcal{D}_2$. In other words, $(\mathfrak{p}, \mathfrak{s}_2) \in \mathcal{M}(\mathfrak{t})$.

As said before, $\|B\| \leq 1$ and then $\mathfrak{p}[f]^{\frac{1}{2}} = \|BO_1f\|_2 \leq \|O_1f\|_1 = \mathfrak{s}_1[O_1f]^{\frac{1}{2}} = \mathfrak{s}_1[f]^{\frac{1}{2}}$ for every $f \in \mathcal{D}_1$. Since $(\mathfrak{s}_1, \mathfrak{s}_2)$ is minimal in $\mathcal{M}(\mathfrak{t})$, we have $\|Bf\|_2 = \mathfrak{p}[f]^{\frac{1}{2}} = \mathfrak{s}_1[f]^{\frac{1}{2}} = \|f\|_1$ for every $f \in \widetilde{\mathcal{D}}_1$, i.e., B is an isometry. Because

$$|\mathfrak{t}(f, g)| \leq \|f\|_1 \|B^*g\|_1, \quad \forall f \in \widetilde{\mathcal{D}}_1, g \in \widetilde{\mathcal{D}}_2,$$

holds too, in the same way, we have that B^* is an isometry. In conclusion, B is unitary.

Now assume that B is a unitary operator. Let $(\mathfrak{p}, \mathfrak{q}) \in \mathcal{M}(\mathfrak{t})$ be such that $\mathfrak{p} \leq \mathfrak{s}_1$ and $\mathfrak{q} \leq \mathfrak{s}_2$. Then there exist $P \in \mathcal{B}(\mathcal{H}_1)$ and $Q \in \mathcal{B}(\mathcal{H}_2)$ satisfying

$$\|P\| \leq 1, \quad \|Q\| \leq 1, \quad (2.8)$$

$\mathfrak{p}[f] = \|Pf\|_1^2$ for every $f \in \widetilde{\mathcal{D}}_1$, and $\mathfrak{q}[g] = \|Qg\|_2^2$ for every $g \in \widetilde{\mathcal{D}}_2$. By [14, Ch. VI, Lemma 3.1], there exists $R \in \mathcal{B}(\mathcal{H}_p, \mathcal{H}_q)$ ⁴ such that

$$\|R\| \leq 1, \quad (2.9)$$

$$\mathfrak{t}(f, g) = \langle R Pf, Qg \rangle_2, \quad \forall f \in \widetilde{\mathcal{D}}_1, g \in \widetilde{\mathcal{D}}_2.$$

Therefore, by uniqueness of the associated operator, $B = Q^*RP$. Moreover,

$$\|f\|_1 = \|Bf\|_2 = \|Q^*RPf\|_2 \leq \|Pf\|_1 \leq \|f\|_1, \quad \forall f \in \widetilde{\mathcal{D}}_1,$$

by (2.8) and (2.9), i.e., $\mathfrak{p}[f] = \|Pf\|_1^2 = \|f\|_1^2 = \mathfrak{s}_1[f]$ for every $f \in \widetilde{\mathcal{D}}_1$. Furthermore, on $\ker(\mathfrak{s}_1)$ (which, as said before, is closed), \mathfrak{s}_1 and \mathfrak{p} are null. Hence, by linearity, $\mathfrak{p} = \mathfrak{s}_1$. Working in a similar way with B^* , we also find that $\mathfrak{q} = \mathfrak{s}_2$. Thus $(\mathfrak{s}_1, \mathfrak{s}_2)$ is minimal in $\mathcal{M}(\mathfrak{t})$. \square

Now we are ready to prove the main result of the paper.

Theorem 2.4. *Let \mathfrak{t} be a sesquilinear form on $\mathcal{D}_1 \times \mathcal{D}_2$, where \mathcal{D}_1 and \mathcal{D}_2 are dense subspaces of \mathcal{H} . The following statements are equivalent.*

- (i) \mathfrak{t} is 0-closed;
- (ii) there exists a pair $(\mathfrak{s}_1, \mathfrak{s}_2)$ of closed strictly-positive sesquilinear forms minimal in $\mathcal{M}(\mathfrak{t})$.

Proof. (i) \implies (ii) Let $\langle \cdot, \cdot \rangle_1$ and $\langle \cdot, \cdot \rangle_2$ be the inner products which make \mathcal{D}_1 and \mathcal{D}_2 two Hilbert spaces based on the definition of 0-closed forms. Let $A = U|A| = |A^*|U$ be the polar decomposition of the bounded operator A in (1.4). In particular, since A is bijective, U is a unitary operator and $|A|^{\frac{1}{2}} : \mathcal{H}_1 \rightarrow \mathcal{H}_1$, $|A^*|^{\frac{1}{2}} : \mathcal{H}_2 \rightarrow \mathcal{H}_2$ are bijective self-adjoint positive operators. By [10, Theorem 2.7], we also have $A = |A^*|^{\frac{1}{2}}U|A|^{\frac{1}{2}}$. Let us define

$$\mathfrak{s}_1(f, f') = \langle |A|^{\frac{1}{2}}f, |A|^{\frac{1}{2}}f' \rangle_1, \quad f, f' \in \mathcal{D}_1,$$

$$\mathfrak{s}_2(g, g') = \langle |A^*|^{\frac{1}{2}}g, |A^*|^{\frac{1}{2}}g' \rangle_2, \quad g, g' \in \mathcal{D}_2.$$

⁴ \mathcal{H}_p and \mathcal{H}_q are the completions of $\widetilde{\mathcal{D}}_1$ and $\widetilde{\mathcal{D}}_2$ with respect to the norms induced by \mathfrak{p} and \mathfrak{q} , respectively.

The sesquilinear forms \mathfrak{s}_1 and \mathfrak{s}_2 are strictly-positive and closed forms because $|A|^{\frac{1}{2}}, |A^*|^{\frac{1}{2}}$ are bijective and $\langle \cdot, \cdot \rangle_1, \langle \cdot, \cdot \rangle_2$ are strictly-positive and closed forms. Moreover,

$$|\mathfrak{t}(f, g)| = |\langle |A^*|^{\frac{1}{2}}U|A|^{\frac{1}{2}}f, g \rangle| = |\langle U|A|^{\frac{1}{2}}f, |A^*|^{\frac{1}{2}}g \rangle| \leq \mathfrak{s}_1[f]^{\frac{1}{2}}\mathfrak{s}_2[g]^{\frac{1}{2}},$$

i.e., $(\mathfrak{s}_1, \mathfrak{s}_2) \in \mathcal{M}(\mathfrak{t})$. Finally, $(\mathfrak{s}_1, \mathfrak{s}_2)$ is minimal in $\mathcal{M}(\mathfrak{t})$ by Lemma 2.3 because the operator B constructed as in (2.5) coincides with U which is a unitary operator.

(ii) \implies (i) Let

$$\langle f, f' \rangle_1 = \mathfrak{s}_1(f, f') \quad \text{and} \quad \langle g, g' \rangle_2 = \mathfrak{s}_2(g, g'), \quad f, f' \in \mathcal{D}_1, g, g' \in \mathcal{D}_2.$$

Since \mathfrak{s}_1 and \mathfrak{s}_2 are closed strictly-positive sesquilinear forms, \mathcal{D}_1 and \mathcal{D}_2 turn into Hilbert spaces continuously embedded in \mathcal{H} with inner products $\langle \cdot, \cdot \rangle_1$ and $\langle \cdot, \cdot \rangle_2$, respectively. Moreover, \mathfrak{t} is bounded with respect to the norms of these spaces. Finally, the last condition required in the definition of 0-closed forms is satisfied as a consequence of Lemma 2.3. \square

3. A supplementary result. In this section, we prove another characterization of minimal forms concerning a different representation in comparison to (2.5).

As previously, let \mathfrak{t} be a sesquilinear form on $\mathcal{D}_1 \times \mathcal{D}_2$ and $(\mathfrak{s}_1, \mathfrak{s}_2) \in \mathcal{M}(\mathfrak{t})$ with \mathfrak{s}_1 and \mathfrak{s}_2 non-negative and closed. By Kato's second representation theorem [14, Theorem VI.2.23], there exist positive self-adjoint operators H_1 and H_2 in \mathcal{H} with $D(H_1) = \mathcal{D}_1$ and $D(H_2) = \mathcal{D}_2$ such that $\mathfrak{s}_1[f] = \|H_1 f\|^2$ and $\mathfrak{s}_2[g] = \|H_2 g\|^2$ for every $f \in \mathcal{D}_1$ and $g \in \mathcal{D}_2$. We write $\overline{R(H_1)}$ and $\overline{R(H_2)}$ for the closures of $R(H_1)$ and $R(H_2)$ in \mathcal{H} , respectively. By (2.1) and [14, Ch. VI, Lemma 3.1], there exists a bounded operator $Q \in \mathcal{B}(\overline{R(H_1)}, \overline{R(H_2)})$ with $\|Q\| \leq 1$ such that

$$\mathfrak{t}(f, g) = \langle QH_1 f, H_2 g \rangle, \quad f \in \mathcal{D}_1, g \in \mathcal{D}_2.$$

Moreover, $\ker(H_1) = \ker(\mathfrak{s}_1)$, $\ker(H_2) = \ker(\mathfrak{s}_2)$, so the restrictions \widetilde{H}_1 and \widetilde{H}_2 of H_1 and H_2 on $\widetilde{\mathcal{D}}_1$ and $\widetilde{\mathcal{D}}_2$ (which are defined in (2.2)), respectively, are injective and we can also write

$$\mathfrak{t}(f, g) = \langle Q\widetilde{H}_1 f, \widetilde{H}_2 g \rangle, \quad f \in \widetilde{\mathcal{D}}_1, g \in \widetilde{\mathcal{D}}_2. \tag{3.1}$$

Proposition 3.1. *Let \mathfrak{t} be a sesquilinear form on $\mathcal{D}_1 \times \mathcal{D}_2$, where \mathcal{D}_1 and \mathcal{D}_2 are dense subspaces of \mathcal{H} and let $(\mathfrak{s}_1, \mathfrak{s}_2)$ be a pair of closed non-negative sesquilinear forms in $\mathcal{M}(\mathfrak{t})$. Then $(\mathfrak{s}_1, \mathfrak{s}_2)$ is minimal in $\mathcal{M}(\mathfrak{t})$ if and only if Q in (3.1) is a unitary operator.*

Proof. Let H_1 and H_2 be the operators introduced above. The operators Q and B of (2.5) are connected by the following relation

$$Q = \widetilde{H}_2 B \widetilde{H}_1^{-1} \quad \text{on } R(H_1).$$

Comparing the representations (2.5) and (3.1) for \mathfrak{t}^* instead of \mathfrak{t} , we also have

$$Q^* = \widetilde{H}_1 B^* \widetilde{H}_2^{-1} \quad \text{on } R(H_2),$$

so

$$QQ^* = \widetilde{H}_2 BB^* \widetilde{H}_2^{-1} \text{ on } R(H_2), \quad Q^*Q = \widetilde{H}_1 B^* B \widetilde{H}_1^{-1} \text{ on } R(H_1),$$

and

$$BB^* = \widetilde{H}_2^{-1} QQ^* \widetilde{H}_2 \text{ on } \widetilde{\mathcal{D}}_2, \quad B^*B = \widetilde{H}_1^{-1} Q^*Q \widetilde{H}_1 \text{ on } \widetilde{\mathcal{D}}_1.$$

Thus, by Lemma 2.3 and taking into account that $R(H_1)$, $R(H_2)$ are dense in their corresponding closures, $(\mathfrak{s}_1, \mathfrak{s}_2)$ is minimal in $\mathcal{M}(\mathfrak{t})$ if and only if Q is unitary. \square

Example. Let us consider $\mathcal{H} = \mathcal{D} = L^2(\mathbb{R})$ and a bounded measurable function $r : \mathbb{R} \rightarrow \mathbb{C}$. We write $N = \{x \in \mathbb{R} : r(x) = 0\}$. Let

$$\mathfrak{t}(f, g) = \int_{\mathbb{R}} r(x) f(x) g(x) dx, \quad \forall f, g \in \mathcal{H}. \quad (3.2)$$

The sesquilinear form \mathfrak{t} is bounded, and it is non-negative if and only if $r(x)$ is non-negative for a.e. $x \in \mathbb{R}$. For any measurable function $p : \mathbb{R} \rightarrow [0, +\infty)$, with $c \leq p(x) \leq d$ for some $c, d > 0$ and every $x \in \mathbb{R}$, we define

$$\mathfrak{s}_1(f, g) = \int_{\mathbb{R}} |r(x)| p(x) f(x) g(x) dx, \quad \forall f, g \in \mathcal{H}$$

and

$$\mathfrak{s}_2(f, g) = \int_{\mathbb{R}} |r(x)| p(x)^{-1} f(x) g(x) dx, \quad \forall f, g \in \mathcal{H}.$$

The sesquilinear forms \mathfrak{s}_1 and \mathfrak{s}_2 are non-negative (trivially closed by boundedness) and, by the Cauchy-Schwarz inequality, $(\mathfrak{s}_1, \mathfrak{s}_2) \in \mathcal{M}(\mathfrak{t})$. Moreover, again independently by p , we have $\ker(\mathfrak{s}_1) = \ker(\mathfrak{s}_2) = \{f \in \mathcal{H} : f(x) = 0 \text{ for a.e. } x \in \mathbb{R} \setminus N\} \equiv L^2(N)$, and so $\widetilde{\mathcal{D}}_1 = \widetilde{\mathcal{D}}_2 = \{f \in \mathcal{H} : f(x) = 0 \text{ for a.e. } x \in N\} \equiv L^2(\mathbb{R} \setminus N)$. The pair $(\mathfrak{s}_1, \mathfrak{s}_2)$ is minimal in $\mathcal{M}(\mathfrak{t})$ for any choice of p by Proposition 3.1. Indeed, one easily checks that the operators \widetilde{H}_1 and \widetilde{H}_2 are the multiplication operators by $|r(x)|^{\frac{1}{2}} p(x)^{\frac{1}{2}}$ and $|r(x)|^{\frac{1}{2}} p(x)^{-\frac{1}{2}}$ on the domain $L^2(\mathbb{R} \setminus N)$, respectively. Hence, by comparing (3.2) and (3.1), the operator Q is the multiplication operator by $\frac{r}{|r|}$ on the domain $L^2(\mathbb{R} \setminus N)$ and it is in particular unitary.

Acknowledgements. The author acknowledges a partial support by the “Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni” (GNAMPA-INdAM).

Funding Open access funding provided by Università degli Studi di Palermo within the CRUI-CARE Agreement.

Open Access. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- [1] Arendt, W., ter Elst, A.F.M.: Sectorial forms and degenerate differential operators. *J. Oper. Theory* **67**, 33–72 (2012)
- [2] Corso, R., Trapani, C.: Representation theorems for solvable sesquilinear forms. *Integral Equations Operator Theory* **89**, 43–68 (2017)
- [3] Corso, R.: A survey on solvable sesquilinear forms. In: *The Diversity and Beauty of Applied Operator Theory. Operator Theory: Advances and Applications*, vol. 268, pp. 167–177. Birkhäuser, Cham (2018)
- [4] Corso, R.: A Kato's second type representation theorem for solvable sesquilinear forms. *J. Math. Anal. Appl.* **462**(1), 982–998 (2018)
- [5] Corso, R.: Maximal operators with respect to the numerical range. *Complex Anal. Oper. Theory* **13**(3), 781–800 (2019)
- [6] Corso, R.: A Lebesgue-type decomposition for non-positive sesquilinear forms. *Ann. Mat. Pura Appl.* **198**(1), 273–288 (2019)
- [7] Corso, R.: A Lebesgue-type decomposition on one side for sesquilinear forms. In: *Theta Series in Advanced Mathematics, Conference Proceedings Timisoara 2018*, vol. 26 (2021)
- [8] Corso, R.: Lebesgue-type decomposition for sesquilinear forms via differences. *Arch. Math. (Basel)* **118**, 151–157 (2022)
- [9] Fleige, A., Hassi, S., de Snoo, H.: A Krein space approach to representation theorems and generalized Friedrichs extensions. *Acta Sci. Math. (Szeged)* **66**, 633–650 (2000)
- [10] Gesztesy, F., Malamud, M., Mitrea, M., Naboko, S.: Generalized polar decompositions for closed operators in Hilbert spaces and some applications. *Integral Equations Operator Theory* **64**, 83–113 (2009)
- [11] Grubišić, L., Kostykin, V., Makarov, K.A., Veselić, K.: Representation theorems for indefinite quadratic forms revisited. *Mathematika* **59**(1), 169–189 (2013)
- [12] Hassi, S., Sebestyén, Z., de Snoo, H.S.V.: Lebesgue type decompositions for nonnegative forms. *J. Funct. Anal.* **257**, 3858–3894 (2009)

- [13] Iborat, A., Llavona, J.G., Lledó, F., Pérez-Pardo, J.M.: Representation of non-semibounded quadratic forms and orthogonal additivity. *J. Math. Anal. Appl.* **495**(2), 124783 (2021)
- [14] Kato, T.: *Perturbation Theory for Linear Operators*. Springer, Berlin (1966)
- [15] McIntosh, A.: Representation of bilinear forms in Hilbert space by linear operators. *Trans. Amer. Math. Soc.* **131**(2), 365–377 (1968)
- [16] McIntosh, A.: Bilinear forms in Hilbert space. *J. Math. Mech.* **19**, 1027–1045 (1970)
- [17] McIntosh, A.: Hermitian bilinear forms which are not semibounded. *Bull. Amer. Math. Soc.* **76**, 732–737 (1970)
- [18] Reed, M., Simon, B.: *Methods of Modern Mathematical Physics I. Functional Analysis*. Academic Press, New York (1972)
- [19] Simon, B.: A canonical decomposition for quadratic forms with applications to monotone convergence theorems. *J. Funct. Anal.* **28**, 371–385 (1978)

ROSARIO CORSO
Dipartimento di Matematica e Informatica
Università degli Studi di Palermo
90123 Palermo
Italy
e-mail: rosario.corso02@unipa.it

Received: 13 August 2022

Revised: 3 September 2022

Accepted: 19 September 2022.