



Fragments of subdesert vegetation in the Mediterranean Region: the *Periploca angustifolia* maquis (*Periplocion angustifoliae*, *Quercetea ilicis*) in Sicily*

Lorenzo Gianguzzi^{1,2}, Antonino La Mantia³, Riccardo Rocca¹, Andrea Volpe⁴, Salvatore Cambria⁵

¹ Department of Agricultural, Food and Forest Sciences, University of Palermo, Palermo, Italy

² NBFC, National Biodiversity Future Center, Palermo, Italy

³ Department of Rural and Territorial Development, Regione Siciliana, Palermo, Italy

⁴ Division of Infectious Diseases, University of Verona, Verona, Italy

⁵ Department of Biological, Geological and Environmental Sciences, University of Catania, Catania, Italy

Corresponding author: Lorenzo Gianguzzi (lorenzo.gianguzzi@unipa.it)

Academic editor: Claudia Angiolini ♦ Received 17 October 2025 ♦ Accepted 15 November 2025 ♦ Published 22 December 2025

Abstract

The paper presents the results of a phytogeographic, ecological, phytosociological, and management-oriented study on the maquis vegetation dominated by *Periploca angustifolia* (class *Quercetea ilicis*) occurring along of southern coast of mainland Sicily. The species is a South-Mediterranean shrub, previously known in Europe only from some small islands surrounding Sicily (Linosa, Lampedusa, Pantelleria, and the Aegadian, and the Maltese Archipelagos), as well as from southeastern Spain and the Aegean islands of Crete, Khri, and Gavdos. Along with providing an updated overview of the species' distribution, the study explores the historical reasons for its presence in Sicily, where it is part of a diverse group of biogeographically related species mainly found along the southern coasts. This further supports the hypothesis of connection routes contracted with North Africa during the drying of the Mediterranean in the Messinian period. Based on a synoptic comparison with similar associations present in the central-Mediterranean, the plant community investigated is proposed as a new syntaxon (association *Asparago albi-Periplocetum angustifoliae* ass. nov.), framed within the alliance *Periplocion angustifoliae* (order *Pistacio-Rhamnetalia alaterni*). Its role as a vicariant association with respect to the association *Periploca angustifoliae-Euphorbietum dendroidis* (occurring on the small islands of the Sicilian Channel) is also discussed, together with floristic, synecological, syndynamic, and conservation data. Finally, issues concerning afforestation with alien species—often carried out on a large scale throughout the Mediterranean area—are discussed, as such interventions may sometimes negatively affect ecosystems typical of endemic or particularly rare species. This is precisely the case of *Periploca angustifolia* in the study area, whose residual population had been brought to the brink of extinction by conifer plantations, leading to the alteration and consequent disappearance of its habitats.

Keywords

Conservation, habitat fragmentation, *Periplocion angustifoliae*, relict Mediterranean maquis, semiarid ecosystems

* Topical Collection: "Vegetation classification: from classic to numeric approaches". Edited by Bianca Ott Andrade, Claudia Angiolini, Lorenzo Lazzaro.

Introduction

The flora of Sicily—including the surrounding islands—is recognized as having considerable phytogeographical interest (Giardina et al. 2007; Cañadas et al. 2014), being a major biodiversity hotspot of the Mediterranean region (Médail and Quézel 1997; Testolin et al. 2024). This is not only because it comprises a significant contingent of endemic species, but also due to the presence of several taxa that reach one of the limits of their distribution ranges precisely in this area (Brullo et al. 1995; Brullo and Brullo 2020). The southern coastal and subcoastal areas of mainland Sicily, for instance, host many species that originated from nearby North Africa and are either entirely absent or extremely rare throughout the rest of Europe. This is the case with several interesting shrubs such as *Searsia tripartita* (Brullo et al. 1998), *Tamarix arborea* (Brullo and Sciandrello 2006), *Soda oppositifolia* (Brullo et al. 2012), *Ziziphus lotus* (La Mantia and Scuderi 2021; La Mantia et al. 2025), *Lycium schweinfurthii* (Portal to the Flora of Italy 2025; POWO 2025), to which *Periploca angustifolia* may be added, recorded in a restricted site on Mount San Calogero, near Sciacca, which is the subject of this contribution. According to Quézel (1983, 1985), the migration of these taxa from arid North African areas into the Mediterranean region is linked to the

paleogeographic and paleoclimatic history of the Mediterranean Basin during the Early Cretaceous and particularly during the arid phases of the Messinian.

According to Plants of the World Online (POWO 2025), the species is widely distributed in North Africa (Mauritania, Western Sahara, Morocco, Algeria, Tunisia, Libya, and Egypt up to the Sinai area), with localized and fragmented occurrences across the Mediterranean Basin, from the Iberian Peninsula to the Aegean region. In particular, the European populations of this species are located in the southeastern part of Spain [Andalusia (provinces of Almería, Granada, and Málaga), Murcia, and the Valencian Community (Mota et al. 1997; Alcaraz et al. 2000; Mendoza-Fernández et al. 2020)], it was until now known from some small islands of the Sicilian Channel (Fig. 1) and from the Aegean islands of Crete, Gavdos, and Khrii (Brullo and Guarino 2000).

An additional record of the species has also been reported in Jordan at Wadi Rum (Taifour et al. 2017), whereas it should be considered extinct in Lebanon–Syria (Abd El-Ghani and Ahmed 2025) (Fig. 2). *Periploca angustifolia* is a characteristic species of xeric sub-Saharan environments, exhibiting distinct ecophysiological adaptations and a notable tolerance to soil water deficiency and drought stress (Noumi et al. 2013). It occurs on a range of soil types, though it shows a preference for calcareous substrates,



Figure 1. Distribution of *Periploca angustifolia* in the islands of Sicily Channel (grey dots) with the new stand of Mount S. Calogero (red dot).




Figure 2. Distribution of *Periploca angustifolia* (from POWO 2025, modif.); in red, the site subject of this survey.

typically in regions with mean annual precipitation below 400 mm, and occasionally as low as 100 mm (Chaieb and Boukhris 1998). Along the North African coasts, it has a broad distribution along the Mediterranean shoreline, from Mauritania to Egypt, extending southwards into desert areas. In Tunisia, it ranges from the northern regions to the Saharan area of Dhiba (Ennaieth et al. 2010), where it is a typical component of open maquis and garrigue, occurring in depressions, gullies, and wadis. It is often associated with *Searsia tripartita*, *Retama raetam* subsp. *raetam*, and *Stipa tenacissima*; in the Algerian Sahara, it is found in communities with *Searsia tripartita* and *Myrtus nivellei* (Benaissa and Djebbar 2018). It is also reported from various habitats of the western Mediterranean desert (sandy plains, ridges and inland plateaus, wadis, and roadsides), where it contributes to maintaining biodiversity and the ecological stability of these environments (Ahmed 2009).

In the eastern portion of its North African range, a gradual decline in populations has been documented, attributed by several authors mainly to anthropogenic pressures (El-Hadidi and Hosni 2000; Gamal et al. 2020). According to Abd El-Ghani and Ahmed (2025), the species shows a significant regression in Egypt and Sinai, with similar reports also concerning certain areas of Tunisia (Noumi et al. 2010). Among the principal factors is intense grazing by goats and camels (Boulos 2000), which adversely affects plant growth and reproductive performance.

Other important pressures are also represented by the degradation and decline of its preferred habitats, due to agricultural expansion and infrastructure development by humans (Salama et al. 2005; Abd El-Ghani and Ahmed 2025).

In this contribution, we first report the important discovery of a new population of the species in mainland Sicily, located on the southern slopes of Mount San Calogero (also known as Mount Kronio), in the municipality of Sciacca (Agrigento province). This discovery is indeed rather

surprising, not only because the new site significantly expands the species' range, but also due to its phytogeographical significance and the notable phytosociological and ecological value, including its potential future importance. Based on these premises, the results of a study are presented here, addressing the following objectives:

- a) to provide an updated distributional framework for the species in the Mediterranean territories;
- b) to discuss the historical factors underlying its occurrence along the southern coasts of the insular area, in comparison with other biogeographically analogous entities;
- c) to develop a phytosociological, synecological, and syndynamic characterization of the vegetation in which it occurs, in relation to other communities of the alliance *Periplocion angustifoliae* so far documented across the islands of the Sicilian Channel;
- d) to assess the pressures and threats affecting its habitat, with the aim of supporting the implementation of targeted conservation measures.

Study area

The new stand of *Periploca angustifolia* was found along the southern slopes of Mount San Calogero (397 m a.s.l.), a small hill that is part of a highly active volcanic region along the southwestern coast of Sicily (37°31'N – 13°6'E). About 60 km further southeast lies the submarine volcano Empedocles, which in 1831 gave rise to the island of Ferdinandea (or Graham Bank), later submerged following collapses caused by waves (Badino 2013). This hill, known since antiquity also as Mount Kronio—legendary home of the Greek deity Cronus (father of Zeus)—is catalogued among the most important geosites of the Sicilian Region, primarily

due to its rich labyrinth of caves affected by intense hydro-thermal phenomena related to secondary volcanism.

For this reason, an oriented nature reserve was established by the Sicilian Region (D.A. Territory and Environment No. 366/44 of 26/07/2000), covering 52.25 hectares, managed by the Department of Rural and Territorial Development (ORBD 2025a). The same area was subsequently included within a Natura 2000 site [SAC ITA040009 – Mount San Calogero (Sciacca); DM 07/04/2021 – Official Gazette 90 of 15/04/2021], together with adjacent territories of the elevation (Fig. 3), for a total area of approximately 127 hectares (ORBD 2025b).

Geomorphological and hydrothermal features

The relief of Mount San Calogero consists of a rock outcrop of the Upper Triassic–Lower Lias (stromatolitic limestones with *Megalodon* of Late Triassic–Early Lias) (Basilone 2018), characterized by Meso-Cenozoic limestones assigned to the “Saccense Unit” (Lentini and Carbone 2014). Its importance, however, is mainly due to a complex system of hypogene karst cavities, which also includes around twenty hydrothermal caves (Guidi and Verde 2001; Badino et al. 2009; Badino 2013; Commissione Grotte Boegan e Associazione Geografica La Venta 2013; Guidi et al. 2014). This geosite is considered a world unique site (ORBD 2025c) due to its speleo-thermal peculiarities (Floriani 1979; Forti 1991) and its significant

historical importance since prehistoric Sicily. Human presence is documented here as early as the 6th century BC in the upper caves, and up to 4000 years ago in the intermediate ones (Verde 2021). The entrances located at the top of the relief, namely the “Stufe di San Calogero,” consisting of two natural steam-filled caves, called the Antro di Dedalo (or Grotta di San Calogero and Grotta degli Animali) are incorporated within an ancient thermal complex built near the Sanctuary of San Calogero, patron saint of Sciacca.

This complex network of cavities and tunnels remains only partly explored due to its extreme conditions, including fluctuating temperatures (15–37 °C) and nearly 100% humidity (Di Piazza et al. 2017). These closed or semi-closed habitats resemble caves, where darkness and nutrient scarcity allow only specialized organisms—such as arthropods and microorganisms—to survive, while others, like bats, use them temporarily (Souza Maysa and Ferreira 2011; Smith et al. 2012). Recent micromycological studies have also identified various fungal taxa, including species from the genera *Penicillium*, *Cladosporium*, and *Trametes* (Di Piazza et al. 2017).

From an ecological perspective, it remains unclear how these internal environmental conditions influence the external characteristics of the site (e.g., the temperature of lithological substrates and soils). Nevertheless, the presence of *Periploca angustifolia* and *Searsia tripartita* (La Mantia et al. 2025)—rare shrub species in Sicily and typical of pre-desert formations—clearly highlights their potential climatic role within the area.




Figure 3. Area of S.A.C. ITA040009 and “Monte San Calogero (M. Kronio)” Nature Reserve, with distribution of *Periploca angustifolia* populations.

Moreover, from a geological perspective, this sector of Sicily has recently been considered by some authors as part of the foreland, similar to the Hyblaean area, whose territories may also have emerged (at least partially) after the deposition of pelagic carbonates, called “Trubi” (Ferranti et al. 2019). According to this hypothesis, the biological colonization of this area could have been even earlier, with a phytogeographical evolution potentially very different and highly significant for Sicily itself (Geraci et al. 2009; Troia et al. 2012; Ilardi et al. 2020).

Climatic features, bioclimate, and biogeographical characterization

From a climatic perspective, the mean annual precipitation is 564 mm, distributed over an average of 65 rainy days; precipitation peaks in winter, while a pronounced and prolonged minimum occurs during the spring–summer period (La Mantia et al. 2025). The mean annual temperature is 17.9 °C, with minimum average temperatures in February (7.9 °C) and maximum average temperatures in July (30.6 °C). According to the bioclimatic classification of Rivas-Martínez et al. (2004), the area falls within the lower dry Thermomediterranean belt (La Mantia et al. 2025).

From a biogeographical standpoint, the area is classified within the Mediterranean Region and the Central Mediterranean Subregion (Rivas-Martínez et al. 2004), and further assigned to the Sicilian Province, Eusiculo Sector, Western Subsector, and Drepano–Panormitano District (Brullo et al. 1995).

Land use and historical landscape evolution

The land use of the Mount San Calogero area reflects a long history of human activity, from prehistoric therapeutic and religious visits to quarrying, agriculture, and afforestation. Historical toponyms such as *Thermae Selnuntinae* and *Aquae Labodes* attest to the region's thermal heritage (Verde 2021). Quarrying of limestone outcrops (“Cave di pietra”) and extensive agriculture, including dry cultivation of carob and almond trees and sheep grazing, once shaped the landscape. Around the rocky areas, naturalized stands of *Opuntia ficus-indica* and *Agave* species mark past human cultivation.

Today, much of the territory forms part of the Sciacca publicly owned Forest within a regional nature reserve managed by the Sicilian Forestry Service. Since 1988, afforestation—mainly with *Pinus halepensis*, *Cupressus sempervirens*, and *Eucalyptus camaldulensis*—has altered the original low maquis dominated by *Periploca angustifolia*. The area has suffered repeated wildfires, notably in 1997, leading to the loss of large, forested zones and recurrent vegetation degradation (La Mantia et al. 2025). In recent years, additional fires have occurred, highlighting a trend of increasing frequency.

According to data from the Regional Forestry Information System, fires in this area have recurred almost annually since 2008, except in 2013 and 2022 (La Mantia et al. 2025). In the same fire-cleared clearings, numerous *Periploca angustifolia* individuals have regenerated from seeds, evidently dispersed by wind from residual plants located along hedges and at the margins of afforested areas. More developed shrubs are sometimes found entwined with dried branches or even the charred trunks of old conifers consumed by fire. The species tends to create sparse garrigue and low maquis, part of a complex phytocoenotic mosaic, interspersed with the more extensive *Hyparrhenia hirta* grasslands, as secondary aspects of two distinct native vegetation series: i) *Olea europaea* on rock outcrops; ii) *Searsia tripartita* on stony-detritic substrates.

In the surrounding coastal belt, ongoing residential expansion has further transformed the landscape, encroaching toward borders of the protected area. Other land-use types are also notable in the adjacent coastal strip, where in recent decades, intense residential urban expansion has been recorded, approaching some parts of the aforementioned protected areas.

Previous botanical investigations

From a floristic and phytosociological viewpoint, the area was little known until now, except in general terms. In fact, the toponym “Sciacca at Monte S. Calogero” (or “Monte Kronio”) is scarcely mentioned in the most important classical floristic works, such as those by Gussone (1827, 1843) and Lojacono (1888–1889, 1891, 1902–1903, 1904–1907, 1906, 1908–1909). As also noted in Giardina et al. (2007), for this locality there are only sporadic bibliographic references, mainly concerning Gussone's collections from 1827 and 1828. In particular, these include: *Searsia tripartita* (La Mantia et al. 2025), *Astragalus caprinus* subsp. *huetii*, *Micromeria microphylla* (Gussone 1834), *Stipa barbata* [Sciacca at S. Calogero (Gussone 1827)].

For other taxa, some specifically listed in the Standard Form of the Natura 2000 site (ORBD 2025c), there are only general bibliographic references (e.g., “Sciacca”), as in the cases of *Lonas annua*, *Malva flava* (sub *Lavatera agrigentina*), and *Glandora rosmarinifolia* [sub *Lithodora rosmarinifolia*]. For these taxa, given the absence of specific exsiccata from the area and/or the rarity of their preferred habitats, confirmation of their actual presence within the Natura 2000 site is necessary.

Indeed, among other geobotanical publications for this area, only a few reports can be mentioned: the sites of the endemic *Brassica villosa* subsp. *tineoi* (Otonello 1978; Raimondo et al. 1991), and forest aspects of *Olea europaea* stands of habitat 9320, *sensu* Directive 92/43/EEC (Rivieccio et al. 2021), as well as a more recent study on a small residual maquis patch of *Searsia tripartita* (La Mantia et al. 2025).

However, comprehensive monographic studies—both on the flora and the vegetation of this protected area—are still lacking. Such investigations would be highly desirable, particularly in light of recent and unexpected discoveries of species with North African affinities, including *Searsia tripartita* and *Periploca angustifolia*.

Materials and methods

Target species

Periploca angustifolia is a shrubby liana belonging to the Apocynaceae family, reaching 2(–3) m in height and adapted to particularly hot and dry climatic conditions (see Suppl. material 1). It has perennial stems with longitudinally striated bark, bearing short, rigid branches, sometimes twisted at the tips, with small terminal shoots. The leaves are lanceolate, coriaceous, either evergreen or summer-deciduous, the upper opposite; the blade is narrowly ovate to oblong-lanceolate, cuneate at the base and obtuse at the apex, glabrous, and subsessile. Flowers are grouped (2–15) in small axillary cymes; they have a corolla about 1 cm in diameter, with glabrous petals, lobes yellowish-green and purplish-brown on the inside. The fruits are narrowly conical follicles (0.6 × 6–10 cm), often paired and divergent. Seeds are dark (7 mm), with white woolly hairs (25–40 mm) that are easily dispersed by wind (anemochory) (Fig. 4a). Although it produces latex, the species is particularly appreciated by livestock, especially in the arid environments of North Africa; in Tunisia it also has medicinal value (Dghim et al. 2018).

Phytosociological investigation

The study was conducted through preliminary field surveys, initiated at the beginning of spring 2025, following the incidental discovery of *Periploca angustifolia* by one of the authors, documented by photographs posted on the web (Acta Plantarum 2025). Several excursions were subsequently carried out, aimed at spatially locating the population, monitoring the ecological-stationary features of the site, and conducting a preliminary survey of the flora and vegetation of the garrigue and low maquis it forms, as well as the role it plays within the landscape. Relevant literature was also consulted, along with herbarium investigations at the Herbaria of Palermo (PAL) and Catania (CAT), to define an updated distribution range of the species in the regional context, previously known only from some small circum-Sicilian islands (Linosa, Lampedusa, Pantelleria, Marettimo, Levanzo, Favignana), as well as the Maltese Islands, but not from Sicily itself.

For the geolithological characterization of the area, we mainly referred to Basilone (2018) and Lentini and Carbone (2014), as well as to various other publications

concerning the geomorphological and hydrothermal aspects of the site (Badino 2013; Verde 2021). Climatic data were derived from the nearby thermopluviometric station of Sciacca (La Mantia et al. 2025), considering the dataset from 1926–1985 analyzed by Duro et al. (1996).

The geobotanical investigation was expanded to include a phytosociological, synecological, and syndynamic characterization of the vegetation landscape, with particular attention to the shrub formations of *Periploca angustifolia*, primarily established in clearings within afforested areas along the southwestern slope of the hill. Twelve phytosociological relevés were performed in spring, on homogeneous patches of low maquis-garrigue vegetation, although partly discontinuous and of limited extent. The sampled surfaces ranged between 50–100 m², at elevations between 143 and 278 m a.s.l. The relevés were then assembled into a phytosociological table (Table 1). Moreover, it was used for further floristic–phytocoenotic comparison with other analogous communities dominated by *Periploca angustifolia* and/or related taxa (included in the alliance *Periplocion angustifoliae*), so far described for the islands of the Sicilian Channel. A synoptic table was subsequently compiled (Table 2), including 22 phytosociological tables and 184 relevés, with the aim of distinguishing groups of characteristic and differential elements within the various syntaxa, as well as the most representative companion species.

Species identification was primarily based on Pignatti et al. (2017–2019), whereas the taxonomic nomenclature followed the Portal to the Flora of Italy (2025), except for *Searsia tripartita* and *Lycium schweinfurthii* [according to Plants of the World Online (POWO 2025)]. The phytosociological nomenclature follows the criteria of the International Code of Phytosociological Nomenclature (ICPN: Theurillat et al. 2021), while syntaxonomic classification was based on Mucina et al. (2016) and Brullo et al. (2009, 2020).

The communities reported in the above synoptic table were further classified through cluster analysis, in which frequency classes were converted into numerical values according to van der Maarel scale (1979). Hierarchical cluster analysis was performed using the Bray–Curtis dissimilarity index. An agglomerative hierarchical clustering algorithm with Ward.D2 linkage was applied, minimizing within-group variance. NMDS (Non-metric Multidimensional Scaling) was used to graphically represent floristic relationships among the relevés in a two-dimensional space. Furthermore, a PERMANOVA (Permutational Multivariate Analysis of Variance) was carried out to test whether the studied associations differed significantly in composition, based on a Bray–Curtis distance matrix. Analyses were conducted with R software, version 4.5.1 (R Core 2025), using the “vegan” package (Oksanen et al. 2006) for NMDS, PERMANOVA, and Bray–Curtis distance matrix computation. The “cluster” and “stats” libraries were used for hierarchical clustering, while “ggplot2” and “matplotlib” (via Python) were employed for graphical visualization.




Figure 4. a. *Periploca angustifolia* at Mt. S. Calogero (Sciacca, Italy): twigs in fruiting and dissemination (anemochory); b. *Asparagus albi*-*Periplocetum* scrub, among the clearings of the afforestation; c. Follicles of *P. angustifolia* with nymphs of the host insect *Caenocoris nerii* (Lygaeidae); d. Other scrub areas along the firebreaks, on the south-eastern edge of Mount S. Calogero; e. Plant established in areas affected by fire; f. Scrub and grassland areas with *Hyparrhenia hirta*, colonizing the slopes denuded by the fires.

Results

The population of *Periploca angustifolia* on Mount S. Calogero (Sciacca) is mainly distributed across the lower and western sectors of the homonymous Nature Reserve (as well as SACITA040009), between the localities of Contrada Isabella, Cave di Pietra, and the easternmost part of the afforestation area (Fig. 3). The subarea occupies approximately 9 hectares, at elevations ranging from roughly 140 to 270 m a.s.l. The population is estimated at approximately 800–1000 individuals and appears to represent a relict, autochthonous nucleus well integrated into the environmental context of the area. The species grows in garrigue and low maquis vegetation, with small groups of residual individuals – including older plants in full fruiting – mostly found in clearings within the afforestation stands (Fig. 4b), as well as along the margins of firebreak strips (Fig. 4d). The population extends up to about 270 m a.s.l., even within rockfall protection barriers close to the cliffs (Fig. 4e, f). Here, the low maquis vegetation in which it occurs is common in areas stripped bare by fire and in clearings within the same afforestation stands, where it plays a clear constructive role in the recovery of more mature natural features of the local vegetation landscape.

Our surveys highlighted the presence of a particularly vigorous population of *Periploca*, with abundant fruiting and active anemochorous seed dispersal, as well as a rich presence of seedlings scattered throughout the area.

Distribution of the species on the islands of the Sicilian Channel

The first records of *Periploca angustifolia* in the Islands of the Sicilian Channel were provided by Gussone (1827) in the Aegadian Archipelago, where he reported it on the three largest islands (Favignana, Levanzo, and Marettimo). The same author later also indicated its presence on the islands of Pantelleria, Lampedusa, and Linosa (Gussone 1832). In the Maltese Archipelago, it was reported by Caruana Gatto (1890), under the name *Periploca laevigata*. Further mentions of the species were given by Riccobono (1906), within an account of his collections from Linosa and Lampedusa, where he also simultaneously indicated it in Sicily at “Modica and Girgenti.” However, for these latter localities, no herbarium exsiccata are known, nor subsequent confirmations by other authors have been provided.

Another specimen of the species collected by the same author must also be noted, preserved at the Palermo Herbarium (PAL63537!). It consists of three portions of twigs of the species, on the label of which—handwritten by Riccobono himself—appears the note: “Foresta Capaci – Cultivated. 4/25/1887, Riccobono.” This clearly indicates the origin of the sample, taken from a cultivated plant (in Contrada Carrubbella, where the toponym “Case Riccobono” still exists today).

In addition to this newly documented population on Mount S. Calogero, the current distribution of the species in the Sicilian Channel area—compiled from

bibliographic references and unpublished data—includes the following additional localities:

- Marettimo island: Punta Bassana (Gianguzzi et al. 2006);
- Favignana island: C.de Boschitto, Montagna Grossa o Grande, Punta Faraglione (Di Martino and Trapani 1967);
- Levanzo Island: Faraglione, Pietre Varate, Costa Dogana, Serra Alberello, Cala Fredda (Di Martino and Trapani 1968; Romano et al. 2006);
- Pantelleria island: C.da Fram, C.da Camilli, C.da Polacca, Punta Polacca, C.da Sataria (Brullo et al. 1977; Gianguzzi 1999);
- Lampedusa (frequent along the coastal belt and in the dry rocky slopes; Bartolo et al. 1988) and Linosa Islands (common in rocky habitat in various localities, as Panzarella, Arena Bianca, Montagna Rossa and Monte Vulcano; Brullo and Siracusa 1996; Malta (Mistra Valley, Salina Bay, Ghar Lapsi, Naxxar, Mnajdra near Qrendi, Victoria Lines, Ghar Lapsi), Gozo (Xlendi Valley, Triq Kalati Punic) and Comino islands (Brullo et al. 2020).

Phytosociology

Table 1 reports 12 relevés carried out in the low maquis community with presence/dominance of *Periploca angustifolia*, localized within the study area, which is here proposed as a new association, described below.

Asparago albi-Periplocetum angustifoliae Gianguzzi, La Mantia et Cambria ass. nov.

HOLOTYPE: Rel. 3, Table 1

Phytosociological data: Table 1 and Table 2 (column 1).

Diagnostic species: *Periploca angustifolia* (dominant), *Asparagus albus*, *Ampelodesmos mauritanicus*, *Carlina sicula* subsp. *sicula*, *Pistacia terebinthus*, *Chamaerops humilis*.

Short description: Primary or secondary low maquis, 2–3 m tall, more or less dense or open, occurring on xeric, rocky-stony slopes, with high frequency and sometimes dominance of *Periploca angustifolia*. It is associated with several elements of the alliance *Oleo-Ceratonion siliquae* and the order *Pistacio lentisci-Rhamnetalia alaterni* (e.g., *Chamaerops humilis*, *Olea europaea*, *Pistacia lentiscus*, *Asparagus albus*, *Teucrium fruticans*, *Stachys major*, as well as of the class *Quercetea ilicis* (e.g., *Ampelodesmos mauritanicus*, *Asparagus acutifolius*, *Pistacia terebinthus*, *Arisarum vulgare*, *Allium subhirsutum*).

Based on the data reported in Table 1 and in the synoptic table (Table 2), the specific species combination that characterizes the new association, here proposed under the name *Asparago albi-Periplocetum angustifoliae*, includes the following differential taxa (all more or less frequent within the community, and rare or virtually absent on the small islands of the Strait of Sicily):

1. *Ampelodesmos mauritanicus*, diagnostic of the order *Pistacio-Rhamnetalia alaterni* and the class *Quercetea ilicis* (Rivas-Martínez et al. 2001, 2002; Gianguzzi et al. 2016). This species is very common in the studied community yet virtually absent from other *Periploca angustifolia* scrub communities of the Strait of

Table 1. *Asparago albi-Periplocetum angustifoliae* ass. nov., Sciacca, Riserva M. S. Calogero, 13.05.2025 [rel. 1 (37.51505°N, 13.1007°E), 2 (37.51513°N, 13.10071°E), 3 (37.51491°N, 13.10235°E), 4 (37.51491°N, 13.10183°E), 5 (37.51569°N, 13.11134°E), 6 (37.51638°N, 13.10997°E), 7 (37.51667°N, 13.11056°E), 8 (37.51646°N, 13.10985°E)] and 19.05.2025 [rels. 9 (37.5168883°N, 13.1105639°E), 10 (37.5163093°N, 13.1111889°E), 11 (37.5155780°N, 13.1113448°E), 12 (37.5154315°N, 13.1116401°E)]. The values indicated in brackets concern species which were identified visually during field surveys performed during vegetative dormancy (dried leaf residues). The asterisk indicates the type relevé. The presence classes are as follows: I = 1–20%; II: 21–40%; III: 41–60%; IV: 61–80%; V = 81–100%.

No. of relevé	1	2	3*	4	5	6	7	8	9	10	11	12		F
Elevation (m a.s.l.)	143	144	146	150	227	239	278	271	261	226	201	190	P	R
Slope (°)	11	15	15	12	20	15	22	25	15	20	10	15	R	E
Aspect	S	S	S	S	S	S	S	S	S	S	S	S	E	Q.
Surface (m ²)	150	150	120	120	150	100	100	150	100	100	50	50	S	
Total cover (%)	95	100	95	95	95	100	85	90	90	90	90	90	E	C
Cover of the woody layer (%)	85	90	80	90	90	80	75	85	70	75	85	80	N	L
Cover of the herb layer (%)	35	35	45	60	65	30	75	50	75	70	70	75	C	A
Average height of the woody layer (m)	1.5	1.3	1.3	1.3	1.7	2.8	2.2	2.5	1.3	1.4	1.7	1.3	E	S
No. of taxa	31	34	35	31	25	24	30	23	24	23	22	24		S
Characteristic and differential association and alliance <i>Periplocion angustifoliae</i> (#)														
# <i>Periplocia angustifolia</i> (dom.)	2	3	3	3	3	3	3	3	2	2	3	2	12	V
<i>Asparagus albus</i>	2	1	2	2	2	3	2	2	1	1	1	+	12	V
<i>Ampelodesmos mauritanicus</i>	2	2	1	1	1	+	1	1	3	+	+	1	12	V
<i>Carlina sicula</i> subsp. <i>sicula</i>	1	+	1	+	.	1	1	.	1	.	1	1	9	IV
<i>Pistacia terebinthus</i>	.	1	1	.	2	1	1	.	.	+	1	.	7	III
Characteristic order <i>Pistacio-Rhamnetalia alaterni</i> and class <i>Quercetalia ilicis</i>														
<i>Chamaerops humilis</i>	3	3	2	2	2	2	1	1	1	1	3	2	12	V
<i>Olea europaea</i>	1	+	2	1	2	2	1	2	+	1	.	2	11	V
<i>Teucrium fruticans</i>	2	1	+	1	1	1	1	.	+	+	.	+	10	V
<i>Pistacia lentiscus</i>	1	2	2	2	.	+	+	.	+	1	1	.	9	IV
<i>Asparagus acutifolius</i>	1	2	1	+	+	.	.	+	+	.	.	1	8	IV
<i>Stachys major</i>	.	+	.	+	2	1	2	2	.	+	.	+	8	IV
<i>Ruta chalepensis</i>	1	+	1	.	.	2	1	.	1	.	+	+	8	IV
<i>Ceratonia siliqua</i> (cult.)	.	1	1	.	.	2	2	1	1	.	1	.	7	III
<i>Arisarum vulgare</i>	.	(+)	(+)	.	.	.	(+)	.	(+)	.	(+)	+	6	III
<i>Allium subhirsutum</i>	(+)	(+)	.	(+)	.	.	.	.	.	(+)	.	+	5	III
<i>Pinus halepensis</i> (cult.)	.	.	.	1	2	1	1	.	.	.	1	.	5	II
<i>Cytisus infestus</i> subsp. <i>infestus</i>	1	1	.	.	.	.	.	.	.	.	.	1	3	II
<i>Euphorbia dendroides</i>	.	.	.	.	.	.	.	2	.	3	.	.	2	I
<i>Myrtus communis</i> (cult.)	1	.	.	.	.	.	.	.	.	.	.	.	1	I
<i>Artemisia arborescens</i>	+	.	.	.	.	.	.	.	.	.	.	.	1	I
<i>Cyclamen repandum</i>	.	.	(+)	.	.	.	.	.	.	.	.	.	1	I
<i>Rubia peregrina</i>	.	.	.	+	.	.	.	.	.	.	.	.	1	I
<i>Osyris alba</i>	.	.	.	.	.	.	1	.	.	.	.	.	1	I
<i>Cercis siliquastrum</i> (cult.)	.	.	.	.	1	.	.	.	.	.	.	.	1	I
Other species														
<i>Hyparrhenia hirta</i>	2	1	4	2	4	+	5	2	1	2	2	2	12	V
<i>Asphodelus ramosus</i>	(+)	(1)	(+)	(1)	(1)	(1)	(1)	(+)	(+)	.	(+)	(1)	11	V
<i>Bituminaria bituminosa</i>	1	1	2	1	+	1	2	1	+	.	1	+	11	V
<i>Foeniculum vulgare</i> subsp. <i>vulgare</i>	1	+	1	+	2	+	2	1	1	+	+	.	11	V
<i>Stipellula capensis</i>	1	1	2	1	+	.	+	+	.	.	1	2	9	IV
<i>Andropogon distachyos</i>	1	1	+	+	.	.	+	+	.	.	+	+	8	IV
<i>Phagnalon saxatile</i>	1	+	+	+	.	.	1	+	+	.	.	+	8	IV
<i>Mercurialis annua</i>	.	+	+	.	+	1	+	+	.	+	.	+	8	IV
<i>Brachypodium distachyon</i>	+	+	1	.	.	+	.	+	+	.	+	.	7	III
<i>Spartium junceum</i> (cult.)	+	2	1	.	.	.	2	1	1	+	.	.	7	III
<i>Squilla pancration</i>	(1)	(1)	.	(1)	.	(1)	.	.	(+)	(1)	.	(+)	7	III
<i>Convolvulus altheoides</i>	+	.	+	.	+	.	+	+	.	+	.	.	6	III
<i>Pallenis spinosa</i>	1	+	1	+	+	.	.	.	.	.	.	1	6	III
<i>Eryngium campestre</i>	1	+	+	+	.	.	.	.	.	+	+	.	6	III
<i>Lathyrus clymenum</i>	.	.	+	.	+	+	+	+	.	.	+	.	6	III
<i>Dactylis glomerata</i> subsp. <i>hispanica</i>	1	.	.	+	1	.	+	.	.	.	.	1	5	II
<i>Acanthus mollis</i>	(+)	.	.	.	(+)	.	.	.	(+)	(+)	.	.	4	II
<i>Galactites tomentosus</i>	.	+	.	.	.	+	.	.	+	.	.	+	4	II
<i>Sonchus oleraceus</i>	.	.	+	.	+	.	.	.	+	+	.	.	4	II

No. of relevé	1	2	3*	4	5	6	7	8	9	10	11	12		F
Elevation (m a.s.l.)	143	144	146	150	227	239	278	271	261	226	201	190	P	R
Slope (°)	11	15	15	12	20	15	22	25	15	20	10	15	R	E
Aspect	S	S	S	S	S	S	S	S	S	S	S	S	E	Q.
Surface (m ²)	150	150	120	120	150	100	100	150	100	100	50	50	S	
Total cover (%)	95	100	95	95	95	100	85	90	90	90	90	90	E	C
Cover of the woody layer (%)	85	90	80	90	90	80	75	85	70	75	85	80	N	L
Cover of the herb layer (%)	35	35	45	60	65	30	75	50	75	70	70	75	C	A
Average height of the woody layer (m)	1.5	1.3	1.3	1.3	1.7	2.8	2.2	2.5	1.3	1.4	1.7	1.3	E	S
No. of taxa	31	34	35	31	25	24	30	23	24	23	22	24		S
<i>Fumana laevipes</i>	2	1	.	+	.	.	.	.	.	.	.	.	3	II
<i>Crupina crupinastrum</i>	+	.	.	+	.	.	.	.	.	.	+	.	3	I
<i>Convolvulus cantabrica</i>	.	+	.	.	.	+	.	.	1	.	.	.	3	I
<i>Sideritis romana</i>	.	+	.	.	.	.	+	.	.	+	.	.	3	I
<i>Echium parviflorum</i>	.	.	+	.	+	.	.	.	+	.	.	.	3	I
<i>Urospermum dalechampii</i>	.	.	+	+	.	.	.	.	.	.	.	.	2	I
<i>Lantana camara</i>	.	.	.	+	+	.	.	.	.	.	.	.	2	I
<i>Micromeria graeca</i>	.	.	.	.	.	.	1	+	.	.	.	.	2	I
<i>Allium ampeloprasum</i>	.	.	.	.	.	.	+	+	.	.	.	.	2	I
<i>Capparis spinosa</i> subsp. <i>rupestris</i>	.	1	.	.	.	.	.	.	.	.	.	.	1	I
<i>Allium commutatum</i>	.	.	r	.	.	.	.	.	.	.	.	.	1	I
<i>Ononis viscosa</i> subsp. <i>breviflora</i>	.	.	.	+	.	.	.	.	.	.	.	.	1	I
<i>Allium obtusiflorum</i>	.	.	.	+	.	.	.	.	.	.	.	.	1	I
<i>Serapias cordigera</i>	.	.	.	+	.	.	.	.	.	.	.	.	1	I
<i>Prunus dulcis</i> (cult.)	.	.	.	.	.	.	.	.	.	2	.	.	1	I
<i>Plantago afra</i>	.	.	.	.	.	.	.	.	.	.	+	.	1	I
<i>Silene colorata</i>	.	.	.	.	.	.	.	.	.	.	+	.	1	I

Sicily. In Sicily, however, it is also listed among the diagnostic elements of the following syntaxa:

- Ampelodesmo mauritanici-Quercetum ilicis* Gianguzzi et al. 2016, a *Quercus ilex* woodland found in inland areas, particularly in the Sicani Mountains (Gianguzzi et al. 2014a, 2014b);
 - Ampelodesmo mauritanici-Juniperetum turbinatae* (Gianguzzi et al. 2012), a relict maquis community dominated by *Juniperus turbinata*, also restricted to the Sicani Mountains (central-western Sicily; Gianguzzi et al. 2016);
 - the alliance *Avenulo cincinnatae-Ampelodesmion mauritanici* Minissale 1994 and the class *Lygeo sparti-Stipetea tenacissimae* (Brullo et al. 2010), which encompasses secondary grassland aspects, typically developed on rocky meso-Mediterranean lithosols, rather steep and degraded by frequent fires. In Sicily, several associations have also been described where this species is structurally dominant (Minissale 1994).
- Asparagus albus*, another diagnostic taxon of the order *Pistacio lentisci-Rhamnetalia alaterni*, frequent within the community but absent on the circum-Sicilian islands (Brullo and Siracusa 1996; Giardina et al. 2007; Gianguzzi et al. 2006; Romano et al. 2006).
 - Pistacia terebinthus*, a shrub relatively frequent in Sicily, also indicated as a characteristic element of the *Pistacio terebinthi-Celtidetum aetnensis* (Gianguzzi et al. 2014b), a maquis association dominated by *Celtis tournefortii*, with a fragmented distribution on Mt. Etna, the Nebrodi Mountains, the Sicani Mountains, and Rocca Busambra (Gianguzzi et al. 2014a).

4. *Carlina sicula* subsp. *sicula*, endemic to Sicily, linked to grassland, garrigue, and low scrub habitats, typical of rocky and xeric environments (Gianguzzi et al. 1996).

5. *Chamaerops humilis*, a diagnostic taxon of the order *Pistacio-Rhamnetalia alaterni*, dominant in some basiphilous scrub associations represented in Sicily (Brullo and Marcenò 1985; Brullo et al. 2009).

Substrates: Rocky calcareous and/or stony-debris slopes, edaphically poor or with shallow soils, occurring at the base of rock walls exposed to the South/South-East.

Bioclimate: Mediterranean oceanic pluviseasonal; lower thermomediterranean thermotype; lower dry ombrotype, tending toward semiarid conditions.

Syntaxonomic notes: The *Asparago albi-Periplocetum angustifoliae* ass. nov. is framed within the alliance *Periplocion angustifoliae* (order *Pistacio-Rhamnetalia alaterni*, class *Quercetea ilicis*). The distribution area of the alliance comprises the various islands of the Strait of Sicily [where 10 different associations have been described, summarized in the synoptic table (Table 2)], the small southern Aegean islets (Brullo and Guarino 2000), and extends across the Mediterranean fringe of Africa as far as southern Spain (Rivas-Martínez 1975).

Vegetation series: In its most typical form, the association *Asparago albi-Periplocetum angustifoliae* ass. nov. constitutes a garrigue or low maquis vegetation, acting as a transitional community linked to the following two xerophilous vegetation series: i) the *Ruto chalepensis-Oleo sylvestris sigmetum* is widespread on rocky carbonate substrates (Gianguzzi and Bazan 2020; Bazan et al.

[illegible]

Association table	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
No. relevés	2	8	14	10	5	7	14	8	6	13	7	5	8	4	4	19	6	5	7	8	14	10
Characteristic of class <i>Quercetea ilicis</i>																						
<i>Asparagus acutifolius</i>	IV	III	V	V	V	I	V	V	V	.	V	III	V	.	.	V	I	IV	V	IV	IV	.
<i>Rubia peregrina</i> subsp. <i>longifolia</i>	I	II	V	IV	.	.	II	.	.	III	.	II	V	3	.	IV	III	IV	V	III	IV	V
<i>Arisarum vulgare</i>	III	III	.	.	V	III	II	V	.	III	II	II	II	.	.	V	.	.	I	V	V	IV
<i>Ruta chalepensis</i>	IV	IV	.	I	IV	I	IV	V	I	.	I	IV	.	.	.	III	V	.	.	.	I	.
<i>Phillyrea latifolia</i>	.	.	IV	I	.	I	II	III	I	.	II	.	III	4	.	III	.	V	.	.	.	.
<i>Smilax aspera</i>	.	.	III	.	.	.	.	.	.	IV	.	I	II	1	.	II	.	.	.	.	II	II
<i>Asparagus aphyllus</i>	.	IV	.	V	.	.	.	.	.	V	.	.	V	.	.	.	.	V	.	III	.	V
<i>Lonicera implexa</i>	.	.	III	.	IV	.	.	.	.	III	.	I	.	3	.	.	.	.	.	.	.	V

2021) and its regressive stages are represented by grasslands dominated by *Hyparrhenia hirta* (*Hyparrhenietum hirta-pubescentis*) and by small therophytic grasslands with *Trachynia distachya* (class *Stipo-Trachynietea distachyae*); ii) the *Calicotomo infestae*-*Searsia tripartitae* *sigmetum* is typical of xeric stony-detrital substrates along eroded slopes. Its secondary aspects are again represented by the same vegetation types described for the previous series. The potential area of these two vegetation series is heavily disturbed by anthropogenic activities (e.g., afforestation, cultivation, more or less urbanized zones). In the upper part of Mount San Calogero, the same series are in catenal contact with the microgeosigmatum of cliffs, marked by sparse chasmophytic aspects referred to the alliance *Dianthion rupicolae* (class *Asplenietea trichomanis*) or to the association *Capparidetum rupestris* (class *Parietietea judaicae*); in the lower part, they likely extend almost to the coast (Fig. 5).

Synchorology: The association is endemic to this portion of the southern Sicilian coast near Sciacca, representing a geographical vicariant of association *Periploco-Euphorbietum dendroidis*, which is found on several small islands of the Strait of Sicily (Brullo and Marcenò 1985; Brullo et al. 2009; Gianguzzi et al. 2016).

Cluster analysis and ordination of the Sicilian communities

The dendrogram resulting from the cluster analysis performed on the associations listed in the synoptic table (Table 2) highlights the presence of two main clusters (Fig. 6). The first includes the associations *Asparago acutifolii-Ziziphietum loti*, *Ephedro fragilis-Pistacietum lentisci*, and *Asparago horridi-Retametum gussonei*, all occurring only in Sicily (where *Periploca angustifolia* is entirely absent), while the second brings together a more complex and diverse set of communities, subdivided into 10 groups.

Within the first group of this second cluster, two associations dominated by *Searsia tripartita* can be distinguished: the *Calicotomo infestae-Rhoetum tripartitae* (also restricted to Sicily and lacking *Periploca angustifolia*) and the *Periploco angustifoliae-Rhoetum tripartitae* (exclusive to the island of Linosa).

The remaining associations are distributed across the other 9 groups, as follows:

1. *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum* (Island of Marettimo; Aegadian Archipelago);
2. *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum* and *Periploco angustifoliae-Euphorbietum dendroidis* subass. *euphorbietosum papillaris* (both on the Island of Levanzo; Aegadian Archipelago);
3. *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum* (Island of Marettimo; Aegadian Archipelago);
4. *Asparago albi-Periplocetum angustifoliae* ass. nov. (southern Sicily);
5. *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum* and *Periploco angustifoliae-Euphorbietum dendroidis* subass. *euphorbietosum papillaris* (Island of Favignana; Aegadian Archipelago);
6. *Asparago aphylli-Tetraclinetum articulatae* and *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum* (both located in the Maltese Islands);
7. *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum* and *Periploco angustifoliae-Juniperetum turbinatae* subass. *brassicetosum insularis* (both on the Island of Pantelleria);
8. *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum* (Island of Linosa);
9. *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum* and *Periploco angustifoliae-Juniperetum turbinatae* subass. *typicum* (both on the Island of Lampedusa).

Overall, it emerges that the most frequent and widely recognized syntaxon across the different islands of the Strait of Sicily—the association *Periploco angustifoliae-Euphorbietum dendroidis*—shows a certain floristic heterogeneity, splitting into different subclusters depending on geographical provenance. All other communities, on the other hand, tend to separate rather clearly from one another, owing to their markedly different floristic composition.

As for the *Asparago albi-Periplocetum angustifoliae* ass. nov., it shows a certain autonomy compared to the other communities considered, displaying greater similarity with the aspects of *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum* found on Favignana Island (Aegadian Archipelago).

The NMDS analysis (Fig. 7) yields similar results, with the relevés of *Periploco angustifoliae-Euphorbietum dendroidis* broadly grouped into a rather wide and irregular




Figure 5. Scheme of the vegetation units co-occurring in the study area: (1) maquis with *Periploca angustifolia* (association *Asparagus albi-Periplocetum angustifoliae* ass. nov.); (2) perennial grassland with *Hyparrhenia hirta* (association *Hyparrhenietum hirtopubescentis*); (3) afforestation with *Pinus halepensis* and *Cupressus* spp. (4) chasmophytic community (alliance *Dianthion rupicolae*); (5) maquis with *Olea europaea* (association *Ruto chalepensis-Oleetum sylvestris*).

cluster, while the other communities remain fairly distant from each other.

Regarding the *Asparago albi-Periplocetum angustifoliae* ass. nov., also in this case it shows greater affinity with the relevés of *Periploco angustifoliae-Euphorbietum dendroidis* from the Aegadian Archipelago, geographically the closest.

Finally, the PERMANOVA analysis highlights significant differences in the floristic composition of the various communities investigated ($F = 3.52$; $p < 0.001$), confirming that the dissimilarities among the different associations are greater than their internal variability, and thus justifying the syntaxonomic treatment proposed.

Discussion

According to some authors (Brullo et al. 1995; Médail and Quézel 1997; Médail 2022), for *Periploca angustifolia* and other species previously confined to Northern Africa, evidence suggests that their northward expansion is linked to the Messinian Salinity Crisis (Late Miocene, ~5.96–5.33 Ma). This major geological event, caused by repeated closures of the Strait of Gibraltar, led to the

desiccation of the Mediterranean Sea (Hsü et al. 1973; Fauquette et al. 2006), creating widespread arid environments and new land connections between North Africa and parts of southern Europe, as Sicily, southern Italy, Sardinia, Corsica, and the Aegean region (Roveri et al. 2014; Médail 2017).

These conditions facilitated biogeographical exchanges and the dispersal of drought-adapted taxa (Krijgsman et al. 1999). Xerophilous and halophilous species expanded northwards from North Africa into newly emergent territories, contributing to the distinctive South-Mediterranean floristic elements still observed today (Brullo et al. 1995). In Sicily, such taxa occupy diverse dry and saline habitats, including dunes, salt marshes, riverbeds, badlands, gypsum outcrops, steppe grasslands, and garrigues.

Within this context, *Periploca angustifolia* represents a relict South-Mediterranean xerophyte, surviving today on Mount San Calogero under marginal ecological conditions. Its limited distribution likely reflects historical habitat loss and human disturbance, as well as intrinsic ecological constraints. The species serves as an indicator of infra- and thermo-Mediterranean semiarid to dry environments, although its narrow elevational range suggests vulnerability to even minor environmental changes.




Figure 6. Cluster analysis of the communities belonging to *Periplocion angustifoliae* occurring in Sicily and nearby islands: *Periplocion angustifoliae*: 1 – *Asparago albi-Periplocetum angustifoliae* ass. nov. 2 – *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum*, Linosa Island; 3 – *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum*, Pantelleria Island; 4 – *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum*, Lampedusa Island 5, 6 – *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum*, Marettimo Island; 7, 9 – *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum*, Levanzo Island; 8 – *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum*, Favignana Island; 10 – *Periploco angustifoliae-Euphorbietum dendroidis* subass. *typicum*, Malta islands; 11 – *Periploco angustifoliae-Euphorbietum dendroidis* subass. *euphorbietosum papillaris*, Levanzo Island; 12 – *Periploco angustifoliae-Euphorbietum dendroidis* subass. *euphorbietosum papillaris*, Favignana Island; 13 – *Periploco angustifoliae-Juniperetum turbinatae* subass. *typicum*, Lampedusa Island; 14 – *Periploco angustifoliae-Juniperetum turbinatae* subass. *brassicetosum insularis*, Pantelleria Island; 15 – *Periploco angustifoliae-Rhoetum tripartitae*, Linosa Island; 16 – *Calicotomo infestae-Rhoetum tripartitae*, Southeastern Sicily; 17 – *Calicotomo infestae-Rhoetum tripartitae*, M. Kronio (Sciacca); 18 – *Ephedro fragilis-Pistacietum lentisci*, Gela at manfria; 19 – *Ephedro fragilis-Pistacietum lentisci*, Ragusa at Cava Randello; 20 – *Asparago horridi-Retametum gussonei*, Gela at Manfria; 21 – *Asparago acutifolii-Ziziphietum loti*, Palermo at Allaura; 22 – *Asparago aphylli-Tetraclinetum articulatae*, Malta islands.




Figure 7. NMDS analysis of the communities belonging to the alliance *Periplocion angustifoliae* occurring in Sicily and nearby islands. Stress value = 0.182.

The new population grows in a low shrubland, which has been proposed as a new syntaxon (*Asparago albi-Periplocetum angustifoliae* ass. nov.), strictly endemic of this area, where it displays a clear relict character.

The new association is classified within the alliance *Periplocion angustifoliae* (order *Pistacio-Rhamnetalia alaterni*, class *Quercetea ilicis*). In agreement with various other authors (Brullo and Marcenò 1985; Mucina et al. 2016), it includes the “thermomediterranean semiarid deciduous relict low matorral of the coastal regions” of the south-western Mediterranean (Rivas-Martínez 1975), characterized by the frequency/dominance of thermo-xerophilous shrubs with a North African affinity. In addition to *Periploca angustifolia*, other characteristic shrubs occurring in Sicily are listed (all summer-deciduous and very rare), such as *Ziziphus lotus*, *Lycium intricatum*, *Lycium schweinfurthii* (Portal to the Flora of Italy 2025; POWO 2025), *Searsia tripartita*, *S. pentaphylla* (Brullo et al. 2009; La Mantia et al. 2025), as well as *Asparagus horridus*; on the island of Malta, *Tetraclinis articulata* is also present, and considered a characteristic species of the alliance.

The classification carried out through cluster analysis as well as NMDS highlighted its clear differentiation from the other associations of the alliance represented in this central-Mediterranean sector, apart from the *Periploco-Euphorbietum dendroidis*. Indeed, the new association is considered to act as vicariant with respect to the latter community, which is also marked by the frequency/dominance of *Periploca angustifolia*. However, the *Asparago albi-Periplocetum angustifoliae* ass. nov. is clearly differentiated by its floristic assemblage, characterized by the presence of several significant species, such as *Ampelodesmos mauritanicus*, *Carlina sicula*, *Pistacia terebinthus*,

Asparagus albus and *Chamaerops humilis*; the first three are completely absent, while the latter two are quite rare, in the other association of the small circum-Sicilian islands.

The distribution of the *Asparago albi-Periplocetum angustifoliae* ass. nov. is concentrated on the south-western slope of Mount San Calogero, where it had so far escaped botanical and naturalistic observations; this despite the fact that the site lies within a protected area (namely the Natural Reserve “Monte San Calogero”, and also the Natura 2000 site ITA040009), apparently well known from a floristic point of view. Equally surprising is that a similar coincidence had already occurred in this area with *Searsia tripartita*, another very interesting xerophytic shrub of pre-desert vegetation, although found in a nearby locality (La Mantia et al. 2025). This further highlights the floristic richness of Sicily, a major biodiversity hotspot of the Mediterranean (Cowling et al. 1996; Myers et al. 2000; Mittermeier et al. 2004; Testolin et al. 2024), where field explorations continue to provide new discoveries.

On some individuals of *P. angustifolia*, we observed a notable presence of *Caenocoris nerii* (Germar, 1847 (Heteroptera: Lygaeidae)) (Fig. 4c), an interesting oligophagous insect associated with Apocynaceae and also commonly found on *Nerium oleander* L. (Pericart 1998; Olivieri 2013), but relatively rare in Sicily (Pokharel et al. 2021). This insect exhibits a South Mediterranean and Palearctic distribution, with occurrences in certain European countries, and has previously been recorded on *Periploca angustifolia* in the islands of the Sicilian Channel (Pisciotta et al. 2011; Zito et al. 2011, 2013).

These species, which are subject to pressures and threats, including afforestation interventions carried out in the area. These were implemented in past decades—partly

even before the establishment of the natural reserve and the SAC—but it is evident that such interventions negatively interfered with native formations, almost completely erasing the relict patches of *Periploca angustifolia* shrublands (here classified to the association *Asparago albi-Periplocetum angustifoliae* ass. nov.) and the even smaller patches of *Searsia tripartita* (in turn referred to the association *Calicotomo-Rhoetum tripartitae*; Brullo et al. 2009; La Mantia et al. 2025). Both associations were confined by human activities—formerly agricultural, more recently forestry—confined to ever smaller spaces, and driven nearly to extinction. It is also evident that they were favored by the wildfires of recent years, which partially destroyed forest plantations.

Indeed, the new spaces opened up by fire have created clearings and grassland habitats more favorable to the *Asparago albi-Periplocetum angustifoliae* ass. nov., starting from the plants and small vegetation nuclei that had survived along hedges or at the margins of afforestations. This has led to an evident recovery of this shrub vegetation on the more xeric slopes, where the aforementioned community shows its climatic potential.

A comparable case to *Periploca angustifolia* is that of *Ptilostemon greuterii*, a Tertiary relict shrub endemic to northwestern Sicily and originally known from a single, very small population on Mount Inici (Raimondo and Domina 2006; Domina and Castellano 2012; Rossi et al. 2013; Gianguzzi et al. 2022). The species was already critically endangered due to anthropogenic conifer afforestation, which altered the local microclimate by increasing shade and inhibiting growth and regeneration. The situation appeared to worsen after a wildfire destroyed much of the pine plantation, but the event ultimately proved beneficial: thanks to a persistent soil seed bank, *P. greuterii* regenerated quickly once open, sunny, nutrient-rich conditions were restored. This led to the development of a new plant community, *Malvo parviflorae-Ptilostemetum greuterii*.

This case exemplifies how disturbance events, such as fire, can unexpectedly promote the resilience and regeneration of rare Mediterranean taxa when they restore natural habitat conditions, emphasizing the ecological importance of seed banks and habitat openness for relict and endemic species survival. These examples highlight the need for careful consideration, given the numerous afforestations efforts across Sicily and the Mediterranean—both historically and today—sometimes even in ecologically unique biotopes. Such interventions, often undertaken for hydrogeological protection or environmental restoration can disrupt native vegetation, alter ecosystem processes, and threaten biodiversity. The popular notion of “planting a million trees to save the planet” is insufficient if actions lack scientific guidance and fail to use species suited to local ecosystems. Inappropriate afforestations can cause habitat regression, degradation, and loss of biodiversity, placing rare communities and endemic species at particular risk due to their heightened sensitivity to ecological changes (Capotorti et al. 2020; Casavecchia et al. 2021). The goal should be the restoration of native vegetation, emphasizing the reintroduction of characteristic species often lost due to human impact. Poorly planned interventions can degrade ecosystems and

threaten sensitive species, particularly in protected areas and Natura 2000 sites, contradicting the conservation objectives of the Habitats Directive (92/43/EEC).

Conclusions

In addition to reporting a new stand of *Periploca angustifolia*, so far unrecorded for mainland Sicily, this study provided an updated distribution of this interesting species, while also clarifying its synecological and phytosociological features. In agreement with the opinion of several authors, its presence at this specific site in Sicily is probably due to the ancient territorial connections with North Africa during the Messinian, when many species reached Sicily.

Finally, the newly described community *Asparago albi-Periplocetum angustifoliae* illustrates how fire can enhance the regeneration of rare Mediterranean taxa in areas impacted by improper afforestations. Therefore, this remark should be considered when formulating the conservation measures planned for this community within the “Monte San Calogero (Sciacca)” Natura 2000 site (SAC ITA040009). This is of particular importance also because it pertains to the priority habitat “5220* Matorral arborescent with *Ziziphus*,” which is listed in Annex I of the Habitats Directive (92/43/EEC) due to its rarity and the need for strict protection.

Syntaxonomic scheme

Quercetea ilicis Br.-Bl. ex A. et O. Bolòs 1950

Pistacio lentisci-Rhamnetalia alaterni Rivas-Martínez 1975

Periplocion angustifoliae Rivas-Martínez 1975

Calicotomo infestae-Rhoetum tripartitae Bartolo, Brullo et Marcenò 1982

Asparago acutifolii-Ziziphietum loti Gianguzzi, Ilardi et Raimondo 1996

Ephedro fragilis-Pistacietum lentisci Brullo, Guarino et Ronsisvalle 2000

Asparago horridi-Retametum gussonei Brullo, Guarino et Ronsisvalle ex Brullo, Giusso, Siracusa et Spampinato 2001

Periploco angustifoliae-Juniperetum turbinatae Bartolo, Brullo, Minissale et Spampinato 1990

subass. *typicum*

subass. *brassicetosum insularis* Brullo, Gianguzzi, La Mantia et Siracusa 2009

Periploco angustifoliae-Rhoetum tripartitae Brullo, Gianguzzi, La Mantia et Siracusa 2009

Periploco angustifoliae-Euphorbietum dendroidis Brullo, Di Martino et Marcenò 1977

subass. *typicum*

subass. *euphorbietosum papillaris* Brullo, Gianguzzi, La Mantia et Siracusa 2009

Asparago aphylli-Tetraclinetum articulatae Brullo, Brullo C., Cambria et Giusso Del Galdo 2019

Asparago albi-Periplocetum angustifoliae Gianguzzi, La Mantia et Cambria ass. nov.

Author contributions

Conceptualization, L.G., A.L.M., S.C.; methodology, L.G., A.L.M., R.R., A.V., S.C.; investigation, L.G., A.L.M., R.R., A.V., S.C.; data curation, L.G., A.L.M., R.R., A.V., S.C.; data elaboration, L.G., A.L.M., S.C.; writing—original draft preparation, L.G., S.C.; writing—review and editing, L.G., A.L.M., R.R., A.V., S.C.; drawings, L.G. All authors have read and agreed to the final version of the manuscript.

Acknowledgements

This work was carried out with the financial support of the University of Palermo (FFR D13_001682, resp. L. Gianguzzi). The authors would especially like to thank Dr. Salvatore Carlino (Department of Rural and Territorial Development - Agrigento district, Sicilian Region) for the data pertaining afforestation and fire affecting the protected area, Prof. Haralabos Tsolakis (Dept. of Agricultural, Food and Forest Sciences, University of Palermo, Italy) and Prof. Attilio Carapezza for the identification of *Caenocoris nerii*.

References

- Abd El-Ghani MM, Ahmed DAEA (2025) Floristic diversity and ecological insights of the threatened liana *Periploca angustifolia* Labill. from the northwestern Mediterranean desert of Egypt: Conservation perspectives. *Journal for Nature Conservation* 85: 126879. <https://doi.org/10.1016/j.jnc.2025.126879>
- Acta Plantarum (2025) *Acta Plantarum*. <https://www.actaplantarum.org/> [Accessed on 1 May 2025]
- Ahmed DA (2009) Current situation of the flora and vegetation of the western Mediterranean desert of Egypt. Ph.D. Thesis, Tanta Univ., Tanta, 424 pp.
- Alcaraz F, Delgado MJ, Inocencio C, López Bernal E, Clemente M, ... Carreño E (2000) Estimación de la biodiversidad de la vegetación de la region de Murcia: asociaciones y hábitats. In: Calvo Sendin JF, Esteve Selma MA, López Bermúdez (Eds) *Biodiversidad, Contribución a su conocimiento y conservación en la region de Murcia*. Servicio de Publicaciones Universidad de Murcia, Murcia, 39–50.
- Badino G, De Vivo A, Prelli R, Sauro F, Savino G, ... Tedeschi R (2009) The Kronio project: a first note. In: *Proceedings of the 15th International Congress of Speleology; 2009 luglio 19–26; Kerville (TX): National Speleological Society, vol. 3, 1760–1763*.
- Badino G (2013) Micrometeorology of mt Cronio caves, Sicily. *Proceedings of the 16th International Congress of Speleology; Jul 21–28; Brno, Czech Republic*.
- Bartolo G, Brullo S, Minissale P, Spampinato G (1988) Flora e vegetazione dell'Isola di Lampedusa. *Bollettino dell'Accademia Gioenia di Scienze Naturali di Catania* 21(334): 119–225.
- Basilone L (2018) *Lithostratigraphy of Sicily*. Springer, Heidelberg, 349 pp. <https://doi.org/10.1007/978-3-319-73942-7>
- Bazan G, Bacchetta G, Bagella S, Bonari G, Bonini F, ... Gianguzzi L (2021) New national and regional Annex I Habitat records: from # 21 to #25. *Plant Sociology* 58(1): 167–178. <https://doi.org/10.3897/pls2020572/05>
- Benaissa A, Djebbar R (2018) *Searsia tripartita* (Ucria) Grande: A rock shrub of Ahaggar: a review. *Biocénose* 9(1): 9–19.
- Boulos L (2000) *Flora of Egypt: volume two (Geraniaceae - Boraginaceae)*. Al-Hadara Publishing, Cairo, 352 pp.
- Brullo C, Brullo S (2020) *Flora Endemica illustrata della Sicilia*. Laruffa, Reggio Calabria.
- Brullo C, Brullo S, Giusso del Galdo G, Guarino R, Minissale P, ... Spampinato G (2010) The *Lygeo-Stipetea* class in Sicily. *Annali di Botanica (Roma)* 4(0): 57–87.
- Brullo S, Gianguzzi L, La Mantia A, Siracusa G (2009) La classe *Quercetea ilicis* in Sicilia. *Bollettino dell'Accademia Gioenia di Scienze Naturali di Catania* 41(369): 1–77.
- Brullo S, Grillo M, Guglielmo A (1998) Considerazioni fitogeografiche sulla flora iblea. *Bollettino dell'Accademia Gioenia di Scienze Naturali di Catania* 29(352): 45–111.
- Brullo S, Guarino R (2000) Contribution to the knowledge of flora and vegetation of Khriis islet (Crete, SE Mediterranean sea). *Flora Mediterranea* 10: 265–282.
- Brullo S, Guarino R, Ronsisvalle G (2000) La vegetazione del litorale di Manfria, presso Gela (Sicilia), area soggetta a vincolo archeologico. *Archivio Geobotanico* 4: 91–107.
- Brullo S, Marcenò C (1983) Osservazioni fitosociologiche sull'isola di Marettimo (Arcipelago delle Egadi). *Bollettino dell'Accademia Gioenia di Scienze Naturali di Catania* 15(320): 201–228.
- Brullo S, Marcenò C (1985) Contributo alla conoscenza della classe *Quercetea ilicis* in Sicilia. *Notiziario della Società Italiana di Fitosociologia* 19(1): 183–229.
- Brullo S, Marcenò C, Di Martino A (1977) La vegetazione di Pantelleria (Studio fitosociologico). *Pubblicazioni dell'Istituto di Botanica dell'Università di Catania* 1–110.
- Brullo S, Minissale P, Spampinato G (1995) Considerazioni fitogeografiche sulla flora della Sicilia. *Ecologia Mediterranea* 21(1–2): 99–117. <https://doi.org/10.3406/ecmed.1995.1759>
- Brullo S, Siracusa G (1996) Studio fitosociologico dell'Isola di Linosa. *Documents phytosociologiques* 16: 123–174.
- Brullo S, Sciandrello S (2006) La vegetazione del bacino lacustre “Biviere di Gela” (Sicilia meridionale). *Fitosociologia*. 43(2): 21–40.
- Brullo S, Brullo C, Cambria S, Giusso del Galdo G (2020) The vegetation of the Maltese Islands. Springer, Cham. <https://doi.org/10.1007/978-3-030-34525-9>
- Brullo S, Giusso del Galdo G, Guarino R, Minissale P, Sciandrello S, Spampinato G (2012) Syntaxonomic survey of the class *Pegano harmalae-Salsoletea vermiculatae* Br.-Bl. & O. Bolos 1958 in Italy. *Plant Biosystems* 147(2): 472–492. <https://doi.org/10.1080/11263504.2012.717544>
- Cañadas EM, Fenu G, Peñas J, Lorite J, Mattana E, Bacchetta G (2014) Hotspots within hotspots: Endemic plant richness, environmental drivers, and implications for conservation. *Biological Conservation* 170: 282–291. <https://doi.org/10.1016/j.biocon.2013.12.007>
- Capotorti G, Zavatiero L, Copiz R, Del Vico E, Facioni L, ... Blasi C (2020) Implementation of IUCN criteria for the definition of the Red List of Ecosystems in Italy. *Plant Biosystems* 154(6): 1007–1011. <https://doi.org/10.1080/11263504.2020.1839806>
- Caruana Gatto A (1890) Di alcune piante rare indigene; nuove varietà e località delle medesime. *Il Naturalista Maltese* I: 7–8.
- Casavecchia S, Allegrezza M, Angiolini C, Biondi E, Bonini F, ... Ciaschetti G (2021) Proposals for improvement of Annex I of Directive 92/43/EEC: Central Italy. *Plant Sociology* 58(2): 99–118. <https://doi.org/10.3897/pls2021582/08>

- Chaieb M, Boukhris M (1998) Flore succincte et illustrée des zones arides et sahariennes de Tunisie. Sfax: Association pour la Protection de la Nature et de l'Environnement, Sfax, Tunisie, 290 pp.
- Commissione Grotte Boegan and Associazione Geografica La Venta (2013) Stufe di San Calogero, una sfida millenaria. *Speleologia*. 68: 76–78. https://www.speleo.it/site/images/speleologia/Speleologia_68_Giu_2013_en.pdf
- Cowling RM, Rundel PW, Lamont BB, Arroyo MK, Arianoutsou M (1996) Plant diversity in Mediterranean-climate regions. *Trends in Ecology and Evolution* 11: 362–366. [https://doi.org/10.1016/0169-5347\(96\)10044-6](https://doi.org/10.1016/0169-5347(96)10044-6)
- Dghim F, Abdellaoui R, Boukhris M, Neffati M, Chaieb M (2018) Physiological and biochemical changes in *Periploca angustifolia* plants under withholding irrigation and rewatering conditions. *South African Journal of Botany* 114: 241–249. <https://doi.org/10.1016/j.sajb.2017.11.007>
- Di Martino A, Trapani S (1967) Flora e vegetazione delle Isole di Favignana e Levanzo nell'Arcipelago delle Egadi. I. Favignana. *Lavori Istituto Botanico del Giardino Coloniale di Palermo* 22: 122–228.
- Di Martino A, Trapani S (1968) Flora dell'Isola di Favignana e Levanzo nell'Arcipelago delle Egadi. 2. Levanzo. *Lavori Istituto Botanico del Giardino Coloniale di Palermo* 23: 37–132.
- Di Piazza S, Isaia M, Vizzini A, Badino G, Voyron S, Zotti M (2017) First mycological assessment in hydrothermal caves of Monte Kronio (Sicily, southern Italy). *Webbia* 72(2): 277–285. <https://doi.org/10.1080/00837792.2017.1347368>
- Domina G, Castellano G (2012) *Ptilostemon greuteri* Raimondo et Domina. In: Schede per una Lista Rossa della Flora vascolare e crittogamica Italiana. *Informatore Botanico Italiano* 44(2): 468–469.
- Duro A, Piccione V, Scalia C, Zampino D (1996) Precipitazioni e temperature medie mensili in Sicilia relative al sessantennio 1926–1985. In: Piccione V, Antonelli C (Eds) *Proceedings of the 5th Workshop del Progetto Strategico CNR Clima Ambiente e Territorio nel Mezzogiorno*, Amalfi April 28–30 1993. Amalfi: CNR eds, 17–103.
- El-Hadidi MN, Hosni HA (2000) *Flora Aegyptiaca*: Volume 1, Part 1. The Palm Press, Cairo, 187 pp.
- Ennajeth MM, Cochard HH, Khemira HH (2010) Re-introduction success of an autochthonous plant species, *Periploca angustifolia*, in the Natural Reserve of Oued Dekouk, Tunisia. *Spanish Journal of Agricultural Research* 8(4): 1005–1011. <https://doi.org/10.5424/sjar/2010084-1205>
- Fauquette S, Suc JP, Bertini A, Popescu SM, Warny S, ... Clauzon G (2006) How much did climate force the Messinian salinity crisis? Quantified climatic conditions from pollen records in the Mediterranean region. *Palaeogeography, Palaeoclimatology, Palaeoecology* 238(1–4): 281–301. <https://doi.org/10.1016/j.palaeo.2006.03.029>
- Ferranti L, Pepe F, Barreca G, Meccariello M, Monaco C (2019) Multi-temporal tectonic evolution of Capo Granitola and Sciacca foreland transcurrent faults (Sicily Channel). *Tectonophysics* 765: 187–204. <https://doi.org/10.1016/j.tecto.2019.05.002>
- Floriani E (1979) Studi idrogeologici sul bacino termale di Sciacca. *Datt. Inedito*, A.T.S. Morciano, 81 pp.
- Forti P (1991) Processi ipercarsici e speleogenesi. *Speleologia* 24: 42–46.
- Gamal E, Khderi G, Morsy A, Ali M, Hashim A, Saleh H (2020) GIS based modelling to aid conservation of two endangered plant species (*Ebenus armitagei* and *Periploca angustifolia*) at Wadi Al-Afreet, Egypt. *Remote Sensing Applications: Society and Environment* 19: 100336. <https://doi.org/10.1016/j.rsase.2020.100336>
- Geraci A, Raimondo FM, Troia A (2009) Genetic diversity and local population structure in *Ambrosina bassii* (Araceae, Ambrosineae), a Mediterranean relict species. *Biochemical Systematics and Ecology* 37: 737–746. <https://doi.org/10.1016/j.bse.2009.12.002>
- Gianguzzi L (1999) Vegetazione e bioclimatologia dell'Isola di Pantelleria (Canale di Sicilia). *Braun-Blanquetia* 22: 3–70.
- Gianguzzi L, Bazan G (2020) A phytosociological analysis of the *Olea europaea* L. var. *sylvestris* (Mill.) Lehr. forests in Sicily. *Plant Biosystems* 154(5): 705–725. <https://doi.org/10.1080/11263504.2019.1681532>
- Gianguzzi L, Caldarella O, Pasta S (2022) A new association of relict maquis with *Ptilostemon greuteri* (Oleo-Ceratonion, *Quercetea ilicis*), located in a circumscribed area of north-western Sicily. *Plant Sociology* 59(1): 67–83. <https://doi.org/10.3897/pls2022591/06>
- Gianguzzi L, Cusimano D, Cuttonaro L, Gianguzzi G, Romano S (2014a) Distribution, ecology and conservation survey on the *Celtis tournefortii* subsp. *aetnensis* (Celtidaceae-Cannabaceae) populations in Sicily. *Webbia* 69(2): 325–334. <https://doi.org/10.1080/00837792.2014.971586>
- Gianguzzi L, Cusimano D, Romano S (2014b) Phytosociological characterization of the *Celtis tournefortii* subsp. *aetnensis* microwoods in Sicily. *Plant Sociology* 51(2): 17–28. <https://doi.org/10.7338/pls2014512/02>
- Gianguzzi L, Ilardi V, Caldarella O, Cusimano D, Cuttonaro P, Romano S (2012) Phytosociological characterization of the *Juniperus phoenicea* L. subsp. *turbinata* (Guss.) Nyman formations in the Italo-Tyrrhenian Province (Mediterranean Region). *Plant Sociology* 49(2): 3–28.
- Gianguzzi L, Ilardi V, Raimondo FM (1996) La vegetazione del promontorio di Monte Pellegrino (Palermo). *Quaderni di Botanica Ambientale e Applicata* 4[1993]: 79–137.
- Gianguzzi L, Papini F, Cusimano D (2016) Phytosociological survey vegetation map of Sicily (Mediterranean region). *Journal of Maps* 12(5): 845–851. <https://doi.org/10.1080/17445647.2015.1094969>
- Gianguzzi L, Scuderi L, Pasta S (2006) La flora vascolare dell'Isola di Marettimo (Arcipelago delle Egadi, Canale di Sicilia): Aggiornamento ed analisi fitogeografica. *Webbia* 61(2): 359–402. <https://doi.org/10.1080/00837792.2006.10670810>
- Giardina G, Raimondo FM, Spadaro V (2007) A catalogue of plants growing in Sicily. *Bocconea* 20: 5–582.
- Guidi P, Prelli R, Torelli L (2014) Le stufe di San Calogero. In *Sicilia un enigma affascinante che gli speleologi cercano di svelare*. *Montagne* 360(2): 56–58.
- Guidi P, Verde G (2001) Il fenomeno carsico del monte Cronio. *Saggio Bibliografico. Atti e Memorie CGEB* 24: 1–150.
- Gussone G (1827) *Florae Siculae Prodromus sive plantarum in Siciliae ulteriori nascentium enumeratio secundum systema Linneanum disposita*. Vol. 1. Neapoli, ex Regia Typographia, vii + 592 + xi pp. <https://doi.org/10.5962/bhl.title.6700>
- Gussone G (1828) *Florae Siculae Prodromus*. II. – pp. 1–581. Regia Tipografia, Napoli.
- Gussone G (1832) *Supplementum ad Florae Siculae Prodromum, quoad et specimen Florae Insularum Siciliane Ulteriori Adjacentium*. I. – pag. 1–168. Regia Tipografia, Napoli.
- Gussone G (1834) *Supplementum ad Florae Siculae Prodromum, quoad et specimen Florae Insularum Siciliane Ulteriori Adjacentium*. II. – pp. 171–208. Regia Tipografia, Napoli.
- Gussone G (1843) *Florae Siculae Synopsis exhibens plantas vasculares in Sicilia insulisque adjacentibus hucusque detectas secundum systema Linneanum dispositas*. Vol. 1. Neapoli, Typ. Tramater, v + 575 pp. <https://doi.org/10.5962/bhl.title.50455>
- Hsü KJ, Ryan WB, Cita MB (1973) Late Miocene desiccation of the Mediterranean. *Nature* 242: 240–244. <https://doi.org/10.1038/242240a0>
- Ilardi V, Troia A, Geraci A (2020) *Brassica tardarae* (Brassicaceae), a new species from a noteworthy biotope of South-western Sicily (Italy). *Plants* 9(8): 947. <https://doi.org/10.3390/plants9080947>

- Ionta GM, Judd WS (2007) Phylogenetic relationships in Periplocoideae (Apocynaceae s.l.) and insights into the origin of pollinia. *Annals of the Missouri Botanical Garden* 94: 360–375. [https://doi.org/10.3417/0026-6493\(2007\)94\[360:PRIPAS\]2.0.CO;2](https://doi.org/10.3417/0026-6493(2007)94[360:PRIPAS]2.0.CO;2)
- Izquierdo JJ, Botella JP (2005) El género *Periploca* en la Comunidad Valenciana. *Flora Montiberica* 31: 39–43.
- Krijgsman W, Hilgen FJ, Raffi I, Sierro FJ, Wilson DS (1999) Chronology, causes and progression of the Messinian salinity crisis. *Nature* 400(6745): 652–655. <https://doi.org/10.1038/23231>
- La Mantia A, Rocca R, Gianguzzi L (2025) Rediscovery of a relict *Searsia tripartita* maquis in Southwestern Sicily (Italy) with data on habitat 5220. *Vegetation Ecology and Diversity* 62: 1–14. <https://doi.org/10.3897/ved.140946>
- La Mantia A, Scuderi L (2021) Present and past distribution of *Ziziphus lotus* (L.) Lam. subsp. *lotus* (Rhamnaceae) in Sicily. *Il Naturalista Siciliano series 4*, 45(1–2): 7–28.
- Lentini F, Carbone S (2014) Geology of Sicily. Descriptive Memories of the Geological Map of Italy. Volume XCV. ISPRA-Servizio Geologico d'Italia.
- Lojacono-Pojero M (1888–1889) Flora Sicula o descrizione delle piante spontanee o indigenate in Sicilia. Vol. 1(1) (Polypetalae-Thalamiflorae). Palermo, Stab. Tipografico Virzi, 234 + XIV pp.
- Lojacono-Pojero M (1891) Flora Sicula o descrizione delle piante spontanee o indigenate in Sicilia. Vol. 1(2) (Polypetalae-Calyciflorae). Tip. dello Statuto, Palermo, 311 + XVI pp.
- Lojacono-Pojero M (1902–1903) Flora Sicula o descrizione delle piante spontanee o indigenate in Sicilia. Vol. 2(1) (Gamopetalae Calyciflorae). Tip. dello Statuto, Palermo, 240 + XIV pp.
- Lojacono-Pojero M (1904–1907) Flora Sicula o descrizione delle piante spontanee o indigenate in Sicilia. Vol. 2(2) (Corolliflorae-Monochlamydeae, Gymnospermae). Tipo-Litografia S. Bizzarrilli, Palermo, 428 pp.
- Lojacono-Pojero M (1906) Notizie. Addenda et emendanda ad Floram Siculam. *Malpighia* (Genova) 20: 37–48, 95–119, 180–218, 290–300.
- Lojacono-Pojero M (1908–1909) Flora Sicula o descrizione delle piante spontanee o indigenate in Sicilia. Vol. 3 (Monocotyledones, Cryptogamae). Scuola Tipograf. “Boccone del Povero”, Palermo, 448 + XVI pp.
- Médail F (2017) The specific vulnerability of plant biodiversity and vegetation on Mediterranean islands in the face of global change. *Regional Environmental Change* 17(6): 1775–1790. <https://doi.org/10.1007/s10113-017-1123-7>
- Médail F (2022) Plant biogeography and vegetation patterns of the Mediterranean islands. *The Botanical Review* 88(1): 63–129. <https://doi.org/10.1007/s12229-021-09245-3>
- Médail F, Quézel P (1997) Hot-spots analysis for conservation of plant biodiversity in the Mediterranean basin. *Annals of the Missouri Botanical Garden* 84: 112–127. <https://doi.org/10.2307/2399957>
- Mendoza-Fernández AJ, Martínez-Hernández F, Salmerón-Sánchez E, Pérez-García FJ, Teruel B, ... Mota JF (2020) The relict ecosystem of *Maytenus senegalensis* subsp. *europaea* in an agricultural landscape: Past, present and future scenarios. *Land* 10(1): 1. <https://doi.org/10.3390/land10010001>
- Minissale P (1994) Studio fitosociologico delle praterie ad Ampelodesmos mauritanicus della Sicilia. *Colloques Phytosociologiques* 21: 615–652.
- Mittermeier R, Robles GP, Hoffmann M, Pilgrim J, Brooks T, ... da Fonseca G (2004) Hotspots revisited: Earth's biologically richest and most endangered ecoregions. CEMEX, Mexico City.
- Mota J, Cabello J, Cueto M, Gómez F, Giménez E, Peñas J (1997) Datos sobre la vegetación del sureste de Almería (Desiertos de Tabernas, Karst en Yesos de Sorbas y Cabo de Gata). Servicio Publicaciones Universidad de Almería, Almería, 130 pp.
- Mucina L, Bültmann H, Dierßen K, Theurillat J-P, Raus T, ... Tichý L (2016) Vegetation of Europe: Hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities. *Applied Vegetation Science* 19(S1): 3–264. <https://doi.org/10.1111/avsc.12257>
- Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* 403: 853–855. <https://doi.org/10.1038/35002501>
- Noumi Z, Dhaou SO, Chaieb M (2010) Seed germination characteristics of *Periploca angustifolia* Labill. and *Searsia tripartita* (Ucria) Grande: effects of temperature, salinity and water stress. *Acta Botanica Gallica* 157(2): 317–327. <https://doi.org/10.1080/12538078.2010.10516209>
- Noumi Z, Ouled Dhaou S, Chaieb M (2013) Seed germination characteristics of *Periploca angustifolia* Labill. and *Rhus tripartita* (Ucria) Grande: effects of temperature, salinity, and water stress. *Acta Botanica Gallica* 157: 317–327. <https://doi.org/10.1080/12538078.2010.10516209>
- Oksanen J, Kindt R, Legendre P, O'Hara RB (2006) *Vegan Community Ecology Package* Version 1.17-4.
- Olivieri N (2013) Segnalazioni Faunistiche Italiane - 570 - *Tropidothorax sternalis sternalis* (Dallas, 1852) (Heteroptera Lygaeidae). *Bollettino Della Società Entomologica Italiana* 145(3): IX–XIII.
- ORBD [Osservatorio Regionale Biodiversità Siciliana] (2025a) Aree naturali protette: Riserva Naturale Monte San Calogero (M. Kronio). <https://orbs.regione.sicilia.it/aree-protette/riserve-naturali-siciliane/152-riserva-naturale-monte-san-calogero-monte-kronio.html> [Accessed on 2025, July 16]
- ORBD [Osservatorio Regionale Biodiversità Siciliana] (2025b) – Rete Natura 2000 Regione Siciliana. <https://orbs.regione.sicilia.it/aree-protette/rete-natura2000.html> [Accessed on 2025, July 16]
- ORBD [Osservatorio Regionale Biodiversità Siciliana] (2025c) – Sistema di cavità carsico-ipogeniche delle Stufe di San Calogero (Monte Kronio). <https://orbs.regione.sicilia.it/aree-protette/geositi/205-sistema-di-cavita-carsico-ipogeniche-delle-stufe-di-san-calogero-monte-kronio.html> [Accessed on 2025, July 16]
- Ottone D (1978) *Brassica villosa* Biv. endemica del settore floristico siciliano rinvenuta a monte Cronio sulla costa sud-occidentale dell'isola. *Il Naturalista Siciliano, serie 4*, 2(3–4): 153–158.
- Pericart J (1998) Hemipteres Lygaeidae euro-mediterraneens. Generalites. Systematique: premiere partie - Faune de France. 84/A. Federation Francaise des Societes des Sciences Naturelles, Paris, Vol. I, 468 pp.
- Pignatti S, Guarino R, La Rosa M (2017–2019) *Flora d'Italia*. 2nd edition. Edagricole - Business Media, Bologna. Milano, 4 vols.
- Pisciotta S, Raspi A, Sajeva M (2011) First records of pollinators of two co-occurring Mediterranean Apocynaceae. *Plant Biosystems* 145: 141–149. <https://doi.org/10.1080/11263504.2010.540779>
- Pokharel P, Steppuhn A, Petschenka G (2021) Dietary cardenolides enhance growth and change the direction of the fecundity-longevity trade-off in milkweed bugs (Heteroptera: Lygaeinae). *Ecology and evolution* 11(24): 18042–18054. <https://doi.org/10.1002/ece3.8402>
- Portal to the Flora of Italy (2025) Portal to the Flora of Italy 2024.2. <http://dryades.units.it/floritaly/> [Accessed on 2025, August 2]
- POWO [Plants of the World Online] (2025) Plants of the World Online. Royal Botanic Garden Kew. <https://powo.science.kew.org/> [Accessed on 2025, August 2]
- Quézel P (1983) Flora and vegetation of North Africa, their significance according to the origin, evolution and migrations of flora and past vegetation structures. *Bothalia* 14: 411–416. <https://doi.org/10.4102/abc.v14i3/4.1186>

- Quézel P (1985) Definition of the Mediterranean region and the origin of its flora. In: Gomez-Campo (Ed.) Plant conservation in the Mediterranean area. Junk, Dordrecht, 9–24.
- R Core Team (2025) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. <https://www.r-project.org/>
- Raimondo FM, Domina G (2006) *Ptilostemon greuteri* (Compositae), a new species from Sicily. Willdenowia 36(1): 169–175. <https://doi.org/10.3372/wi.36.36114>
- Raimondo FM, Mazzola P, Ottonello D (1991) On the taxonomy and distribution of *Brassica* sect. *Brassica* (Cruciferae) in Sicily. Flora Mediterranea 1(1): 63–86.
- Riccobono A (1906) Le piante della flora siciliana più acconce all'ornamento dei giardini. Priulla, Palermo vol. I, 138 pp.
- Rivas-Martínez S (1975) La vegetación de la clase *Quercetea ilicis* en España y Portugal. Anales del Instituto Botánico Cavanilles 31(2): 205–259.
- Rivas-Martínez S, Díaz TE, Fernández-González F, Izco J, Lousa M, Penas A (2002) Vascular plant communities of Spain and Portugal. Addenda to the syntaxonomical checklist of 2001. Part. II. Itinera Geobotanica 15: 433–922.
- Rivas-Martínez S, Fernández-González F, Loidi J, Lousa M, Penas A (2001) Syntaxonomical checklist of vascular plant communities of Spain and Portugal to association level. Itinera Geobotanica 14(2): 5–341.
- Rivas-Martínez S, Penas A, Díaz TE (2004) Bioclimatic map of Europe, bioclimates, cartographic service. University of León, Spain.
- Rivieccio G, Aleffi M, Angiolini C, Bagella S, Bazan G, ... Gigante D (2021) New national and regional Annex I Habitat records: from #26 to #36. Plant Sociology 58(2): 77–98. <https://doi.org/10.3897/pls2021582/07>
- Romano S, Tobia G, Gianguzzi L (2006) Rassegna della flora vascolare dell'Isola di Levanzo (Arcipelago delle Egadi, Canale di Sicilia). Informatore Botanico Italiano 38(2): 481–502.
- Rossi G, Montagnani C, Gargano D, Peruzzi L, Abeli T, ... Orsenigo S [Eds] (2013) Lista Rossa della Flora Italiana. 1. Policy Species e altre specie minacciate. Comitato Italiano IUCN e Ministero dell'Ambiente e della Tutela del Territorio e del Mare.
- Roveri M, Flecker R, Krijgsman W, Lofi J, Lugli S, ... Govers R (2014) The Messinian Salinity Crisis: past and future of a great challenge for marine sciences. Marine Geology 352: 25–58. <https://doi.org/10.1016/j.margeo.2014.02.002>
- Salama FM, Abd El-Ghani MM, El-Naggar SM, Baayo Kh (2005) Vegetation structure and environmental gradients in the Sallum area, Egypt. Ecologia Mediterranea 31: 15–32. <https://doi.org/10.3406/ec-med.2005.1476>
- Smith GB, Eberhard SM, Perina G, Finston T (2012) New species of short range endemic troglobitic silverfish (Zygentoma: Nicoletiidae) from subterranean habitats in Western Australia's semi-arid Pilbara region. Records of the Western Australian Museum 27(2): 101–116. [https://doi.org/10.18195/issn.0312-3162.27\(2\).2012.101-116](https://doi.org/10.18195/issn.0312-3162.27(2).2012.101-116)
- Souza Maysa FVR, Ferreira RL (2011) A new troglobitic *Eukoenenia* (Palpigradi: Eukoeneiidae) from Brazil. Journal of Arachnology 39(1): 185–188. <https://doi.org/10.1636/ha10-43.1>
- Taifour H, El-Oqlah A, Ghazanfar SA (2017) The plants of Jordan: an annotated checklist. Royal Botanic Gardens, Kew, 242 pp.
- Testolin R, Attorre F, Bruzzaniti V, Guarino R, Jiménez-Alfaro B, ... Chiarucci A (2024) Plant species richness hotspots and related drivers across spatial scales in small Mediterranean islands. Journal of Systematics and Evolution 62(2): 242–256. <https://doi.org/10.1111/jse.13034>
- Theurillat JP, Willner W, Fernández-González F, Bültmann H, Čarni A, ... Weber H (2021) International Code of Phytosociological Nomenclature. 4th edn. Applied Vegetation Science 24: e12491. <https://doi.org/10.1111/avsc.12491>
- Troia A, Raimondo FM, Geraci A (2012) Does genetic population structure of *Ambrosina bassii* L. (Araceae, Ambrosineae) attest a post-Messinian land-bridge between Sicily and Africa? Flora 207: 646–653. <https://doi.org/10.1016/j.flora.2012.06.017>
- Turrisi RE, Galletti I, Ilardi V (2002) Contributo alla conoscenza della vegetazione di Cava Randello. Quaderni di Botanica ambientale applicata 12(2001): 117–130.
- van der Maarel E (1979) Transformation of Cover–Abundance Values in Phytosociology and Its Effects on Community Similarity. Vegetatio 39: 97–114. <https://doi.org/10.1007/BF00052021>
- Venter HJT (1997) A revision of *Periploca* (Periplocaceae). South Africa Journal of Botany 63: 123–128. [https://doi.org/10.1016/S0254-6299\(15\)30723-7](https://doi.org/10.1016/S0254-6299(15)30723-7)
- Verde G (2021) Cronio The geo-cultural site of Monte San Calogero in Sciacca (Sicily). Atti e Memorie della Commissione Grotte “E.Boegan” 50: 3–21.
- Zito P, Sajeva M, Bruno M, Maggio A, Rosselli S, Senatore F, Formisano C (2011) Essential oil composition of the fruits of *Periploca laevigata* Aiton subsp. *angustifolia* (Labill.) Markgraf (Apocynaceae – Periplocoideae). Natural Product Research 25: 1339–1346. <https://doi.org/10.1080/14786419.2010.535157>
- Zito P, Sajeva M, Bruno M, Rosselli S, Maggio A, Senatore F (2013) Essential oils composition of *Periploca laevigata* Aiton subsp. *angustifolia* (Labill.) Markgraf (Apocynaceae–Periplocoideae). Natural Product Research 27(3): 255–265. <https://doi.org/10.1080/14786419.2012.671319>

Supplementary material 1

Taxonomic treatment and selected specimens examined

Authors: Lorenzo Gianguzzi, Antonino La Mantia, Riccardo Rocca, Andrea Volpe, Salvatore Cambria

Data type: docx

Explanation note: The supplementary file includes the taxonomic treatment and selected specimens examined.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/ved.175232.suppl1>