

Doxorubicin anti-tumor mechanisms include Hsp60 post-translational modifications leading to the Hsp60/p53 complex dissociation and instauration of replicative senescence

Antonella Marino Gammazza ¹ - Claudia Campanella ¹ - Rosario Barone ¹ - Celeste Caruso Bavisotto ¹ - Magdalena Gorska ² - Michal Wozniak ² - Francesco Carini ¹ - Francesco Cappello ¹ - Antonella D'Anneo ³ - Marianna Lauricella ⁴ - Giovanni Zummo ¹ - Everly Conway De Macario ⁵ - Alberto J.L. Macario ⁵ - **Valentina Di Felice** ¹

¹ Department of Experimental Biomedicine and Clinical Neurosciences, University of Palermo, Palermo, Italy - ² Department of Medical Chemistry, Medical University of Gdansk, Gdansk, Poland - ³ Department of Biological, Chemical and Pharmaceutical Sciences and Technologies, Laboratory of Biochemistry, University of Palermo, Palermo, Italy - ⁴ Department of Experimental Biomedicine and Clinical Neurosciences, Laboratory of Biochemistry, University of Palermo, Palermo, Italy - ⁵ Department of Microbiology and Immunology, School of Medicine, University of Maryland at Baltimore; and Imet, Columbus Center, Baltimore, Md, USA

Hsp60 is a pro-carcinogenic chaperonin in certain tumor types by interfering with apoptosis and with tumor cell death. In these tumors, it is not known whether or not doxorubicin anti-tumor effects include a blockage of the pro-carcinogenic action of this protein. We used the human lung mucoepidermoid cell line NCI-H292 and different doses of doxorubicin to measure cell viability, cell cycle progression, cell senescence indicators, Hsp60 levels and its post-translational modifications as well as the release of the chaperonin into the extracellular environment. Cell viability was reduced in relation to doxorubicin dose and this was paralleled by the appearance of cell senescence markers. Concomitantly, intracellular Hsp60 levels decreased while its acetylation levels increased. The data suggest that Hsp60 acetylation may interfere with the formation of the Hsp60/p53 complex and/or promote its dissociation, both causing an increase in the levels of free p53, which can then activate cell senescence. On the other hand, acetylated Hsp60 is ubiquitinated and degraded and, thus, the anti-apoptotic effect of Hsp60 is impaired with subsequent tumor cell death.

References

- [1] Campanella C, D'Anneo A, Gammazza AM, Bavisotto CC, Barone R, Emanuele S, Lo Cascio F, Mocciano E, Fais S, De Macario EC, Macario AJ, Cappello F, Lauricella M. The histone deacetylase inhibitor SAHA induces HSP60 nitration and its extracellular release by exosomal vesicles in human lung-derived carcinoma cells. *Oncotarget*. 2015 Dec 19. doi: 10.18632/oncotarget.6680.
- [2] Rodolico V, Tomasello G, Zerilli M, Martorana A, Pitruzzella A, Gammazza AM, David S, Zummo G, Damiani P, Accomando S, Conway de Macario E, Macario AJ, Cappello F. Hsp60 and Hsp10 increase in colon mucosa of Crohn's disease and ulcerative colitis. *Cell Stress Chaperones*. 2010 Nov;15(6):877-84. doi: 10.1007/s12192-010-0196-8.

Keywords

Doxorubicin; Hsp60; p53; replicative senescence; post-translational modifications.