



Bronchial responsiveness and airway inflammation in trained subjects

M R Bonsignore, N Scichilone and G Morici

Thorax 2008;63;90-91
doi:10.1136/thx.2007.084855

Updated information and services can be found at:
<http://thorax.bmj.com/cgi/content/full/63/1/90>

These include:

References

This article cites 7 articles, 5 of which can be accessed free at:
<http://thorax.bmj.com/cgi/content/full/63/1/90#BIBL>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Notes

To order reprints of this article go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to *Thorax* go to:
<http://journals.bmj.com/subscriptions/>

relationship between bronchial hyperresponsiveness (BHR) and physical activity. Several hypotheses are invoked to explain this association, including a suggestion that physical activity reduces bronchial inflammation by altering airway physiology.

Their major hypothesis is that obesity reduces physical activity and that it is this reduction in physical activity which causes, in some mysterious way, the increase in BHR observed. Their proposed mechanism—that this lack of exercise is associated with a decrease in deep inspiration—is truly breathtaking.

We suggest a much more obvious explanation, which is supported by the published evidence. In our recent survey reported in *Thorax*² we demonstrated a highly significant association of body mass index with chronic cough. Other associations observed in this study infer that the cough of obesity is reflux in nature. If obesity leads to reflux-related respiratory symptoms, can this form of upper airway reflux cause BHR?

Unfortunately, Shaaban *et al*¹ do not provide us with any information concerning the incidence of classic reflux symptoms in their population. In a study of patients with dyspepsia and endoscopically proven gastroesophageal reflux by Bagnato *et al*,³ over one-third had significant BHR. These subjects had no personal/family history or symptoms suggestive of asthma.

However, about two-fifths of patients in the study by Shaaban *et al* had asthma-like symptoms, defined as wheeze and sedentary breathlessness. We suggest that these patients could still have reflux-related symptoms as one-third of patients with chronic reflux cough, as demonstrated by pH monitoring, complain of exertional wheeze and dyspnoea.⁴

With the rising levels of obesity in the population, the accurate recognition of the aetiology of the associated BHR is vital to avoid the spurious diagnosis of “late onset” asthma. Perhaps reflux asthma would be a better—but, as yet, unproven—term.

N M Haley, A H Morice

Academic Respiratory Unit, Castle Hill Hospital, Hull, UK

Correspondence to: Professor A H Morice, Academic Department of Medicine, Castle Hill Hospital, Cottingham East, Yorkshire HU16 5JQ, UK; a.h.morice@hull.ac.uk

Competing interests: None.

Thorax 2008;**63**:89–90. doi:10.1136/thx.2007.084350

REFERENCES

1. Shaaban R, Leynaert B, Soussan D, *et al*. Physical activity and bronchial hyperresponsiveness: ECRHS II. *Thorax* 2007;**62**:403–10.
2. Ford AC, Forman D, Moayyedi P, *et al*. Cough in the community: a cross sectional survey and the relationship to gastrointestinal symptoms. *Thorax* 2006;**61**:975–9.
3. Bagnato GF, Gulli S, Giacobbe O, *et al*. Bronchial hyperresponsiveness in subjects with gastroesophageal reflux. *Respiration* 2000;**67**:507–9.
4. Everett CF, Morice AH. Clinical history in gastroesophageal cough. *Respir Med* 2007;**101**:345–8.

Bronchial responsiveness and airway inflammation in trained subjects

We read with interest the paper by Shaaban and coworkers¹ on the protective effect of physical activity against bronchial hyperreactivity (BHR) in the general population. The authors suggest that a beneficial effect of deep inspirations during exercise could account for the lower prevalence of BHR in physically active subjects compared with sedentary subjects, while the accompanying editorial² favours an “anti-inflammatory” effect of exercise as the most plausible explanation.

We have studied lung function and airway cell biology in non-asthmatic amateur athletes^{3,4} and found that both modulation of airway responsiveness and downregulation of airway inflammation occur with training. At rest, the response to single-dose methacholine inhalation in the absence of deep breaths was significantly lower in amateur runners than in age-matched sedentary controls.³ Shortly after a marathon race the response to methacholine was further blunted, suggesting a causal relationship between endurance exercise and low bronchial responsiveness,³ possibly mediated by ventilation at increased lung volumes.

We have previously reported large numbers of neutrophils in induced sputum of runners.⁴ However, this finding was not associated with evidence of neutrophil activation after intense exercise, since expression of adhesion molecules by airway neutrophils decreased and the elastase concentration in sputum supernatants was unchanged after a marathon race compared with baseline.⁴ Similarly, inflammatory cell infiltration in the airways was not associated with activation of the NFκB pathway in endurance-trained mice,⁵ while airway inflammation was found to decrease strikingly in ovalbumin-sensitised trained mice compared with sedentary mice.⁶ Exercise therefore appears as a model of tightly regulated airway inflammation, possibly secondary to exercise-induced mild bronchial epithelial damage.⁵ Along the same line, physically active smokers appear to be protected against lung function decline and the risk of developing chronic obstructive pulmonary disease compared with sedentary smokers,⁷ supporting a role for regular exercise in blunting airway inflammation.

We acknowledge that athletes, even at the amateur level, do not represent the general population. On the other hand, a publication bias may have favoured preferential reporting of exercise-associated BHR in athletes, especially those training under extreme environmental conditions (such as “ski asthma”) or exposed to irritants (such as swimmers). It is time to reconsider the beneficial effects of regular exercise as a strategy to preserve respiratory health.

Studies like that by Shaaban and coworkers will certainly help us to move in this direction.

M R Bonsignore,^{1,2} N Scichilone,¹ G Morici^{2,3}

¹ Department of Medicine, Pneumology, Physiology and Nutrition (DIMPEFINU), University of Palermo, Italy;

² Institute of Biomedicine and Molecular Immunology (IBIM), National Research Council (CNR), Palermo, Italy;

³ Department of Experimental Medicine, (DIMES), University of Palermo, Italy

Correspondence to: Dr M R Bonsignore, Department of Medicine, Pneumology, Physiology and Nutrition (DIMPEFINU), University of Palermo, Palermo 90100, Italy; marisa@ibim.cnr.it

Competing interests: None.

Thorax 2008;**63**:90. doi:10.1136/thx.2007.084855

REFERENCES

1. Shaaban R, Leynaert B, Soussan D, *et al*. Physical activity and bronchial hyperresponsiveness: ECRHS II. *Thorax* 2007;**62**:403–10.
2. Mahler DA. Is physical activity anti-inflammatory on the airways? *Thorax* 2007;**62**:376.
3. Scichilone N, Morici G, Marchese R, *et al*. Reduced airway responsiveness in non-elite runners. *Med Sci Sports Exerc* 2005;**37**:2019–25.
4. Bonsignore MR, Morici G, Riccobono L, *et al*. Airway inflammation in nonasthmatic amateur runners. *Am J Physiol* 2001;**281**:L668–76.
5. Chimenti L, Morici G, Paternò A, *et al*. Endurance training under standard laboratory conditions damages small airway epithelium in mice. *Am J Respir Crit Care Med* 2007;**175**:442–9.
6. Pastva A, Estell K, Schoeb TR, *et al*. Aerobic exercise attenuates airway inflammatory responses in a mouse model of atopic asthma. *J Immunol* 2004;**172**:4520–6.
7. Garcia-Aymerich J, Lange P, Benet M, *et al*. Regular physical activity modifies smoking-related lung function decline and reduces risk of chronic obstructive pulmonary disease: a population-based cohort study. *Am J Respir Crit Care Med* 2007;**175**:458–63.

Propionibacterium acnes in granulomas of a patient with necrotising sarcoid granulomatosis

Necrotising sarcoid granulomatosis (NSG) was first described by Liebow¹ in 1973. It is defined by three pathological features: the presence of a conglomerate mass of sarcoid-like granulomas; varying degrees of necrosis within the confluent granulomas; and vasculitis with granulomas and giant cells involving the walls of muscular arteries and veins. The relationship between NSG and classic sarcoidosis is controversial. In NSG hilar lymphadenopathy is not seen as frequently as in sarcoidosis, extrapulmonary involvement is rare and serum levels of angiotensin-converting enzyme (ACE) are not necessarily raised.²

The cause of sarcoidosis is unknown, but it has been hypothesised that it results from exposure of a genetically susceptible individual to specific environmental agents. Abe *et al*³ isolated *Propionibacterium acnes* (*P acnes*) in culture from sarcoidosis biopsy specimens, and recently the *P acnes* genome has been detected in sarcoid lymph nodes by