

## PSEUDOANEURYSM OF SUPERIOR THYROID ARTERY FOLLOWING A TRANSESOPHAGEAL ECHOCARDIOGRAPHY: A CASE PRESENTATION

Frank Rikki Canevari<sup>1</sup>, Francesco Martines<sup>2</sup>, Raffaele Sorrentino<sup>1</sup>, Matteo Nicolotti<sup>1</sup>, Federico Sireci<sup>2</sup>

<sup>1</sup> Otorinolaryngology Section, SS Antonio Biagio e Cesare Arrigo Hospital, Alessandria, Italy

<sup>2</sup> Otorhinolaryngology Section, Department of Experimental Biomedicine and Clinical Neurosciences, (BioNeC), University of Palermo, Italy

### ARTICLE INFO

#### Article history:

Received 23 October 2016

Revised 23 December 2016

Accepted 10 January 2017

#### Keywords:

Pseudoaneurysm, ENT complication,

Endovascular coil embolization

### ABSTRACT

The pseudo-aneurysm is an encapsulated hematoma of post-traumatic origin which is in communication with the lumen of the artery of relevance. We present a rare case of pseudo-aneurysm occurring after superior thyroid trans-esophageal echocardiography (TEE) and external cardioversion. Singular occurrence in otolaryngology, if not recognized early a pseudo-aneurysm can result in dramatic events such cataclysmic bleeding or acute occlusion of the upper airway .

© EuroMediterranean Biomedical Journal 2017

## 1. Introduction

Pseudoaneurysm is a pulsating hematoma that results from a tangential injury to an arterial vessel wall and could represent a complication after trauma. In the head and neck region pseudoaneurysm is rare but can have catastrophic consequences; therefore, must be identified speedily in order to prevent events such as hemorrhage or acute occlusion of the airway. A pseudoaneurysm arising from superior thyroid artery (STA) has been reported in only three cases in the literature, one after ultrasonographically guided chemical parathyroidectomy, one after radiotherapy for hypopharyngeal cancer and the last after fine needle aspiration biopsy (FNAC) of thyroid nodule. We report a case of STA after transesophageal echocardiography guided cardioversion.

## 2. Case presentation

A 62-year-old man was admitted to our Otolaryngology Department presenting sore throat, dysphagia and mild dyspnea after a transesophageal echocardiography guided cardioversion performed 3 days earlier. The endoscopic examination of upper aerodigestive tract showed a


diffuse soft edema of arytenoids and pharyngolaryngeal tract without airway obstruction. A corticosteroid therapy was started and the symptoms quickly improved, but two days later, the patient suddenly presented a haemorrhage from the superior aerodigestive tract. The endoscopic examination revealed a swelling of the left lateral wall of the hypopharynx and a haemorrhage from the apex of the swelling which stopped spontaneously after a few minutes. A CT scan of the neck revealed active arterial bleeding with pseudoaneurysmal dilatation of 1,2 cm in diameter arising from a fine branch of external carotid artery and a well-defined homogeneously enhancing mass, 9,5 x 3,2 cm diameter in left neck spaces extending superiorly to parapharyngeal spaces, inferiorly to hypopharynx displacing hyoid bone, thyroid cartilage, posteriorly to cricoid cartilage displacing cervical esophagus and laterally to subcutaneous tissue displacing sternocleidomastoid muscle (Figure 1). The mass was suggestive of hematoma. The angiography revealed a pseudoaneurysm with active bleeding of the terminal tract of the superior thyroid artery (STA) (Figure 2). An endovascular procedure was performed with a superselective microcatheter and an endovascular occlusion by coil embolization of STA with exclusion of the pseudoaneurysm demonstrated at the end of the procedure (Figure 3). The symptoms resolved one day from the endovascular procedure.

\* Corresponding author: Federico Sireci, federicosireci@hotmail.it


DOI: 10.3269/1970-5492.2017.12.3

All rights reserved. ISSN: 2279-7165 - Available on-line at www.embj.org

The patient was discharged after two days without evidence of bleeding and an endoscopic control after 4 weeks revealed that swelling had disappeared.



**Figure 1 - Axial CT scan illustrating pseudoaneurysmal dilatation arising from a fine branch of external carotid artery**



**Figure 2 - Angiography illustrating the pseudoaneurysm with active bleeding of the terminal tract of superior thyroid artery (STA)**



**Figure 3 - Angiography illustrating endovascular occlusion by coil embolization**

### 3. Discussion

Pseudoaneurysm, also called “false aneurysm”, is a hematoma contained by adventitia or perivascular tissue communicating with an arterial vessel that results from a tangential injury to the vessel (1,2). Most often it is secondary to blunt and penetrating trauma (1,3,4) or surgery (5-7) and is a well-known iatrogenic complication of vascular catheterization (8,9) or percutaneous biopsy of solid organs (10).

Pseudoaneurysm of head and neck are quite rare events but could lead to catastrophic complications. Clinical manifestations can include pulsating mass in the neck, dysphagia, dyspnea, pain in the neck, bleeding from oral cavity and cerebrovascular symptoms. The differential diagnosis of an expansive cervical mass include branchial cleft cyst, neck abscess and cervical adenitis. Other cervical masses such as tumors, lymphadenopathy or neurinomas can be excluded by timing the growth of the mass. Many imaging techniques are available to differentiate the diagnosis. Color Doppler ultrasound is an excellent technique to detect the presence of an aneurismal sac and the presence of active bleeding. MRI and CT scan of the neck give information regarding the involvement of neck anatomical structures. Angiography is the best technique to investigate the morphology of the vessel involved and the source of the aneurismal sac.

Management options for pseudoaneurysm include conservative measures with or without sonographically guided compression therapy (10,12), surgical repair (13), transarterial coil embolization (14) and percutaneous thrombin or alcohol injection (15). There is no standard treatment protocol because of the rarity of neck pseudoaneurysm and therefore, appropriate treatment is determined on a case-by-case basis.

The rapid development of an expanding mass under the angle of the mandible or in lateral pharyngeal wall after surgical procedure of the neck, percutaneous biopsy and neck trauma should always raise the suspicion of an extracranial arterial pseudoaneurysm.

To our knowledge only three cases of pseudoaneurysm of superior thyroid artery have been reported previously. The first case was a pseudoaneurysm occurring after ultrasonographically guided chemical parathyroidectomy (16). The diagnosis was made by angiography and treated by selective coil embolization. The second case reported a pseudoaneurysm in a patient with hypopharyngeal squamous cell carcinoma during simultaneous radiotherapy and chemotherapy (14). The diagnosis was made by a CT scan and treated using selective coil embolization. The third case reported a pseudoaneurysm after ultrasonographically guided biopsy of a thyroid nodule (10). The diagnosis was made after ultrasonographic and color Doppler examination and treated waiting spontaneous thrombosis.

In our patient the pseudoaneurysm occurred after a transesophageal echocardiography guided cardioversion (TEE-guided). The diagnosis was made by a computed tomography scan and treated by selective coil embolization. Transesophageal echocardiography guided cardioversion with short-term anticoagulation can be considered safe and clinically effective for patients with atrial fibrillation. Complications reported after TEE-guided procedure include bleeding events. We describe a unique case of superior thyroid artery pseudoaneurysm after TEE-guided.

Pseudoaneurysm in the neck is quite a rare event but a rapidly growing expanding mass in the neck after surgical procedure of the neck, percutaneous biopsy and neck trauma or transoral procedures should

always raise the suspicion of an extracranial arterial pseudoaneurysm (17). A rapidly growing mass in the neck can lead to an acute airway obstruction and quick diagnosis and treatment is required in order to prevent catastrophic consequences. The literature so far details no protocols for diagnosis and treatment. In our opinion CT scan for differential diagnosis and visualization of anatomical structures involved and a transarterial coil embolization can be considered safe and effective diagnostic and treatment procedures.

## References

1. Krempf GA, Noorily AD. Pseudoaneurysm of the descending palatine artery presenting as epistaxis. *Otolaryngol Head Neck Surg.* 1996 Mar;114(3):453-6
2. Reiber ME, Burkey BB. Intracavernous carotid pseudoaneurysm after blunt trauma: case report and discussion. *Head Neck.* 1994 May-Jun;16(3):253-8.
3. Canevari FR, Giourgos G, Pistochini A. The endoscopic transnasal parasseptal approach to a sphenoid sinus osteoma: case report and literature review. *Ear Nose Throat J.* 2013 Dec;92(12):E7-E10. Review. Erratum in: *Ear Nose Throat J.* 2014 Apr-May;93(4-5):148.
4. Han MH, Sung MW, Chang KH, Min YG, Han DH, Han MC. Traumatic pseudoaneurysm of the intracavernous ICA presenting with massive epistaxis: imaging diagnosis and endovascular treatment. *Laryngoscope.* 1994 Mar;104(3 Pt 1):370-7.
5. Smelt JL, Alhamarneh O, Dyer JD, Liew L. Endovascular coiling of a mycotic external carotid artery pseudoaneurysm following pharyngolaryngectomy with a free jejunal graft. *J Laryngol Otol.* 2012 Feb;126(2):214-6.
6. Karas DE, Sawin RS, Sie KC. Pseudoaneurysm of the external carotid artery after tonsillectomy. A rare complication. *Arch Otolaryngol Head Neck Surg.* 1997 Mar;123(3):345-7.
7. Cockroft KM, Carew JF, Trost D, Fraser RA. Delayed epistaxis resulting from external carotid artery injury requiring embolization: a rare complication of transsphenoidal surgery: case report. *Neurosurgery.* 2000 Jul;47(1):236-9.
8. Kua JS, Tan IK. Airway obstruction following internal jugular vein cannulation. *Anaesthesia.* 1997 Aug;52(8):776-80.
9. Parry W, Dhillon R, Salahudeen A. Carotid pseudoaneurysm from inadvertent carotid artery catheterization for haemodialysis. *Nephrol Dial Transplant.* 1996 Sep;11(9):1853-5.
10. Celik H, Yücel C, Oktar S, Karadag Z, Ozdemir H. Iatrogenic pseudoaneurysm of the superior thyroid artery: color Doppler ultrasonographic diagnosis and treatment approach. *J Ultrasound Med.* 2004 Dec;23(12):1675-8.
11. Attard A, Geraci G, Santoro V, Modica G, Attard M, Ferrauto S, Speciale R, Marchese D. Retropharyngeal abscess from fishbone in adult immunocompetent host presenting as acute thyroiditis. *EuroMediterranean Biomedical Journal* 2015, 10(18): 203-208
12. Schaub F, Theiss W, Heinz M, Zigel M, Schömig A. New aspects in ultrasound-guided compression repair of postcatheterization femoral artery injuries. *Circulation.* 1994 Oct;90(4):1861-5.
13. Walker AT, Chaloupka JC, Putman CM, Abrahams JJ, Ross DA. Sentinel transoral hemorrhage from a pseudoaneurysm of the internal maxillary artery: a complication of CT-guided biopsy of the masticator space. *AJNR Am J Neuroradiol.* 1996 Feb;17(2):377-81.
14. Ernemann U, Herrmann C, Plontke S, Schäfer J, Plasswilm L, Skalej M. Pseudoaneurysm of the superior thyroid artery following radiotherapy for hypopharyngeal cancer. *Ann Otol Rhinol Laryngol.* 2003 Feb;112(2):188-90. Mann GS, Heran MK. Percutaneous thrombin injection of a superficial temporal artery pseudoaneurysm. *Pediatr Radiol.* 2007; 37:578-80
15. Perona F, Barile A, Oliveri M, Quadri P, Ferro C. Superior thyroid artery lesion after US-guided chemical parathyroidectomy: angiographic diagnosis and treatment by embolization. *Cardiovasc Intervent Radiol.* 1999 May-Jun;22(3):249-50.
16. Ferreli F, Turri-Zanoni M, Canevari FR, Battaglia P, Bignami M, Castelnuovo P, Locatelli D. Endoscopic endonasal management of non-functioning pituitary adenomas with cavernous sinus invasion: a 10-year experience. *Rhinology.* 2015 Dec;53(4):308-16.
17. Martines F, Salvago P, Costanzo R, Di Marzo M, Ferrara S, Iovane A, Messina G, Mucia M, Mulè A, Palma S, Rizzo S, Sireci F. Extramedullary plasmocytoma of the tonsil: a new management. *Acta Medica Mediterranea.* 2015 Oct; 31: 431-434